

Iron biogeochemistry across marine systems – progress from the past decade

E. Breitbarth^{1,2}, E. P. Achterberg³, M. V. Ardelan⁴, A. R. Baker⁵, E. Bucciarelli^{6,7}, F. Chever^{6,7}, P. L. Croot⁸, S. Duggen⁹, M. Gledhill³, M. Hassellöv², C. Hassler¹⁰, L. J. Hoffmann^{1,11}, K. A. Hunter¹, D. A. Hutchins¹², J. Ingri¹³, T. Jickells⁵, M. C. Lohan¹⁴, M. C. Nielsdóttir³, G. Sarthou^{6,7}, V. Schoemann¹⁵, J. M. Trapp¹⁶, D. R. Turner², and Y. Ye¹⁷

¹Department of Chemistry, University of Otago, Dunedin, New Zealand

²Department of Chemistry, University of Gothenburg, Gothenburg, Sweden

³National Oceanography Center Southampton, University of Southampton, Southampton, UK

⁴Norwegian University of Science and Technology, Department of Chemistry, Trondheim, Norway

⁵School of Environmental Sciences, University of East Anglia, Norwich, UK

⁶Université Européenne de Bretagne, France

⁷Université de Brest, CNRS, IRD, UMR 6539 LEMAR, IUEM, Plouzané, France

⁸IFM-GEOMAR, Leibniz-Institute of Marine Sciences, Division Marine Biogeochemistry, Kiel Germany

⁹IFM-GEOMAR, Leibniz-Institute of Marine Sciences, Division Dynamics of the Ocean Floor, Kiel, Germany

¹⁰Centre for Australian Weather and Climate Research (CAWCR), Hobart, TAS, Australia

¹¹Department of Plant and Environmental Sciences, University of Gothenburg, Gothenburg, Sweden

¹²Department of Biological Sciences, University of Southern California, Los Angeles, CA, USA

¹³Luleå University of Technology, Division of Applied Geology, Luleå, Sweden

¹⁴Marine Institute, University of Plymouth, Plymouth, UK

¹⁵Ecologie des Systèmes Aquatiques, Université Libre de Bruxelles, Bruxelles, Belgium

¹⁶University of Miami, Rosenstiel School of Marine and Atmospheric Science, Department of Marine and Atmospheric Chemistry, Miami, USA

¹⁷Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

Received: 31 May 2009 – Published in Biogeosciences Discuss.: 7 July 2009

Revised: 20 January 2010 – Accepted: 31 January 2010 – Published: 19 March 2010

Abstract. Based on an international workshop (Gothenburg, 14–16 May 2008), this review article aims to combine interdisciplinary knowledge from coastal and open ocean research on iron biogeochemistry. The major scientific findings of the past decade are structured into sections on natural and artificial iron fertilization, iron inputs into coastal and estuarine systems, colloidal iron and organic matter, and biological processes. Potential effects of global climate change, particularly ocean acidification, on iron biogeochemistry are discussed. The findings are synthesized into recommendations for future research areas.

1 Preface

An international workshop addressing the biogeochemistry of iron in the context of global change across marine ecosystems was held in Gothenburg, Sweden (14–16 May 2008). Largely driven by geographic separation, iron biogeochemistry in the open ocean and in coastal seas are often addressed as two distinct fields and the workshops organized over the past two decades have normally either been system- or task-specific. This has led to the development of system-specific expertise and research approaches, with potential separation of know-how. The aim of this workshop was to conduct a broader cross-system review of marine iron biogeochemistry by bringing together scientists from a wide range of coastal, shelf and deep-ocean environments to merge their system-specific knowledge into a truly cross-disciplinary and cross-system synthesis. This lead article is an attempt to



Correspondence to: E. Breitbarth
(ebreitbart@chemistry.otago.ac.nz)

summarize the scientific milestones of the past 10 years discussed during the workshop.

The Gothenburg workshop was convened almost ten years after a workshop meeting in Amsterdam, sponsored by SCOR and IUPAC which formed the basis for the book “The Biogeochemistry of Iron in Seawater” (Turner and Hunter, 2001). The Gothenburg workshop revisited the topics listed in the “Summary and Recommendations” of this book and took up two further cross-cutting aspects: (A) What can we learn from comparing Fe biogeochemistry in coastal and open ocean systems? And (B), how are global change processes expected to affect Fe biogeochemistry?

This article aims to synthesize the cross-system and interdisciplinary knowledge from atmospheric, chemical, biological, and geological angles discussed during the Gothenburg workshop and ties the manuscripts of the special issue “Iron biogeochemistry across marine systems at changing times” into this overall context. Due to this wide range of topics, it is not intended to be a comprehensive, in-depth review on all aspects of marine iron biogeochemistry. We follow the structure of the workshop topics, which were: Natural Fe fertilization (Sect. 2, articles: Ardelan et al., 2010; Chever et al., 2010; Duggen et al., 2010; Ye et al., 2009); artificial Fe fertilization (3: Bucciarelli et al., 2010; Chever et al., 2010); Fe inputs into coastal and estuarine systems (4: Gelting et al., 2009; Breitbarth et al., 2009); Colloidal iron and organic matter (5); Linking biological processes to iron chemistry (6: Breitbarth et al., 2009; Bucciarelli et al., 2010; Hassler and Schoemann, 2009; Steigenberger et al., 2010); and Iron and Climate Change (7: Breitbarth et al., 2010; Rose et al., 2009). Each section concludes with recommendations for future research.

2 Natural iron fertilization

The past decade brought major advances in the understanding of natural iron fertilization processes to the open ocean. The field is generally subdivided into two major areas: atmospheric deposition with the main focus on dust deposition from the continents and more recently addressing volcanic ash and pumice depositions; and marine processes, where particular areas of interest have been ice melting, hydrothermal vents, continental margins, and the island mass effects.

2.1 Atmospheric deposition – dust

Over the last 10 years, the importance of dust transport and deposition within the Earth System has become clear (Jickells et al., 2005). This includes the role of dust in transporting iron to the oceans, but also the transport of nutrients to land and impacts on albedo. Dust supply is episodic and predominantly from desert regions, and satellite advances have allowed these sources to be better characterized (Prospero et al., 2002). These satellite advances also allow some im-

provement in understanding of dust transport and deposition, but this is still limited to high dust regions where the total aerosol is dominated by dust (Mahowald et al., 2005). In regions remote from the desert sources, aerosols may be dominated by sea and acid salts. Furthermore close to a source region, particularly over the ocean off North Africa, the dust is transported at altitude, so the satellite detection of a dust plume, does not necessarily imply deposition to the oceans at that location (Mahowald et al., 2005). Since dust transport is episodic, field data to validate models and provide direct estimates of dust loading should ideally cover periods of months to years. Obviously though, shorter campaign style measurements can be useful for studying processes, and if repeated can provide long term average concentrations. The number of long-term dust monitoring stations is still very limited and broadly the same as identified in Jickells and Spokes (2001). This data set is dominated by the Prospero network (e.g. Ginoux et al., 2004), and the lack of data in the low dust regions, where ocean euphotic zone iron limitation is evident, is notable. Recent campaigns in some of these regions (Baker et al., 2006; Planquette et al., 2007; Wagener et al., 2008) do provide some confidence in the dust transport models, but the uncertainties in parameterizations within the models are still considerable and hence the uncertainties in flux estimates are still substantial. The work of Measures and colleagues (e.g. Han et al., 2008) has demonstrated the validity of a novel indirect approach of using surface water Al as a tracer of atmospheric deposition which provides data averaged over long time scales (months to years) in remote regions. Again this approach has significant uncertainties, but the broad agreement between this, long term field data, campaign data and models provides reassurance that the estimates of total dust deposition to the oceans and the regional patterns are realistic.

A major continuing source of uncertainty in estimating dust deposition to the oceans is associated with the parameterisation of wet and dry deposition, except in the few cases where wet deposition has been measured directly. The congruence of data and models noted above does provide some confidence that, at the global scale, the average deposition parameterization is approximately correct. This does not mean that the resultant dust flux from these averages is estimated correctly at the regional scale, or in the low dust regions of water column iron limitation. Duce et al. (1991) estimated uncertainties of a factor of three in the deposition flux and this uncertainty largely remains. Jickells et al. (1998) demonstrated that the use of ocean sediment trap data can provide a valuable constraint on the uncertainties in deposition fluxes and Mahowald et al. (2005) considered this further. However, the use of this technique in low dust regions does require high quality measurements of a dust tracer such as Al in the sediment traps and this is not always available. If this became routine it would offer a mechanism to significantly reduce uncertainties in deposition parameterization. Such an improvement would allow dust and iron mass

balance in individual regions and comparison to productivity estimates.

The conversion of dust deposition to soluble iron fluxes requires the solubility of iron from dust to be known. This is required ideally over the timescale of the residence times of dust in the water column (tens of days, see Jickells et al., 2005) and at realistically low dust loadings, although this is very difficult in practice and more pragmatic short term simple aerosol leaching schemes are usually applied (Baker and Croot, 2009).

Considerable effort has been put into studies of aerosol dust solubility over the last 10 years and this has tended to confirm that on a global average Fe solubility is low (Jickells et al., 2005; Mahowald et al., 2005), but also demonstrated that the use of a single solubility estimate is probably inappropriate and there does appear to be a systematic increase in solubility from high to low dust regions (Baker and Croot, 2009; Baker and Jickells, 2006). There is still considerable debate surrounding the drivers of this variation in solubility with four main possibilities; (i) atmospheric chemical processing during dust transport (Fan et al., 2006; Jickells and Spokes, 2001), (ii) systematic changes in aerosol particle size leading to changes in surface area and solubility (Baker and Jickells, 2006), (iii) an additional source of iron beside crustal dust (Jickells et al., 2005; Schroth et al., 2009); (iv) active biological acquisition and uptake mechanisms such as siderophores and grazing that can circumvent abiotic dissolution limitations (Barbeau and Moffett, 2000; Yoshida et al., 2002; Frew et al., 2006).

There is good evidence that solubility of iron from anthropogenic aerosol is higher than from soil dust (Schroth et al., 2009; Journet et al., 2008) but the significance of this high solubility anthropogenic dust to the global iron cycle is uncertain, and in particular it seems unlikely to be responsible for high iron solubilities in aerosols in remote regions seen for instance by Baker and Jickells (2006). However, recent measurements of iron speciation from African dust collected in the Trade Winds at Barbados support the case for anthropogenic iron controls over iron solubility. Trapp et al. (2010) show that Fe^{3+} dominates the iron solubility over the entire range of particle sizes. However, at low mineral dust concentrations Fe^{2+} , believed to be largely derived from anthropogenic sources, becomes increasingly important. Air-mass back trajectories indicate biomass burning in southern Africa and potentially also South America as the source of this anthropogenic iron and dust samples had an $\text{Fe}^{2+}/\text{Fe}^{3+}$ ratio twice that measured in dust-laden aerosols from North Africa. (Trapp et al., 2010).

In their contribution to this special issue, Ye et al. (2009) aim to improve the understanding of the impact of dust deposition on Fe bioavailability and marine primary productivity in modeling iron speciation and biogeochemistry at TENATSO (Tropical Eastern North Atlantic Time-series Observatory). Based on recent studies on Fe speciation and the existing model for the Bermuda Atlantic Time-series Study

(BATS) (Weber et al., 2007), this model aims at studying the role of dust particles in Fe removal and providing a better description of the sources and fate of organic Fe-binding ligands.

Dry deposition probably dominates dust and iron deposition over some regions of the ocean, particularly those where winds flow off the land including areas downwind of major deserts such as the Sahara. Wet deposition is probably particularly important for total deposition in remote regions of the ocean. The data set for dust and soluble iron in wet deposition in the marine atmosphere is very small, and this requires improvement. To date, this discussion has largely considered only dust and iron deposition. However, atmospheric deposition delivers significant amounts of iron (Jickells et al., 2005) and nitrogen (Duce et al., 2008) and relatively small amounts of phosphorus (Mahowald et al., 2008) relative to phytoplankton requirements. Assorted trace metals that may play a role in phytoplankton productivity, including some that are potentially inhibitory, will also be deposited (e.g. Paytan et al., 2009). It is important that we evaluate the impacts of atmospheric deposition holistically, and not artificially separate the contributions of individual nutrients.

2.2 Other atmospheric and marine processes of natural Fe fertilization

All short-term artificial Fe fertilization experiments unequivocally showed the importance of Fe on the carbon cycle, in particular on the food web structure and functioning (e.g. Boyd, 2004; Boyd et al., 2007, 2000; Coale et al., 1996, 2004; Gervais et al., 2002; Tsuda et al., 2003; de Baar et al., 2005, see Sect. 2). However, it is difficult to reliably assess the magnitude of carbon export to the ocean interior using such methods (Blain et al., 2007). Recent natural Fe fertilization experiments carried out in the Southern Ocean showed that the efficiency of fertilization was at least 10 to 20 times greater than that of a phytoplankton bloom induced artificially by adding iron, (KEOPS and CROZEX, Blain et al., 2007; Pollard et al., 2009). Large losses of purposefully added iron can explain the lower efficiency of the induced bloom, as well as the mode of iron addition and the requirement of concomitant supply with major nutrients (Pollard et al., 2009; Blain et al., 2007). In the open ocean, a large variety of naturally iron-fertilized sites exist, which could allow for improved forecasting of the oceanic response to Fe fertilization and a better knowledge of Fe sources to the open ocean. Chever et al. (2010) provide a Fe budget for the naturally fertilized area above the Kerguelen Plateau, using total dissolvable Fe as an additional tracer to better constrain the Fe cycle in this area. They show that horizontal advection of water from South of the Plateau seems to be the predominant source of apparent particulate and dissolved iron above the plateau, over atmospheric and vertical inputs. Further, Ardelan et al. (2010) illustrate natural Fe enrichment processes from the South Shetland Islands-Antarctic Peninsula region.

As discussed in Sect. 2.1, aeolian inputs may have different origins, such as (i) the arid and semi-arid regions (Jickells et al., 2005), (ii) combustion sources (fossil fuel burning, incinerator use, biomass burning; (Spokes and Jickells, 2002; Guieu et al., 2005; Sedwick et al., 2007; Luo et al., 2008), but also by (iii) meteoritic material and extraterrestrial dust (Johnson, 2001), and (iv) volcanic origin (Benitez-Nelson et al., 2003; Duggen et al., 2007; Boyd et al., 1998). All atmospheric input may have an effect on biological productivity in the ocean (Schroth et al., 2009), in particular on bacterial activity (Pulido-Villena et al., 2008), although the causative link is not always obvious as shown by (Boyd et al., 2009).

While the meteoritic contribution is difficult to assess due to the sporadic events, the amount of soluble (presumably bioavailable) iron input into the ocean from extraterrestrial dust is estimated to be $7 \times 10^9 \text{ g year}^{-1}$ (Johnson, 2001) and is thus not insignificant. More so, volcanic eruptions can transport volcanic ash up to several tens of kilometres high into the atmosphere and fine ash may encircle the globe for years, thereby reaching even the remotest and most iron-starved oceanic areas (Schmincke, 2004). The implication of volcanism for the marine biogeochemical iron-cycle is poorly constrained so far. Recent studies demonstrate that volcanic ash from volcanoes worldwide quickly releases soluble and bio-available iron on contact with water (e.g. Jones and Gislason, 2008; Duggen et al., 2007; Frogner et al., 2001). Drill core data from scientific ocean drilling show that volcanic ash layers and dispersed ash particles are frequently found in marine sediments and that volcanic ash deposition and therefore iron-injection into the oceans took place throughout much of the Earth's history (Straub and Schmincke, 1998). It may thus well be possible that the contribution of volcanic ash to the marine biogeochemical iron-cycle is generally underestimated. A review paper (Duggen et al., 2010) summarises the development and the knowledge in a fairly young research field covering a wide range of chemical and biological issues and gives recommendations for future directions. The approach by Duggen et al. (2010) contributes to understanding of the role of volcanic ash for the marine biogeochemical iron-cycle, marine primary productivity and the ocean-atmosphere exchange of CO_2 and other gases relevant for climate throughout the Earth's history.

Melting of sea ice, icebergs and glacial inputs may contribute as Fe sources in polar regions. Estimates of these sources' magnitudes are poorly constrained. Recent studies have highlighted the importance of these sources (Lannuzel et al., 2008; Lannuzel et al., 2007; Statham et al., 2008; Aguilar-Islas et al., 2008; Raiswell et al., 2008, 2006; Smith et al., 2007; Croot et al., 2004). Iron accumulates in sea ice with concentrations one to two orders of magnitude higher than the underlying seawater. Atmospheric iron can be one source but flux estimates by Lannuzel et al. (2008, 2007) seem to indicate that iron must come mostly from below. The exact mechanism remains unclear, but recent evidence

suggests that organic matter could play an essential role in trapping Fe in the sea ice not only during sea ice formation but also during ice algae proliferation in the bottom ice after its formation (Schoemann et al., 2008). Its release into the seawater during ice melting can occur in short time spans such as weeks. For example, Lannuzel et al. (2008) showed that 70% of the accumulated Fe in the sea ice could be released due to brine drainage in a 10 days period, while the sea ice cover was still present. This represents a significant Fe flux to the surface ocean that may be instrumental in sustaining springtime ice edge blooms in the marginal ice zone and polynyas. Dense phytoplankton blooms have been observed in combination with the receding ice edge or in coastal shelf areas (e.g. Smith and Nelson, 1985; Holm-Hansen et al., 1989). Moreover, both sea ice and icebergs may constitute vectors of Fe transport far away from its initial source (Smith et al., 2007; Lancelot et al., 2009). Further to this, Edwards and Sedwick (2001) addressed the contribution of snow bound aerosol iron in the Antarctic seasonal sea ice zone.

The continental margins may also play a key role as a Fe source (Elrod et al., 2004; Laës et al., 2003, 2007; Chase et al., 2005; Blain et al., 2008). As an example, Lam and Bishop (2008) clearly showed that the continental margin was a key source of Fe to the HNLC (high nutrient low chlorophyll) North Pacific Ocean, since the lateral source of Fe is shallow enough to be accessible to phytoplankton by winter mixing and Fe can be transported at distance over 900 km from the continental shelf.

Our current challenge in regions where natural iron fertilization occurs is to have a better knowledge and quantification of these various Fe sources. For example, the global atmospheric iron fluxes are reasonably well known, but the fluxes to remote low iron regions are rather uncertain. Moreover, the aerosol iron solubility varies systematically, but the underlying causes of this are uncertain. In the deep waters, Fe can be transported far away from the source, especially in waters with anoxic conditions (Blain et al., 2008). Local and remote sources of Fe may not have the same impact on carbon cycle. We also need to understand how (i) the different sources of Fe influence its speciation and bioavailability; (ii) they contribute to the global Fe budget; (iii) they will be affected by global change, and (iv) what are the physical mechanisms that allow long distance Fe transport: advection (strong currents, ACC, EUC, de Baar et al., 1995; Mackey et al., 2002; Lam and Bishop, 2008; Loscher et al., 1997), internal waves and slope circulation (Laës et al., 2003), and eddies (Johnson et al., 2005). Finally, the physical mechanisms that allow Fe to be accessible for the food web should also be better understood and quantified (upwelling, diapycnal mixing, winter mixing, Blain et al., 2007).

To assess these challenging issues, there is a crucial need for (i) multi-disciplinary studies (physics/biogeochemistry/biology), (ii) multi-proxy approaches, such as the one promoted by the international

GEOTRACES program, including oceanic sections and intercalibration experiments for seawater and aerosols; (iii) the development of biogeochemical models that correctly take into account the various Fe sources and their impact on Fe speciation and bioavailability, and (iv) the development of regional iron budgets.

3 Artificial iron fertilization

The largest source of iron for the HNLC surface waters comes from deep water supply (Watson, 2001). However, the Fe:N or Fe:P ratio of the upwelled deep waters is often not high enough for optimum phytoplankton growth (Moore et al., 2006). Consequently, an additional source of iron is required, which could be derived from suboxic or anoxic sediments (Las et al., 2007) or dust inputs (Jickells et al., 2005). Fertilization of the Southern Ocean with dust has been suggested as an explanation for past glacial periods (Martin, 1990). During these periods iron dust inputs to the oceans were strongly enhanced, with the Southern Ocean receiving up to 10 times more dust-derived iron (Wolff et al., 2006), and consequent stimulation of phytoplankton growth and the biological carbon pump. Nevertheless, it has been estimated that the increase in iron stimulated productivity could have contributed perhaps 15–25% of the 80–100 ppm drawdown in atmospheric CO₂ observed during glacial maxima by enhancing the biological carbon pump (Sigman and Boyle, 2000; Bopp et al., 2003).

When trace metal clean techniques became available in the late 1980s it was possible to directly test the effect of iron additions on phytoplankton growth in HNLC regions. Ship-board iron-addition bottle experiments clearly showed that these additions stimulated phytoplankton growth (e.g. de Baar et al., 1990). However, the potential for bottle-effects during these experiments led researchers to plan and undertake mesoscale Lagrangian-type oceanic experiments to study the influence of iron additions on primary productivity, and investigate the consequences for nutrient utilization, ecosystem dynamics and carbon export. More than a dozen of these large scale (typically 10×10 km grid) iron addition experiments have been conducted to date in HNLC regions and were reviewed by de Baar et al. (2005) and Boyd et al. (2007). Recent experiments involved conducting iron, carbon, nutrient, climatically active gasses, and ecosystem observations, the latest being Lohafex (January–March, 2009; Editorial Nature Geoscience, Editorial, 2009) following a 300 km² iron addition in a stable Southern Ocean mesoscale eddy for >7 weeks. The longer time scale allowed a thorough examination of biogeochemical and ecosystem changes and carbon export.

In summary, all artificial iron experiments have confirmed that iron supply limits primary production and has impact on phytoplankton species composition and bloom dynamics in tropical as well as in polar HNLC waters (Boyd et al., 2007;

de Baar et al., 2005). Iron limitation also induces a decoupling in the use of macronutrients by phytoplankton, likely to influence the cycling of the major biogeochemical cycles (C, N, P, Si, S) over geological time scales (de Baar and La Roche, 2003). In addition to iron, light has been shown to play an important role in the regulation of phytoplankton production in HNLC regions (Moore et al., 2007a, b; Maldonado et al., 1999; Boyd et al., 2001; Hoffmann et al., 2008; de Baar et al., 2005; Bucciarelli et al., 2010). Overall, in situ iron fertilization experiments have greatly enhanced our knowledge about iron solubility, organic iron complexation, and the importance of iron redox states (e.g. Rue and Bruland, 1995; Croot et al., 2001; Rue and Bruland, 1997), which apply to Fe biogeochemistry in the ocean in general.

Large scale iron oceanic addition has been suggested as an option for mitigating the present day increasing atmospheric CO₂ concentrations (Kintisch, 2007). The Southern Ocean is the HNLC region where iron stimulation of CO₂ sequestration would be most efficient and yield long-term carbon storage in deeper waters (Sarmiento and Orr, 1991). Currently there are a number of uncertainties surrounding intentional, large-scale, ocean iron fertilization, which will require further research for clarification. These have already been critically assessed by Chisholm et al. (2001) as well as more recently by Buesseler et al. (2008). Potential side effects include that the mineralization of the enhanced sinking phytoplankton biomass could result in local anoxia and consequent negative effects to oceanic ecosystems and the production of the harmful greenhouse gases nitrous oxide and methane (Cullen and Boyd, 2008; Furman and Capone, 1991). Other climate active gases, like dimethylsulfide (DMS) might increase following Fe fertilisation (Liss et al., 2005). Direct ecosystem shifts resulting in for example proliferation of jellyfish have also been suggested. Furthermore, purposeful iron fertilization may result in a reduced nutrient inventory and consequently reduced productivity and potentially fisheries in oceanic systems downstream of the fertilization areas (Gnanadesikan et al., 2003).

A further key unknown is the efficiency of carbon removal. The amount of carbon sequestered per unit addition of iron is crucial to the effectiveness of iron fertilization (de Baar et al., 2008). The artificial experiments have indicated an efficiency of biological carbon export into deeper water (100–250 m) ranging from 650 (SERIES, Boyd et al., 2004) to 3300 (mol C/mol Fe) (SOFEX – south, Buesseler et al., 2004). The seasonal sequestration efficiencies estimated for natural Fe fertilization are much higher, 8640 for CROZEX (Pollard et al., 2009) and 154 000 for KEOPS (Chever et al., 2010). The discrepancies in effectiveness between natural and purposeful fertilizations might be partly due to the ~75% immediate loss of added Fe in artificial fertilisations (de Baar et al., 2008). These values will need to be much more tightly constrained to allow a thorough assessment of the potential success of iron fertilization as a means to reduce the increasing atmospheric CO₂ concentrations and cost

(Boyd, 2008). The success of the large scale oceanic additions of iron has furthermore been put into doubt by modeling studies. Recent work by Dutkiewicz et al. (2005) and Aumont and Bopp (2006) suggests that large scale iron additions would only reduce atmospheric CO₂ concentrations by ca. 10 ppm, as other limiting factors such as light and zooplankton grazing become more important. It appears that large uncertainties remain with respect to the efficiency of iron fertilization that require further investigations using observation and models. For a recent in depth assessment of the topic see Boyd et al. (2007), as well as Boyd (2008) and associated publications. From a marine trace-metal research perspective, the attendants of the workshop came to the conclusion, that priority should be given to small scale open ocean Fe biogeochemistry studies that are specifically designed to address clearly defined research questions of trace-metal chemistry.

4 Fe inputs into coastal and estuarine systems

The coastal area is a key environment in the global iron cycle, where the brackish water environment changes the physicochemical speciation, and thus mobility, of river-introduced iron via aggregation, sedimentation and redox processes. The coastal waters also are a highly dynamic transition zone, resulting in very diverse temporal and spatial chemical and biological changes. Total concentrations of iron in coastal waters though are generally several orders of magnitude higher than open ocean values and at a first glance, iron limitation of primary production in coastal areas seems not very likely. However, temporal growth limitation by iron can occur in some coastal upwelling regions (Bruland et al., 2001; Hutchins and Bruland, 1998) and fjord systems (Öztürk et al., 2002).

The world's largest estuary, the Baltic Sea, serves as an excellent large scale laboratory to study trace metal chemistry over a wide salinity gradient. Here the total iron concentration decreases by more than an order of magnitude from the low salinity north-east (Bothnian Bay), via the Bothnian Sea to its central part (Baltic Proper), thus forming a natural well defined iron concentration gradient for studying physicochemical speciation of iron and the role of iron for primary production at different total (unfiltered) iron concentrations (Gelting et al., 2009). The authors observed significant variations in the physicochemical speciation, including the iron isotopes, at high temporal resolution from the euphotic zone. Other large river systems such as the Columbia River and Mississippi also show large gradients in iron concentrations but also act as significant sources of Fe to coastal regions (Powell and Wilson-Finelli, 2003; Lohan and Bruland, 2006).

In addition to photochemical processes and organic complexation it is the cycling of iron between particles, colloids and the truly dissolved fraction (<1 kD), rather than the to-

tal concentration, that determines the bioavailability of iron in coastal surface water. The truly dissolved fraction can rapidly be consumed during bloom conditions if this fraction is small and exchange processes between particulate-colloidal matter and the truly dissolved fraction are slow. Hence, knowledge about distribution and cycling of iron between these phases in the coastal zone is fundamental for predictions about iron limitation for plankton growth, and is key to understanding iron export pathways to the open ocean. For the Baltic Sea, Gelting et al. (2009) show that iron in the <1 kDa fraction never reached critical low levels during summer phytoplankton bloom conditions. Further, Fe(II) is generally not considered as an abundant source of bioavailable iron due to its short residence time in oxygenated water. However, a relatively high standing concentration of Fe(II), large enough to cover the demand for iron by cyanobacteria in Baltic Sea waters, was observed by Breitbarth et al. (2009) in a study paralleling Gelting et al. (2009).

Measurements of the physicochemical speciation of iron in freshwater during the last five years suggest that iron transport in rivers is associated with two types of carrier phases (besides detrital particles), an oxyhydroxide phase with associated CDOM (chromophoric dissolved organic matter, mostly consisting of humic acids) and an organic carbon (fulvic) phase (e.g. Lyvén et al., 2003; Andersson et al., 2006). Much of this fulvic phase is present as small colloids and in the truly dissolved fraction (<1 kD). When these phases reach the saline coastal water substantial aggregation of the Fe-oxyhydroxide fraction with associated CDOM is observed, whereas iron associated to the fulvic fraction show little aggregation (Stolpe and Hassellöv, 2007) and survives the sequential sequestration from the water column during gradual mixing with seawater (Krachler et al., 2005). It is possible that this land-derived fraction can reach the open ocean, as indicated by recent data (Laglera and van den Berg, 2009). With fulvic acid as one important carrier mechanism for riverine Fe, the influence of this Fe source reaches further out to sea than previously expected. Tovar-Sanchez et al. (2006) for example, suggested based on metal composition, that riverine and not dust born material was the main source of trace metal accumulation in a diazotroph (*Trichodesmium* sp.) dwelling the surface waters of the subtropical and tropical North Atlantic Ocean.

Back at the river-seawater interface, Gerringa et al. (2007) argue that particularly the weak iron ligand groups (L₂) may impede the precipitation of Fe in the Scheldt Estuary upon mixing with seawater and that the strong ligand (L₁) generally observed in the open ocean, albeit also present, were insufficient in concentration. Powell and Wilson-Finelli (2003) though point out that the latter is of crucial importance for Fe transport in the Mississippi river plume. Likewise, Buck et al. (2007) demonstrated the predominant importance L₁ type ligands for Fe transport into the sea from the Columbia River and San Francisco Bay plumes. The stability constants of these strong L₁ ligands are very similar to those reported by

Laglera and van den Berg (2009) for Fe bound to fulvic acids, indicating the importance of these ligands in controlling the solubility of dissolved iron in riverine and coastal systems. Clearly, Fe speciation in estuarine and near-shore waters can not be addressed in a generalized manner and systems may differ depending on watershed characteristics (e.g., pristine versus anthropogenically impacted) as well as the level and type of riverine input (Öztürk and Bizsel, 2003; Krachler et al., 2005) (see Sect. 5 for colloidal matter).

Ingri et al. (2006) suggest that iron isotopes could be used to roughly identify the two major suspended fractions for iron in river water, the oxyhydroxides phase, which shows positive δ^{56} values, and the fulvic fraction that has a more negative signal. River water-seawater mixing experiments by Bergquist and Boyle (2006) showed that aggregated Fe was enriched in heavy isotopes. Hence, aggregation and sedimentation of the oxyhydroxide fraction during estuarine mixing should remove heavy isotopes from surface suspended matter, resulting in a more negative signal in the suspended phase, as indicated by field data from the River Lena freshwater plume (J. Ingri, personal communication, 2009). Cycling of iron in coastal areas appears to result in export of a negative iron isotope signal in the truly dissolved fraction, suggesting that open ocean water generally has a negative dissolved isotope iron signal thus explaining the negative δ^{56} value in ferromanganese crusts in the deep-sea. However, recent data indicate that bottom water in the open ocean has a positive δ^{56} value (Lacan et al., 2008), although it should be noted that the dataset is limited to one depth profile.

Iron isotope data from surface water in the Baltic Sea reveal systematic temporal variations in the Fe-isotope signal. For example, the δ^{56} value changed from -0.1 to $+0.25\%$ during a diatom spring-bloom resulting in subsequent sedimentation of iron with a negative isotope signal (Gelting et al., 2009). During the summer a relatively stable positive δ^{56} value was measured in suspended matter at different locations. This was likely due to a combination of river introduced aggregated oxyhydroxides and particulate iron formed from oxidation of dissolved Fe(II) in surface water. In this low salinity system, river introduced Fe-oxyhydroxides aggregate, but may not sediment in the river estuaries due to the lack of detrital sinking and flocculation processes and hence can spread far into the Baltic Sea (Gustafsson et al., 2000). This system is in sharp contrast to recently revised very rapid aggregation and sedimentation processes for direct river – seawater mixing (Nowostawska et al., 2008).

Recent advances suggest that iron isotope measurements have a large potential to provide new information on iron cycling and iron transport from coastal areas to the open ocean (de Jong et al., 2007). Fe/Ti or Fe/Al ratios close to average crust material do not necessarily indicate that the suspended phase mainly reflects detrital particles. Both positive and negative iron isotope values have been measured although the sample has a Fe/Ti or Fe/Al ratio close to average crust material. Furthermore, a δ^{56} value around zero does not necessarily

mean that the sample consists of mainly detrital rock fragments, as it usually is a mixture of iron particles with positive and negative δ^{56} values (Gelting et al., 2009). Recommendations for future work thus consist of a focus on this field including continuing the characterization of the carrier phase for Fe across the salinity gradient and into the open ocean.

5 Colloidal iron and organic matter

Ten years ago the focus of Fe biogeochemistry was on dissolved (filterable) iron speciation and quite specific iron complexes. Once overlooked and neglected (Wells, 1998), progress has been made in understanding the nature and importance of organic colloidal material in seawater and coastal systems and challenged the simple discrimination into particulate and dissolved iron (0.45 or $0.2\ \mu\text{m}$ filtered). Furthermore, dynamic exchange between larger iron particles, colloidal iron, and soluble iron (defined as passing either a $0.02\ \mu\text{m}$ or a $1\ \text{kDa}$ filter) also directs interest towards the particulate and soluble phase. The FeCycle study, a mesoscale SF6 tracer release experiment without iron perturbation in HNLC waters southeast of New Zealand (Boyd et al., 2005), showed that iron recycling rates due to biological iron uptake and regeneration exceeded input of new iron by 10-fold. Further, particulate Fe would undergo a transformation from lithogenic to biogenic iron during settling through the mixed layer. Rapid biological processing (bacterivory and herbivory, subsequent biological uptake) after dissolution of dust deposited iron hydroxides and presumably also photolysis of siderophore complexed Fe(III)-hydroxide (Borer et al., 2005) resulted in exchange from the lithogenic particulate phase via the soluble to the biogenic particulate phase (Frew et al., 2006; Strzepek et al., 2005; Maldonado et al., 2005). The rapid exchange with particulate iron phases provides new insight into iron cycling and export dynamics since the role of particulate iron in iron biogeochemistry appears more important than previously assumed. During the Gothenburg workshop however, the main center of attention was on colloidal iron and we therefore focus thereon hereafter.

Moran et al. (1996) measured iron, among other bioactive trace metals, in colloidal matter obtained by cross-flow filtration of seawater. The major proportion of the dissolved Fe in open ocean seawater (here defined as $0.4\ \mu\text{m}$ filtered) was found to be in the colloidal form (here defined as $>0.02\ \mu\text{m}$ – $0.4\ \mu\text{m}$) (Wu et al., 2001), with continuing debate about the bioavailability of this fraction. Chen and Wang (2001) showed that freshly precipitated colloids were available to phytoplankton but aging processes (15 days) reduced markedly their availability. Wang and Dei (2003) demonstrated that Fe availability from colloidal matter to cyanobacteria (*Synechococcus*, *Trichodesmium*) is largely dependent on the size and origin of the material, with the tendency of Fe bound to smaller colloids and biogenic colloidal

material derived from the same species being more available. The transfer from the soluble to the colloidal fraction appears rapid for iron in comparison to for example Zn, resulting in dynamic cycling including particle formation, and the drawdown of colloidal Fe indicated uptake by phytoplankton (Hurst and Bruland, 2007). Further, colloidal Fe is photoreactive and thus also contributes to the bioavailable pool of Fe(II) in surface waters (Barbeau, 2006; Fan, 2008).

Dissolved organic matter (DOM, which contains the colloidal fraction) in seawater has previously been considered to be old (~ 6 KY) and refractory (Bauer et al., 1992). This refractory pool is also known to be rich in aromatic chromophoric material therefore often called chromophoric DOM (CDOM). However, in the last ten years the picture has changed somewhat and now it is believed that in addition to the refractory pool seawater DOM also consists of in situ biologically derived material, rich in proteic matter and saccharides and saccharide derivatives as building blocks (Aluwihare et al., 1997). In addition significant findings have been made to understand that fractions of marine DOM possess a gel forming character, including spontaneous assembling into microgels after filtration, where calcium bridging is shown to be important (Chin et al., 1998). In addition, new microscopy based techniques have shown that fibrillar type materials, hypothesized to consist of acid polysaccharides, are abundant in many open ocean regimes (Santschi et al., 1998). This marine gel phase can be an important transfer route from truly dissolved to particulate pool of matter (Verdugo et al., 2004). These findings in the dissolved fraction seem to link well to the marine snow formation of transparent exopolymeric particles (TEP), which are important for carbon export (Engel et al., 2004), although direct experimental evidence linking the fibrillar material to TEP and sedimentation has been lacking.

To what extent these processes and phase transfers in organic matter are controlling the physicochemical states and vertical distributions of iron and other trace elements has previously only been hypothesized. Stolpe and Hassellöv (2010) coupled Flow Field Flow Fractionation (FIFFF) with ICPMS, on-line “humic” fluorescence and UV-absorbance detectors, and subsamples for Atomic Force Microscopy (AFM) to fractionate and identify different colloidal size classes and associated trace metals during phytoplankton bloom events in a Fjord on the North Sea coast of Sweden. They found both seasonal and vertical variations in the colloidal size distributions for iron and other trace elements and could use these in order to explain the apparent iron solubility and vertical distribution to a large extent (Stolpe and Hassellöv, 2010). During the winter season colloidal size distribution for iron (and many other elements) were only appearing in the CDOM fraction (~ 0.5 – 3 nm), while during the spring bloom and summer bloom in two consecutive seasons the colloidal size distributions for iron were shifting dramatically. In addition to the CDOM phase, iron partitioned into two larger size classes. With AFM these two colloidal popu-

lations were identified to be semispherical (3–7 nm) and fibrillar (~ 0.5 nm thick and 30–200 nm long). From the partitioning of other elements and their size and shape it was hypothesized that the semispherical colloids were mainly thiol rich proteic biopolymers, while the fibrillar materials were polysaccharide rich exudates that could be the precursors of the microgels proposed by Chin et al. (1998). The conclusion that the seasonal variations of iron association with different colloidal phases to some extent control the apparent iron solubility in estuarine water is in line with the findings from Bergquist et al. (2007), implying that colloids in the open ocean control iron solubility. Likewise, Boehme and Wells (2006) and Fløge and Wells (2007), using FIFFF coupled to excitation emission matrix spectroscopy and a UV-absorbance detector, observed a shift in colloidal size class distribution between protein-like and humic-like fluorescence of CDOM during phytoplankton blooms in an estuary.

Progress has been made in studying the behavior of iron oxide nanoparticles in different freshwater and salt matrixes and drawing conclusions for the inorganic phase within the filterable fraction (Hassellöv and von der Kammer, 2008, and references therein). Partly based on this work, the understanding of flocculation processes has improved and previous concepts (Sholkovitz, 1978; Sholkovitz et al., 1978) have been confirmed. Mylon et al. (2004) using natural organic matter (NOM) coated synthesized hematite colloids, show that the rate of colloid aggregation reaches a maximum at a salinity of 12, resulting in a removal of 80–90% of dissolved iron in a process occurring on a time scale of seconds (Nowostawska et al., 2008). The colloidal particles are stabilized by NOM due to electrostatic and repulsive forces (Mosley et al., 2003; Sander et al., 2004). Theoretically in seawater the conditions would favor attachment, but low particle concentrations result in a low collision frequency. Further, colloidal matter undergoes a transformation in size distribution and elemental composition upon introduction from fresh water into a seawater system (Stolpe and Hassellöv, 2007). The efficiency of transport through salinity gradients needs more investigation and isotope studies may be of significant importance to form a proper understanding of fluvial iron inputs into the sea (see Sect. 4).

As aforementioned, recent methodological advancements include the application of field flow fractionation (FFF), in conjunction with size fractionation by membrane and/or ultrafiltration techniques, to studies of the metal-colloidal phase (Boehme and Wells, 2006; Stolpe et al., 2005). FFF is generally applied to samples from coastal systems and detection limits necessitate the use of pre-concentration steps. While being a powerful tool to characterize size fractionated material, FFF can also help in developing robust filtering methods particularly at the lower end of the size range, as results reveal artefacts from membrane filtration can result in unintended removal of undersized material (Howell et al., 2006). The relevance of this for open ocean seawater

requires further testing. Further, cross-flow ultra-filters are defined as a molar cut-off, which may result in retention of undersized components and permeation of oversized components, as well as separation of size and chemical composition (Assemi et al., 2004). An intercalibration of cross-flow filtration techniques was carried out previously (Buesseler et al., 1996), but a new approach including classical membrane filtration and utilizing FIFFF coupled to ICPMS may yield valuable information about the robustness of different filter membranes with regard to fractionation of colloidal size classes and their elemental composition.

We conclude that future research directions should encompass further in depth characterizations of the different phases (particulate, colloidal, soluble), which may lead to a redefinition of the term dissolved iron. This will also lead to a better structural definition of bioavailable iron. We need to learn how iron is fractionated into specific size classes, what the exchange kinetics between these phases are, and what controls/catalyzes them. Specifically, the origin and nature of iron binding ligands needs to be further addressed to elucidate the role and characteristics of different ligand classes (L_1 , L_2). In that, we may need to overcome measurement artifacts due to pre-concentration procedures that are necessitated due to the detection limits especially in open ocean applications (see also last two paragraphs of Sect. 6).

6 Linking biological processes to iron chemistry

Most areas of the open ocean have surface trace metal concentrations between picomolar and nanomolar levels, which are about one millionth of the concentration in phytoplankton cells (Morel and Price, 2003). Iron is required for many important cellular processes such as photosynthesis, respiration, nitrogen fixation and nitrate reduction. A recent laboratory study involving 15 neritic and oceanic phytoplankton species produced an elemental ratio of $C_{124}N_{16}P_1Fe_{0.0075}$ (Ho et al., 2003), similar to previous reviews of Fe:C ratios which have found a range of 2.3–370 $\mu\text{mol}:\text{mol}$ (Sarhou et al., 2005; see also Twining et al., 2004). Research has linked the oxygenation of the oceans and the subsequent drop in iron solubility and thus iron availability to the evolution of more iron efficient phytoplankton (Quigg et al., 2003; Saito et al., 2003) that are able to cope with the low iron open ocean conditions. Phytoplankton species have evolved very effective acquisition mechanisms with high trace metal affinities that involve interactions with organic iron binding ligands. Uncertainties remain on the nature of such ligands, which control Fe chemistry and bioavailability in marine systems (Hunter and Boyd, 2007).

Culture experiments have established that marine phyto- and bacterioplankton have different iron requirements that are linked to their biogeographical sources (Sunda and Huntsman, 1995; Brand et al., 1983). More recent work has shown that picophytoplankton, which dominate the olig-

otrophic regions of the oceans, are able to grow optimally in culture at extremely low inorganic iron concentrations of 10–15 pM inorganic Fe, (Timmermans et al., 2005). Our ability to relate these studies to the real environment is however limited by our understanding of the chemical speciation of iron in the ocean (Gledhill and van den Berg, 1994; Rue and Bruland, 1995). These studies indicated that dissolved iron is strongly complexed in the ocean, results which have been confirmed on many occasions since (as discussed in Hunter and Boyd, 2007). The composition of this organic fraction is still not well understood, although it appears likely that it will consist of autochthonous complexing ligands produced by marine phyto- and bacterioplankton (Mawji et al., 2008; Boye et al., 2005; Kondo et al., 2008; Vong et al., 2007) and complex organics such as humic/fulvic acids (Laglera and van den Berg, 2009). Calculations of the inorganic iron concentration based on measurements carried out by competitive equilibration cathodic stripping voltammetry show that inorganic iron concentrations in the ocean are of the order of 10^{-14} – 10^{-11} M (Morel et al., 2008), although these calculations neglect the contribution of Fe(II), which may also be present at concentrations of the order of 10^{-11} M in surface waters (e.g. Hansard et al., 2009; Roy et al., 2008b; Croot et al., 2001). It is not clear how much of the organically complexed iron is available to marine phyto- and bacterioplankton, and parameters controlling Fe bioavailability to primary producers are still poorly understood.

Fe bioavailability is influenced by its chemical forms (speciation, redox state), biological cycling, and the different uptake strategies of the phyto- and bacterio-plankton communities (Barbeau et al., 1996; Hutchins et al., 1999a; Strzepek et al., 2005). Competition for available Fe is strongest when Fe is in short supply (e.g. Worms et al., 2006). Recent advances in our understanding and abilities to model iron uptake by marine phytoplankton (Morel et al., 2008; Shaked et al., 2005; Salmon et al., 2006) indicate that even at these low inorganic iron concentrations, open ocean phytoplankton will have sufficient iron to grow. Initially iron uptake was thought to be proportional to the concentration of inorganic Fe species (Fe^+) (Hudson and Morel, 1990). However, this model proved to be too simplistic to explain phytoplankton growth in natural systems where concentrations of inorganic iron species were extremely low due to organic complexation. Thus either the iron-ligand complex (FeL) is directly taken up, or the inorganic Fe availability is increased, e.g. by reduction to Fe(II). More recently two models have been published to describe the kinetics of Fe uptake. The Fe(II) model by Shaked et al. (2005) and the FeL model by Salmon et al. (2006). There are significant distinctions between these models which lead to differences in the predictions of phytoplankton iron limitation in culture experiments. While the Fe(II) model considers the surface Fe(II) concentration and explicitly includes unchelated Fe(III) as a source of Fe(II) for phytoplankton uptake, the FeL model considers the bulk concentration of Fe(II) in the

media as the controlling parameter and excludes unchelated Fe(III) as an irrelevant source (Morel et al., 2008). Morel et al. (2008) point out that the observed decrease in Fe uptake rates with increasing EDTA concentrations can only be explained by the Fe(II) model, which results in the conclusion that unchelated Fe(III) is indeed an important source of Fe(II) for phytoplankton uptake. However, phytoplankton species behave differently under Fe limitation and it is likely that future experiments under more natural conditions without the presence of EDTA will result in more realistic iterations of the iron uptake models. The role of other trace metals and organic material in the partly species specific adaptations of the iron acquisition system are not completely understood. As one example, Peers and Price (2006) have shown that copper is essential for electron transport in *T. oceanica* regardless of Fe status implying that selection pressure imposed by Fe limitation has resulted in the use of a Cu protein for photosynthesis in an oceanic diatom.

Adaptations to low iron environments have been found to include a reduction in cell size (Sarhou et al., 2005), changes in photosynthetic architecture (Strzepek and Harrison, 2004; Peers and Price, 2006) and substitution of iron containing proteins for non-iron containing proteins (Peers and Price, 2006; McKay et al., 1999). Further possible adaptations include the induction of high affinity uptake mechanisms such as the production of siderophores by marine prokaryotes (Vraspir and Butler, 2009) and uptake mechanisms that target specific iron containing compounds such as hemes (Hopkinson et al., 2008) or the production of iron storage proteins (Marchetti et al., 2009).

Microorganisms can exert a feedback effect on Fe chemistry, for example by releasing organic matter which is able to react with Fe (e.g. siderophores, exopolymeric substances (EPS), cell lysis material or fecal pellets), which can enhance iron bioavailability (e.g. Hutchins et al., 1999b). Høgdal et al. (1996), for example, visualized and quantified metals bound to bacterial extracellular matrixes in applying X-ray transmission electron microscopy. The role of grazing as a source of organic, iron binding material via sloppy feeding and/or as a direct source of iron is often discussed. Several studies address this topic and a general consensus about the importance of grazing for iron recycling in surface seawater exists (Sato et al., 2007; Barbeau et al., 1996; Dalbec and Twining, 2009; Sarhou et al., 2008; Hutchins and Bruland, 1994; Hutchins et al., 1995; Tovar-Sanchez et al., 2007; Zhang and Wang, 2004). However, some results are inconsistent and the detailed mechanisms as well as the contribution of different grazer types such as protozoa, copepods, krill, and salps and their specific feeding mechanisms are poorly understood. Therefore, it is difficult today to estimate the overall function of grazing on the biogeochemical cycles of iron especially in HNLC regions.

Most marine microorganisms (bacterio- and phytoplankton) produce polysaccharides that are either stored as energy reserves or secreted as exopolymeric substances

(EPS) (Schoemann et al., 2001; Decho, 1990; Hoagland et al., 1993). It has recently been shown that iron starvation is coupled to transparent exopolymer particles (TEP) production in *Trichodesmium* (Berman-Frank et al., 2007). Recent studies also provide evidence that high concentrations of saccharides or carbon-rich organic matrices can enhance the growth of phytoplankton (Vasconcelos et al., 2002) and efficiently retain Fe (II) (Öztürk et al., 2004; Toner et al., 2009), a highly bioavailable form (Morel et al., 2008). Steigenberger et al. (2010) show that polysaccharides and cell exudates of *Phaeodactylum* sp. can also result in high hydrogen peroxide production, while the authors still observe a net stabilizing effect of Fe(II) potentially due to a combination of organic Fe(II) retention paralleled by superoxide production.

Hassler and Schoemann (2009) explore a Fe-related biogeochemical role for polysaccharides, by examining the influence of various organic ligands (siderophore, porphyrin, mono- and poly-saccharides) on iron solubility and its bioavailability to four keystone phytoplankton species of the Southern Ocean, representing different phytoplankton functional groups and size classes (*Phaeocystis* sp., *Chaetoceros* sp., *Fragilariopsis kerguelensis* and *Thalassiosira antarctica* Comber). Results show that saccharides can increase Fe uptake rates and Fe solubility above the level observed for inorganic Fe. Similar observations were made on natural plankton community from the Southern Ocean (Hassler et al., 2007). Given the ubiquitous presence of saccharides in the ocean, these compounds might represent an important factor to control the basal level of soluble and bioavailable Fe.

Over the past years, the Fe(II) pool has been recognized as an important source of bioavailable Fe and intermediate in Fe cycling. Albeit short-lived due to rapid re-oxidation to Fe(III), significant concentrations of Fe(II) were detected in different oceanic and coastal provinces (Breitbarth et al., 2009; Croot and Laan, 2002; Croot et al., 2008, 2005; Roy et al., 2008b; Hopkinson and Barbeau, 2007; Ussher et al., 2007). There has been emerging evidence that Fe(II) is retained in oxygenated water by organic ligands (Croot et al., 2001), which may be a product of marine biota (Roy et al., 2008b) or also of other origin and rain introduced (Willey et al., 2008). See Barbeau (2006) for a comprehensive review and also Sect. 7. The role of Fe(II) for phytoplankton nutrition and Fe(II) organic complexation provide interesting and relevant research topics for the near future.

Iron limitation also induces a decoupling in the use of macronutrients by phytoplankton, likely to influence the cycling of the major biogeochemical cycles (C, N, P, Si, S) over geological time scales (de Baar and La Roche, 2003). Further, light intensity can play an important role (Hoffmann et al., 2008; Maldonado et al., 1999; de Baar et al., 2005; Moore et al., 2007a and b). Moreover, Bucciarelli et al. (2010) examined the effect of Fe-light co-limitation on cellular silica, carbon and nitrogen in two marine diatom species, *Thalassiosira oceanica* and *Ditylum brightwellii*, observing a

1.4-fold increase in C:N ratio with a decrease in growth rate by 70% in both species and a decrease in biogenic silica per cell under severe Fe or Fe-light limitation. These results however are seemingly in contradiction with many previous lab and field studies showing increased diatom silicification under Fe limitation (Hutchins and Bruland, 1998; Takeda, 1998; Firme et al., 2003; Franck et al., 2003).

A significant contribution to the increasing knowledge on the interaction of biological processes with iron chemistry is made by the improvement of methods in this field. Inter-calibrations of Fe detection methods were carried out and measurements of Fe are now possible in near real time in the field at picomolar level (Bowie et al., 2002, 2005, 2006; Johnson et al., 2007), including Fe(II) (Croot and Laan, 2002). More sophisticated shipboard incubation systems are available (Hare et al., 2007a; Hutchins et al., 2003; Pickell et al., 2009; Hare et al., 2005), allowing for more realistic experimental designs to assess Fe phytoplankton interactions. Methods were developed to detect cell surface Fe reduction and uptake (Shaked et al., 2004) and to measure cellular Fe (Hassler et al., 2004). New highly sensitive electrochemical methods have pushed our understanding of organic iron complexation in new directions (Croot and Johansson, 2000; Laglera and van den Berg, 2009). Utilization of laboratory based extensive instrumentation such as FIFFF, x-ray spectroscopy with TEM microscopy, as well as bioreporters, molecular techniques and genomic information allow for in depth studies and visualization of Fe limitation and Fe organic matter interactions (e.g. Heldal et al., 1996; Toner et al., 2009; Stolpe et al., 2005) and particularly also of iron bioavailability (Lam et al., 2006; Boyanapalli et al., 2007; Hassler et al., 2006).

Nevertheless, some methods still depend on high material/biomass concentrations and future development and work may lead towards more direct measurement techniques overcoming pre-concentration artifacts. The majority of phytoplankton iron interaction studies have been carried out *in vitro* and with a limited range of species, and mostly did not include co-effects of other trace metals. Strong iron chelators such as EDTA and DFOB are commonly used to induce iron limitation in culture experiments and experiments are in part difficult to compare due to the variety and combination of factors (e.g. light intensity, temperature) applied. Thus, it is not clear how predominant the known low iron regime adaptations are in the oceanic environment. Albeit very challenging, future experiments should aim towards using more realistic media chemistries and natural biomass densities of cultures that were recently isolated. Methods and experiments need to be designed to link Fe chemistry to biological processes, including potential biological feedback mechanisms on Fe chemistry as also discussed with regard to climate change in Sect. 7. Further, our increasing ability to detect and characterize iron in seawater and in organisms (Mawji et al., 2008; Gledhill, 2007; Laglera and van den Berg, 2009; Vong et al., 2007) coupled to developments in

techniques such as shotgun genomics (Rusch et al., 2007; Venter et al., 2004; Yooseph et al., 2007) and the potential of proteomics (Nunn and Timperman, 2007; Dupont et al., 2006) should lead to great advances over the coming years in our understanding of how organisms have adapted to low iron environments, and the implications of these adaptations to overall marine productivity and biodiversity. The development of *in-situ* measurement technology as for example suggested in the approach of Roy et al. (2008a), with the potential for deployment on moored sensor arrays, will greatly improve the spatial and temporal resolution of Fe measurements.

7 Iron and climate change

Global climate change will greatly influence atmospheric and hydrographic processes in the future. Most prominent features include changes in thermohaline circulation of the North Atlantic, warming of the polar regions, changing wind patterns resulting in reduced upwelling and wind driven mixing, as well as increased sea-surface temperatures and stratification (Boyd and Doney, 2003). Projected changes in relative humidity and land vegetation cover, affecting soil moisture and local dust availability, together with changed patterns in wind and precipitation, as well as riverine transport, will ultimately modify the iron supply to the open ocean (Boyd and Doney, 2003; Jickells et al., 2005). Further, rising atmospheric CO₂ acidifies the oceans, leading to changes in saturation state with respect to calcium carbonate and shifts the aragonite and calcite saturation depths (Feely et al., 2004) and potentially trace metal solubility. The abovementioned processes, albeit uncertainty over their magnitude and exact interrelations in the future exists, will affect marine biota, causing regime shifts, and modifications of biogeochemical cycling (Boyd and Doney, 2003). While climate change needs to be understood holistically, there is a need to evaluate regional and small scale physical, chemical, and biological processes in order to derive potential biogeochemical feedback mechanisms.

We here focus on direct local effects acting upon iron chemistry in seawater and primarily discuss the emerging field of trace metal biogeochemistry research encompassing two main areas, temperature shifts and changing seawater pH. Both temperature and pH are master variables for chemical and biological processes and effects on trace metal biogeochemistry may be multifaceted and complex. Ten years ago, this research field did not exist and data are scarce. Assessing the potential effects of sea-surface warming and ocean acidification on iron biogeochemistry is crucial and predictions to date are based on our understanding of the current ocean system. Despite the expanding knowledge and increasing awareness for trace metal chemistry in open ocean research during the past 20 years and the recently defined

field and intensifying work on ocean acidification research, there is yet little communication between these fields.

A decrease of the surface seawater pH from pre-industrial 8.25 to 7.85 within this century, and further by up to 0.7 units until 2300 is predicted (Caldeira and Wickett, 2003; Jacobson, 2005). In general, the H^+ ion concentration can directly affect metal uptake by phytoplankton via altered membrane transport activity or via direct competition of the H^+ ion with metal ions for membrane transporters or other metabolically active sites on the cell surface (Sunda and Huntsman, 1983; Vigenault and Campbell, 2005). Further main aspects are the inorganic solubility of iron, changes in organic complexation, phytoplankton – trace metal feedback mechanisms, and differences in redox chemistry.

$Fe(OH)_3$ solubility and $Fe(III)$ inorganic speciation are expected to change with ocean acidification (Liu and Millero, 2002, 1999). When seawater pH falls below 8, changes in the inorganic speciation result in an increase of the thermodynamic $Fe(III)$ hydroxide solubility. Enhanced solubility above pH 7 in seawater of the warm or temperate ocean though is mainly due to organic ligands and suggests any change in solubility arising from acidification will be mainly related to the organic complexes (Liu and Millero, 2002). However, in cold water the solubility of Fe can exceed FeL concentrations (P. Croot, personal communication, 2009, calculated based on Liu and Millero, 2002), bringing inorganic speciation shifts due to pH and temperature back into the game. Interesting questions arise concerning whether ocean acidification could potentially also affect metal leaching from atmospheric deposits (see Sects. 2.1 and 2.2) and how the metastable colloidal Fe phase may be affected (see Sect. 5). The potential effect of pH acting directly on FeL complexes depends on the nature of Fe-binding functional groups. The H^+ stoichiometry of the $Fe(III)$ binding sites defines the magnitude of acid dissociation constants (pK_a). Carboxyl groups have a $pK_a \sim 5$ and thus the conditional stability constant of the FeL complex ($\log K_{FeL}$) should remain unchanged above pH 6. In contrast, phenolic groups have a $pK_a \sim 9$ and $\log K_{FeL}$ will increase with pH (Sillen and Martell, 1971). Both groups can be found in siderophores. While to date no published experimental data on the pH effect for FeL can be found, Averyt et al. (2004) show a decrease of $\log K_{CuL}$ with lower pH in two lakes. Similar effects were observed for Cd ligands, but less so for Zn ligands (Sander et al., 2007). Further, iron chelates are more photolabile at lower pH (Sunda and Huntsman, 2003), which directly involves effects on Fe photochemistry (see below). Overall though, while FeL complexes may or may not be directly pH affected, alterations of organic iron complexation may still arise from biological ligand production processes, should those be affected by pH and/or temperature (see below). Several models of Fe uptake mechanisms for phytoplankton exist (Morel et al., 2008; Shaked et al., 2005; Salmon et al., 2006, see also Sect. 6) and their pH dependence may be largely connected on their reliance on $Fe(II)$ as

the actual species taken up and on the species capability to regulate pH at the cell surface.

It can be expected that pH driven changes in trace metal availability will trigger biological feedback mechanisms, which regulate trace metal availability to marine phytoplankton. These can be in form of exudates, cell lysates, or chlorophyll degradation products, and can serve as trace metal ligands to prevent toxic effects or to increase trace metal uptake rates. The capability of eukaryotic phytoplankton species to produce trace metal binding ligands either to prevent toxic effects or to increase uptake has been addressed (Ahner et al., 1997; Barbeau et al., 2001; Hutchins et al., 1999b). However, information on biological feedback mechanisms in response to climate change that affect trace metal chemistry is very limited. It should be noted that in contrast to the open ocean, estuaries and coastal areas might show a wide range in pH (5 to >9) (e.g. Chen and Durbin, 1994; Sunda and Huntsman, 1998) and obviously in temperature, to which phytoplankton species are adapted to. However, even considering that phytoplankton blooms may cause temporal increases in surface water pH due to CO_2 uptake, open ocean species are adapted to a very narrow range in pH. Further, some coastal areas such as the Oregon shelf temporarily experience subsurface input of low pH water and such systems could be valuable analogs for acidification and temperature effects in natural settings.

Several studies were carried out during the past years assessing changes in phytoplankton physiology using laboratory batch cultures and mesocosm pCO_2 perturbations. Changes in carbon and nitrogen fixation rates, calcification rates, and carbon export are reflective of pH effects on the biogeochemistry of the manipulated system (Riebesell et al., 2007; Orr et al., 2005, see also Biogeosciences Special Issue “The ocean in the high- CO_2 world II”), which unequivocally will also affect trace metal cycling. Further studies also reported combined effects of pCO_2 and temperature change (Hare et al., 2007b), and modeling studies also suggest potential interactions with irradiance effects due to changing stratification in the future ocean on phytoplankton physiology and species composition (Boyd and Doney, 2002). Seen in coherence with biological effects on organic Fe complexation, and in return again with Fe availability effects on phytoplankton, these studies indicate that phytoplankton physiology and species composition could exert biological feedback mechanisms on trace metal cycling as a function of pCO_2 and temperature in seawater.

Data from a coastal mesocosm CO_2 enrichment experiment (Breitbarth et al., 2010) suggest increasing dissolved iron concentrations with ocean acidification. The authors invoke a biological feedback mechanism at future seawater pCO_2 resulting in increased organic $Fe(III)$ complexation, which requires further testing. More so, changes in $Fe(II)$ chemistry were observed. In part, the underlying processes can theoretically be derived based on established relationships of $Fe(II)$ oxidation rates and inorganic $Fe(II)$

speciation in presence of different oxidizers over environmentally relevant ranges in pH, temperature, and salinity (Santana-Casiano et al., 2006; Gonzalez-Davila et al., 2006; Santana-Casiano et al., 2005; Millero and Sotolongo, 1989; Millero et al., 1987; Croot and Laan, 2002). For example, over a seawater pH decrease of 0.5 units, a 10-fold increase in the half-life of Fe(II) can be expected and the effects of ocean acidification may thus override the influences from sea-surface temperature changes (Santana-Casiano et al., 2005). Fe(II) oxidation kinetics are seemingly affected by organic complexation (e.g. Croot et al., 2001; Rose, 2003; Roy et al., 2008b, see also Sect. 6). Fe(II) ligands may be biologically mediated and potential biological feedback mechanisms in the future could thus further complicate the picture, requiring focused research in this field. Moreover, changing light regimes are expected to affect photochemical cycling of Fe in sunlit surface waters (Boyd and Doney, 2002). Both, light intensity and the light spectrum penetrating the water will influence photochemical processes and this field requires further attention.

Similar to seawater pH, temperature effects have been rarely studied in coherence with trace metal biogeochemical measurements in open ocean systems. It has been standard to date to carry out measurements of organic iron complexation at room temperature, but temperature has profound effects on metal speciation and solubility. Further, Rose et al. (2009) demonstrate synergistic effects of temperature and iron additions on phytoplankton physiology and community dynamics in Ross Sea waters. Likewise, Fu et al. (2008) demonstrate that $p\text{CO}_2$ perturbations alone may not give the sole answer to potential physiological changes in phytoplankton, since these can be modified by interactions with Fe limitation. CO_2 and N_2 fixation rates in the future ocean may be controlled by a combination of Fe availability and $p\text{CO}_2$, further stressing the need to elucidate future changes in seawater iron chemistry.

Overall, climate change effects on iron speciation and biological limitation are likely not going to be driven by a single factor, and Rose et al. (2009) stress the importance of multivariate studies in order to understand ecosystem changes. It also remains to be shown how climate change may alter the interrelations of iron with other trace metals and macronutrients. For example, laboratory experiments showed that cadmium toxicity can be reduced under high iron availability, suggesting that cadmium is a competitive inhibitor of the iron uptake system or iron dependent cellular processes (Foster and Morel, 1982; Sunda and Huntsman, 2000). Similar observation are made for iron limited natural phytoplankton assemblages from the Southern Ocean by Cullen et al. (2003) who suggest that Fe limited phytoplankton take up more Cd resulting in lower Cd:PO₄ ratios in surface waters. Iron co-limitations and interactions with other nutrients and trace metals have been observed (e.g. Schulz et al., 2004; Mills et al., 2004; Wu et al., 2003; Wells et al., 2005) and apparently the composition of trace metals and macro nutrients greatly

affect natural Fe fertilization efficiency (see Sect. 2.2). Moreover, Statham et al. (2008) recently addressed glacier meltwater input of iron and colloidal matter from the Greenland Ice Sheet. In the context of the expected changes for Fe biogeochemistry discussed here, their study illustrates how atmospheric warming can act on various levels, evidentially affecting iron biogeochemistry in the sea.

We conclude that ocean acidification may result in increased Fe(III) solubility, is likely to decrease stability of some FeL complexes, and is likely to increase Fe(II) stability. It may also change the mechanisms of Fe acquisition by cells, which though depends on the Fe status of the regime and the type of phytoplankton species present. Temperature effects may be smaller in comparison, with most pronounced changes though to be expected in polar waters. Recommendations for future research directions are systematic measurements of Fe(III) solubility in pH range 7–9 and effects of Fe-binding ligands along with the study of temperature effects thereon, and field experiments in upwelling regions with a focus on low pH regimes. Moreover, the role of organic ligands in enhancing Fe(II) stability needs to be investigated as well as effects of pH and temperature on the photoreactivity of Fe(III)L complexes. It is largely unknown what the pH controls in organisms are, and how they affect Fe acquisition. More emphasis is needed on measurements and control of the seawater carbonate system, including pH, in field studies and laboratory cultures. Protocols carried out to achieve pH control need to be reported and researchers are urged to report pH data on the total or seawater pH scale to ensure comparability of different studies. The comprehensive “Guide for Best Practices in Ocean Acidification Research and Data Reporting” was recently published and should be adapted for trace metal research (<http://www.epoca-project.eu/index.php/Home/Guide-to-OA-Research/>).

Acknowledgements. Funding for the workshop was provided by EUR-OCEANS and the Swedish Research Council (VR). E. B. and L. J. H. acknowledge current funding by the German Research Foundation (DFG, BR 3794 and HO 4217). S. D. is funded by the DFG and the multi-disciplinary research group NOVUM (Nutrients Originating in Volcanoes and their effect on the eUphotic zone of the Marine ecosystem) by the Leibniz Institute of Marine Sciences, IFM-GEOMAR. D. H. acknowledges funding by US NSF ANT 0741411 and OCE 0825319. The authors greatly appreciate the comments received from P. W. Boyd on the manuscript.

Edited by: U. Riebesell

References

- Aguilar-Islas, A. M., Rember, R. D., Mordy, C. W., and Wu, J.: Sea ice-derived dissolved iron and its potential influence on the spring algal bloom in the Bering Sea, *Geophys. Res. Lett.*, 35, L24601, doi:10.1029/2008gl035736, 2008.
- Ahner, B. A., Morel, F. M. M., and Moffett, J. W.: Trace metal control of phytochelatin production in coastal waters, *Limnol. Oceanogr.*, 42, 601–608, 1997.
- Aluwihare, L. I., Repeta, D. J., and Chen, R. F.: A major biopolymeric component to dissolved organic carbon in surface sea water, *Nature*, 387, 166–169, 1997.
- Andersson, K., Dahlqvist, R., Turner, D., Stolpe, B., Larsson, T., Ingri, J., and Andersson, P.: Colloidal rare earth elements in a boreal river: Changing sources and distributions during the spring flood, *Geochim. Cosmochim. Acta*, 70, 3261–3274, doi:10.1016/j.gca.2006.04.021, 2006.
- Ardelan, M. V., Holm-Hansen, O., Hewes, C. D., Reiss, C. S., Silva, N. S., Dulaiova, H., Steinnes, E., and Sakshaug, E.: Natural iron enrichment around the Antarctic Peninsula in the Southern Ocean, *Biogeosciences*, 7, 11–25, 2010, <http://www.biogeosciences.net/7/11/2010/>.
- Assemi, S., Newcombe, G., Hepplewhite, C., and Beckett, R.: Characterization of natural organic matter fractions separated by ultrafiltration using flow field-flow fractionation, *Water Res.*, 38, 1467–1476, doi:10.1016/j.watres.2003.11.031, 2004.
- Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochem. Cy.*, 20, GB2017, Gb2017, doi:10.1029/2005gb002591, 2006.
- Averyt, K. B., Kim, J. P., and Hunter, K. A.: Effect of pH on measurement of strong copper binding ligands in lakes, *Limnol. Oceanogr.*, 49, 20–27, 2004.
- Baker, A. R. and Jickells, T. D.: Mineral particle size as a control on aerosol iron solubility, *Geophys. Res. Lett.*, 33, L17608, doi:10.1029/2006GL026557, 2006.
- Baker, A. R., Jickells, T. D., Witt, M., and Linge, K. L.: Trends in the solubility of iron, aluminium, manganese and phosphorus in aerosol collected over the Atlantic Ocean, *Mar. Chem.*, 98, 43–58, doi:10.1016/j.marchem.2005.06.004, 2006.
- Baker, A. R., and Croot, P. L.: Atmospheric and marine controls on aerosol iron solubility in seawater, *Mar. Chem.*, in press, 2009.
- Barbeau, K., Moffett, J. W., Caron, D. A., Croot, P. L., and Erdner, D. L.: Role of protozoan grazing in relieving iron limitation of phytoplankton, *Nature*, 380, 61–64, 1996.
- Barbeau, K. and Moffett, J. W.: Laboratory and field studies of colloidal iron oxide dissolution as mediated by phagotrophy and photolysis, *Limnol. Oceanogr.*, 45, 827–835, 2000.
- Barbeau, K., Rue, E. L., Bruland, K. W., and Butler, A.: Photochemical cycling of iron in the surface ocean mediated by microbial iron(III)-binding ligands, *Nature*, 413, 409–413, 2001.
- Barbeau, K.: Photochemistry of organic iron(III) complexing ligands in oceanic systems, *Photochem. Photobiol.*, 82, 1505–1516, doi:10.1562/2006-06-16-ir-935, 2006.
- Bauer, J. E., Williams, P. M., and Druffel, E. R. M.: C-14 activity of dissolved organic-carbon fractions in the North-Central Pacific and Sargasso Sea, *Nature*, 357, 667–670, 1992.
- Benitez-Nelson, C. R., Vink, S. M., Carrillo, J. H., and Huebert, B. J.: Volcanically influenced iron and aluminum cloud water deposition to Hawaii, *Atmos. Environ.*, 37, 535–544, 2003.
- Bergquist, B. A. and Boyle, E. A.: Iron isotopes in the Amazon River system: Weathering and transport signatures, *Earth Planet. Sci. Lett.*, 248, 54–68, doi:10.1016/j.epsl.2006.05.004, 2006.
- Bergquist, B. A., Wu, J., and Boyle, E. A.: Variability in oceanic dissolved iron is dominated by the colloidal fraction, *Geochim. Cosmochim. Acta*, 71, 2960–2974, doi:10.1016/j.gca.2007.03.013, 2007.
- Berman-Frank, I., Rosenberg, G., Levitan, O., Haramaty, L., and Mari, X.: Coupling between autocatalytic cell death and transparent exopolymeric particle production in the marine cyanobacterium *Trichodesmium*, *Environ. Microbiol.*, 9, 1415–1422, doi:10.1111/j.1462-2920.2007.01257.x, 2007.
- Blain, S., Queguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C., Brussaard, C., Carlotti, F., Christaki, U., Corbiere, A., Durand, I., Ebersbach, F., Fuda, J.-L., Garcia, N., Gerringa, L., Griffiths, B., Guigue, C., Guillerme, C., Jacquet, S., Jeandel, C., Laan, P., Lefevre, D., Lo Monaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.-H., Picheral, M., Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M., Thuiller, D., Timmermans, K., Trull, T., Uitz, J., van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L., and Wagener, T.: Effect of natural iron fertilization on carbon sequestration in the Southern Ocean, *Nature*, 446, 1070–1074, 2007.
- Blain, S., Bonnet, S., and Guieu, C.: Dissolved iron distribution in the tropical and sub tropical South Eastern Pacific, *Biogeosciences*, 5, 269–280, 2008, <http://www.biogeosciences.net/5/269/2008/>.
- Boehme, J. and Wells, M.: Fluorescence variability of marine and terrestrial colloids: Examining size fractions of chromophoric dissolved organic matter in the Damariscotta River estuary, *Mar. Chem.*, 101, 95–103, doi:10.1016/j.marchem.2006.02.001, 2006.
- Bopp, L., Kohfeld, K. E., Le Quere, C., and Aumont, O.: Dust impact on marine biota and atmospheric CO₂ during glacial periods, *Paleoceanography*, 18, 1046, doi:10.1029/2002pa000810, 2003.
- Borer, P. M., Sulzberger, B., Reichard, P., and Kraemer, S. M.: Effect of siderophores on the light-induced dissolution of colloidal iron(III) (hydr)oxides, *Mar. Chem.*, 93, 179–193, 2005.
- Bowie, A. R., Achterberg, E. P., Sedwick, P. N., Ussher, S., and Worsfold, P. J.: Real-time monitoring of picomolar concentrations of iron(II) in marine waters using automated flow injection-chemiluminescence instrumentation, *Environ. Sci. Technol.*, 36, 4600–4607, 2002.
- Bowie, A. R., Achterberg, E. P., Ussher, S., and Worsfold, P. J.: Design of an automated flow injection-chemiluminescence instrument incorporating a miniature photomultiplier tube for monitoring picomolar concentrations of iron in seawater, *Journal of Automated Methods and Management in Chemistry*, 2005, 37–43, doi:10.1155/JAMMC.2005.1137, 2005.
- Bowie, A. R., Achterberg, E. P., Croot, P. L., de Baar, H. J. W., Laan, P., Moffett, J. W., Ussher, S., and Worsfold, P. J.: A community-wide intercomparison exercise for the determination of dissolved iron in seawater, *Mar. Chem.*, 98, 81–99, 2006.
- Boyanapalli, R., Bullerjahn, G. S., Pohl, C., Croot, P. L., Boyd, P. W., and McKay, R. M. L.: Luminescent whole-cell cyanobacterial bioreporter for measuring Fe availability in diverse marine environments, *Appl. Environ. Microbiol.*, 73, 1019–1024, doi:10.1128/aem.01670-06, 2007.

- Boyd, P.: Ironing out algal issues in the southern ocean, *Science*, 304, 396–397, 2004.
- Boyd, P. W., Wong, C. S., Merrill, J., Whitney, F., Snow, J., Harrison, P. J., and Gower, J.: Atmospheric iron supply and enhanced vertical carbon flux in the NE subarctic Pacific: Is there a connection?, *Global Biogeochem. Cy.*, 12, 429–441, 1998.
- Boyd, P. W., Watson, A. J., Law, C. S., and Abraham, E. R.: A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization, *Nature*, 407, 695–702, 2000.
- Boyd, P. W., Crossley, A. C., DiTullio, G. R., Griffiths, F. B., Hutchins, D. A., Queguiner, B., Sedwick, P. N., and Trull, T. W.: Control of phytoplankton growth by iron supply and irradiance in the subantarctic Southern Ocean: Experimental results from the SAZ Project, *J. Geophys. Res.-Oceans*, 106, 31573–31583, 2001.
- Boyd, P. W. and Doney, S. C.: Modeling regional responses by marine pelagic ecosystems to global climate change, *Geophys. Res. Lett.*, 29, 5351–5354, 2002.
- Boyd, P. W. and Doney, S. C.: The impact of climate change and feedback processes on the ocean carbon cycle, in: *Ocean Biogeochemistry, Global Change – the IGBP Series*, Springer-Verlag Berlin, Berlin, 157–193, 2003.
- Boyd, P. W., Law, C. S., Wong, C. S., Nojiri, Y., Tsuda, A., Levasseur, M., Takeda, S., Rivkin, R., Harrison, P. J., Strzepek, R., Gower, J., McKay, R. M., Abraham, E., Arychuk, M., Barwell-Clarke, J., Crawford, W., Crawford, D., Hale, M., Harada, K., Johnson, K., Kiyosawa, H., Kudo, I., Marchetti, A., Miller, W., Needoba, J., Nishioka, J., Ogawa, H., Page, J., Robert, M., Saito, H., Sastri, A., Sherry, N., Soutar, T., Sutherland, N., Taira, Y., Whitney, F., Wong, S. K. E., and Yoshimura, T.: The decline and fate of an iron-induced subarctic phytoplankton bloom, *Nature*, 428, 549–553, 2004.
- Boyd, P. W., Law, C. S., Hutchins, D. A., Abraham, E. R., Croot, P. L., Ellwood, M., Frew, R. D., Hadfield, M., Hall, J., Handy, S., Hare, C., Higgins, J., Hill, P., Hunter, K. A., LeBlanc, K., Maldonado, M. T., McKay, R. M., Mioni, C., Oliver, M., Pickmere, S., Pinkerton, M., Safi, K., Sander, S., Sanudo-Wilhelmy, S. A., Smith, M., Strzepek, R., Tovar-Sanchez, A., and Wilhelm, S. W.: FeCycle: Attempting an iron biogeochemical budget from a mesoscale SF6 tracer experiment in unperturbed low iron waters, *Global Biogeochem. Cy.*, 19, GB4S20, doi:10.1029/2005GB002494, 2005.
- Boyd, P. W., Jickells, T., Law, C. S., Blain, S., Boyle, E. A., Buesseler, K. O., Coale, K. H., Cullen, J. J., de Baar, H. J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N. P. J., Pollard, R., Rivkin, R. B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., and Watson, A. J.: Mesoscale Iron Enrichment Experiments 1993–2005: Synthesis and Future Directions, *Science*, 315, 612–617, doi:10.1126/science.1131669, 2007.
- Boyd, P. W.: Implications of large-scale iron fertilization of the oceans – Introduction and synthesis, *Mar. Ecol.-Prog. Ser.*, 364, 213–218, 2008.
- Boyd, P. W., Mackie, D. S., and Hunter, K. A.: Aerosol iron deposition to the surface ocean – Modes of iron supply and biological responses, *Mar. Chem.*, in press, doi:10.1016/j.marchem.2009.01.008, 2009.
- Boye, M., Nishioka, J., Croot, P. L., Laan, P., Timmermans, K. R., and de Baar, H. J. W.: Major deviations of iron complexation during 22 days of a mesoscale iron enrichment in the open Southern Ocean, *Mar. Chem.*, 96, 257–271, 2005.
- Brand, L. E., Sunda, W. G., and Guillard, R. R. L.: Limitation of marine-phytoplankton reproductive rates by zinc, manganese, and iron *Limnol. Oceanogr.*, 28, 1182–1198, 1983.
- Breitbarth, E., Gelting, J., Walve, J., Hoffmann, L. J., Turner, D. R., Hassellöv, M., and Ingri, J.: Dissolved iron (II) in the Baltic Sea surface water and implications for cyanobacterial bloom development, *Biogeosciences*, 6, 2397–2420, 2009, <http://www.biogeosciences.net/6/2397/2009/>.
- Breitbarth, E., Bellerby, R. J., Neill, C. C., Ardelan, M. V., Meyerhöfer, M., Zöllner, E., Croot, P. L., and Riebesell, U.: Ocean acidification affects iron speciation during a coastal seawater mesocosm experiment, *Biogeosciences*, 7, 1065–1073, 2010, <http://www.biogeosciences.net/7/1065/2010/>.
- Bruland, K. W., Rue, E. L., and Smith, G. J.: Iron and macronutrients in California coastal upwelling regimes: Implications for diatom blooms, *Limnol. Oceanogr.*, 46, 1661–1674, 2001.
- Bucciarelli, E., Pondaven, P., and Sarthou, G.: Effects of an iron-light co-limitation on the elemental composition (Si, C, N) of the marine diatoms *Thalassiosira oceanica* and *Ditylum brightwellii*, *Biogeosciences*, 7, 657–669, 2010, <http://www.biogeosciences.net/7/657/2010/>.
- Buck, K. N., Lohan, M. C., Berger, C. J. M., and Bruland, K. W.: Dissolved iron speciation in two distinct river plumes and an estuary: Implications for riverine iron supply, *Limnol. Oceanogr.*, 52, 843–855, 2007.
- Buesseler, K. O., Bauer, J. E., Chen, R. F., Eglinton, T. I., Gustafsson, O., Landing, W., Mopper, K., Moran, S. B., Santschi, P. H., VernonClark, R., and Wells, M. L.: An intercomparison of cross-flow filtration techniques used for sampling marine colloids: Overview and organic carbon results, *Mar. Chem.*, 55, 1–31, 1996.
- Buesseler, K. O., Andrews, J. E., Pike, S. M., and Charette, M. A.: The Effects of Iron Fertilization on Carbon Sequestration in the Southern Ocean, *Science*, 304, 414–417, 2004.
- Buesseler, K. O., Doney, S. C., Karl, D. M., Boyd, P. W., Caldeira, K., Chai, F., Coale, K. H., de Baar, H. J. W., Falkowski, P. G., Johnson, K. S., Lampitt, R. S., Michaels, A. F., Naqvi, S. W. A., Smetacek, V., Takeda, S., and Watson, A. J.: Ocean Iron Fertilization – Moving Forward in a Sea of Uncertainty, *Science*, 319, 162, doi:10.1126/science.1154305, 2008.
- Caldeira, K. and Wickett, M. E.: Anthropogenic carbon and ocean pH, *Nature*, 425, 365–365, 2003.
- Chase, Z., Johnson, K. S., Elrod, V. A., Plant, J. N., Fitzwater, S. E., Pickell, L., and Sakamoto, C. M.: Manganese and iron distributions off central California influenced by upwelling and shelf width, *Mar. Chem.*, 95, 235–254, doi:10.1016/j.marchem.2004.09.006, 2005.
- Chen, C. Y., and Durbin, E. G.: Effects on pH on the growth and carbon uptake of marine phytoplankton, *Marine Ecology Progress Series*, 109, 83–94, 1994.
- Chen, M. and Wang, W. X.: Bioavailability of natural colloid-bound iron to marine plankton: Influences of colloidal size and aging, *Limnol. Oceanogr.*, 46, 1956–1967, 2001.
- Chever, F., Sarthou, G., Bucciarelli, E., Blain, S., and Bowie, A. R.: An iron budget during the natural iron fertilisation experiment KEOPS (Kerguelen Islands, Southern Ocean), *Biogeosciences*, 7, 455–468, 2010,

- <http://www.biogeosciences.net/7/455/2010/>.
- Chin, W.-C., Orellana, M. V., and Verdugo, P.: Spontaneous assembly of marine dissolved organic matter into polymer gels, *Nature*, 391, 568–572, 1998.
- Chisholm, S. W., Falkowski, P. G., and Cullen, J. J.: Oceans – Discrediting ocean fertilization, *Science*, 294, 309–310, 2001.
- Coale, K., Johnson, K., Fitzwater, S., Gordon, R., Tanner, S., Chavez, F., Ferioli, L., Sakamoto, C., Rogers, P., Millero, F., Steinberg, P., Nightingale, P., Cooper, D., Cochlan, W., Landry, M., Constantinou, J., Rollwagen, G., Trasvina, A., and Kudela, R.: A massive phytoplankton bloom induced by an ecosystem-scale iron fertilization experiment in the equatorial Pacific Ocean, *Nature*, 383, 495–501, 1996.
- Coale, K. H., Johnson, K. S., Chavez, F. P., Buesseler, K. O., Barber, R. T., Brzezinski, M. A., Cochlan, W. P., Millero, F. J., Falkowski, P. G., Bauer, J. E., Wanninkhof, R. H., Kudela, R. M., Altabet, M. A., Hales, B. E., Takahashi, T., Landry, M. R., Bidigare, R. R., Wang, X., Chase, Z., Strutton, P. G., Friederich, G. E., Gorbunov, M. Y., Lance, V. P., Hilting, A. K., Hiscock, M. R., Demarest, M., Hiscock, W. T., Sullivan, K. F., Tanner, S. J., Gordon, R. M., Hunter, C. N., Elrod, V. A., Fitzwater, S. E., Jones, J. L., Tozzi, S., Koblizek, M., Roberts, A. E., Herndon, J., Brewster, J., Ladizinsky, N., Smith, G., Cooper, D., Timothy, D., Brown, S. L., Selph, K. E., Sheridan, C. C., Twining, B. S., and Johnson, Z. I.: Southern Ocean Iron Enrichment Experiment: Carbon Cycling in High- and Low-Si Waters, *Science*, 304, 408–414, 2004.
- Croot, P. L., and Johansson, M.: Determination of iron speciation by cathodic stripping voltammetry in seawater using the competing ligand 2-(2-thiazolylazo)-p-cresol (TAC), *Electroanalysis*, 12, 565–576, 2000.
- Croot, P. L., Bowie, A. R., Frew, R. D., Maldonado, M. T., Hall, J. A., Safi, K. A., La Roche, J., Boyd, P. W., and Law, C. S.: Retention of dissolved iron and Fe-II in an iron induced Southern Ocean phytoplankton bloom, *Geophys. Res. Lett.*, 28, 3425–3428, 2001.
- Croot, P. L. and Laan, P.: Continuous shipboard determination of Fe(II) in polar waters using flow injection analysis with chemiluminescence detection, *Analytica Chimica Acta*, 466, 261–273, 2002.
- Croot, P. L., Andersson, K., Öztürk, M., and Turner, D. R.: The distribution and speciation of iron along 6° E in the Southern Ocean, *Deep Sea Res.*, 51, 2857–2879, 2004.
- Croot, P. L., Laan, P., Nishioka, J., Strass, V., Cisewski, B., Boye, M., Timmermans, K. R., Bellerby, R. G., Goldson, L., Nightingale, P., and de Baar, H. J. W.: Spatial and temporal distribution of Fe(II) and H₂O₂ during EisenEx, an open ocean mesoscale iron enrichment, *Mar. Chem.*, 95, 65–88, 2005.
- Croot, P. L., Bluhm, K., Schlosser, C., Streu, P., Breitbarth, E., Frew, R. D., and Ardelan, M. V.: Cycling of Fe(II) in Southern Ocean Iron Mesoscale Enrichment Experiments: EIFEX and SOFEX, *Geophys. Res. Lett.*, 35, L19606. doi:10.1029/12008GL035063, 2008.
- Cullen, J. J. and Boyd, P. W.: Predicting and verifying the intended and unintended consequences of large-scale ocean iron fertilization, *Mar. Ecol. Progress Ser.*, 364, 295–301, doi:10.3354/meps07551, 2008.
- Cullen, J. T., Chase, Z., Coale, K. H., Fitzwater, S. E., and Sherrell, R. M.: Effect of iron limitation on the cadmium to phosphorus ratio of natural phytoplankton assemblages from the Southern Ocean, *Limnol. Oceanogr.*, 48, 1079–1087, 2003.
- Dalbec, A. A. and Twining, B. S.: Remineralization of bioavailable iron by a heterotrophic dinoflagellate, *Aquatic Microbial Ecol.*, 54, 279–290, doi:10.3354/ame01270, 2009.
- de Baar, H. J. W., Buma, A. G. J., Nolting, R. F., Cadée, G. C., Jacques, G., and Tréguer, P. J.: On iron limitation of the Southern Ocean: experimental observations in the Weddell and Scotia Seas, *Mar. Ecol. Progr. Series*, 65, 105–122, 1990.
- de Baar, H. J. W., Dejong, J. T. M., Bakker, D. C. E., Loscher, B. M., Veth, C., Bathmann, U., and Smetacek, V.: Importance of iron for plankton blooms and carbon dioxide drawdown in the Southern Ocean, *Nature*, 373, 412–415, 1995.
- de Baar, H. J. W., and La Roche, J.: Trace Metals in the Oceans: Evolution, Biology and Global Change, in: *Marine Science Frontiers for Europe*, edited by: Wefer, G., Lamy, F., and Mantoura, F., Springer Verlag, Berlin, 79–105, 2003.
- de Baar, H. J. W., Boyd, P. W., Coale, K. H., Landry, M. R., Tsuda, A., Assmy, P., Bakker, D. C. E., Bozec, Y., Barber, R. T., Brzezinski, M. A., Buesseler, K. O., Boye, M., Croot, P. L., Gervais, F., Gorbunov, M. Y., Harrison, P. J., Hiscock, W. T., Laan, P., Lancelot, C., Law, C. S., Levasseur, M., Marchetti, A., Millero, F. J., Nishioka, J., Nojiri, Y., van Oijen, T., Riebesell, U., Rijkenberg, M. J. A., Saito, H., Takeda, S., Timmermans, K. R., Veldhuis, M. J. W., Waite, A. M., and Wong, C. S.: Synthesis of iron fertilization experiments: From the Iron Age in the Age of Enlightenment, *J. Geophys. Res.*, 110, C09S16, doi:10.1029/2004JC002601, 2005.
- de Baar, H. J. W., Gerringa, L. J. A., Laan, P., and Timmermans, K. R.: Efficiency of carbon removal per added iron in ocean iron fertilization, *Mar. Ecol. Progr. Series*, 364, 269–282, doi:10.3354/meps07548, 2008.
- de Jong, J., Schoemann, V., Tison, J. L., Becquevort, S., Masson, F., Lannuzel, D., Petit, J., Chou, L., Weis, D., and Mattioli, N.: Precise measurement of Fe isotopes in marine samples by multi-collector inductively coupled plasma mass spectrometry (MC-ICP-MS), *Analytica Chimica Acta*, 589, 105–119, doi:10.1016/j.aca.2007.02.055, 2007.
- Decho, A. W.: Microbial exopolymer secretions in ocean environments – their role(s) in food webs and marine processes, *Oceanogr. Mar. Biol.*, 28, 73–153, 1990.
- Duce, R. A., Liss, P. S., Merrill, J. T., Atlas, E. L., Buat-Menard, P., Hicks, B. B., Miller, J. M., Prospero, J. M., Arimoto, R., Church, T. M., Ellis, W., Galloway, J. N., Hansen, L., Jickells, T. D., Knap, A. H., Reinhardt, K. H., Schneider, B., Soudine, A., Tokos, J. J., Tsunogai, S., Wollast, R., and Zhou, M.: The atmospheric input of trace species to the world ocean, *Global Biogeochem. Cy.*, 5, 193–259, 1991.
- Duce, R. A., LaRoche, J., Altieri, K., Arrigo, K. R., Baker, A. R., Capone, D. G., Cornell, S., Dentener, F., Galloway, J., Ganeshram, R. S., Geider, R. J., Jickells, T., Kuypers, M. M., Langlois, R., Liss, P. S., Liu, S. M., Middelburg, J. J., Moore, C. M., Nickovic, S., Oschlies, A., Pedersen, T., Prospero, J., Schlitzer, R., Seitzinger, S., Sorensen, L. L., Uematsu, M., Ulloa, O., Voss, M., Ward, B., and Zamora, L.: Impacts of atmospheric anthropogenic nitrogen on the open ocean, *Science*, 320, 893–897, doi:10.1126/science.1150369, 2008.
- Duggen, S., Croot, P., Schacht, U., and Hoffmann, L.: Subduction zone volcanic ash can fertilize the surface ocean and stim-

- ulate phytoplankton growth: Evidence from biogeochemical experiments and satellite data, *Geophys. Res. Lett.*, 34, L01612, doi:10.1029/2006GL027522, 2007.
- Duggen, S., Olgun, N., Croot, P., Hoffmann, L., Dietze, H., Delmelle, P., and Teschner, C.: The role of airborne volcanic ash for the surface ocean biogeochemical iron-cycle: a review, *Biogeosciences*, 7, 827–844, 2010, <http://www.biogeosciences.net/7/827/2010/>.
- Dupont, C. L., Yang, S., Palenik, B., and Bourne, P. E.: Modern proteomes contain putative imprints of ancient shifts in trace metal geochemistry, *Proc. Natl. Acad. Sci. USA*, 103, 17822–17827, doi:10.1073/pnas.0605798103, 2006.
- Dutkiewicz, S., Follows, M. J., and Parekh, P.: Interactions of the iron and phosphorus cycles: A three-dimensional model study, *Global Biogeochem. Cy.*, 19, GB1021, doi:10.1029/2004gb002342, 2005.
- Editorial: The Law of the Sea, *Nature Geosci.*, 2, 153–153, doi:10.1038/ngeo464, 2009.
- Edwards, R. and Sedwick, P.: Iron in East Antarctic snow: Implications for atmospheric iron deposition and algal production in Antarctic waters, *Geophys. Res. Lett.*, 28, 3907–3910, 2001.
- Elrod, V. A., Berelson, W. M., Coale, K. H., and Johnson, K. S.: The flux of iron from continental shelf sediments: A missing source for global budgets, *Geophys. Res. Lett.*, 31(4), L12307, doi:10.1029/2004gl020216, 2004.
- Engel, A., Thoms, S., Riebesell, U., Rochelle-Newall, E., and Zondervan, I.: Polysaccharide aggregation as a potential sink of marine dissolved organic carbon, *Nature*, 428, 929–932, 2004.
- Fan, S.-M.: Photochemical and biochemical controls on reactive oxygen and iron speciation in the pelagic surface ocean, *Mar. Chem.*, 109, 152–164, 2008.
- Fan, S. M., Moxim, W. J., and Levy, H.: Aeolian input of bioavailable iron to the ocean, *Geophys. Res. Lett.*, 33, L07602, doi:10.1029/2005gl024852, 2006.
- Feely, R. A., Sabine, C. L., Lee, K., Berelson, W., Kleypas, J., Fabry, V. J., and Millero, F. J.: Impact of Anthropogenic CO₂ on the CaCO₃ System in the Oceans, *Science*, 305, 362–366, doi:10.1126/science.1097329, 2004.
- Firme, G. F., Rue, E. L., Weeks, D. A., Bruland, K. W., and Hutchins, D. A.: Spatial and temporal variability in phytoplankton iron limitation along the California coast and consequences for Si, N, and C biogeochemistry, *Global Biogeochem. Cy.*, 17(13), 1016, doi:10.1029/2001gb001824, 2003.
- Floge, S. A. and Wells, M. L.: Variation in colloidal chromophoric dissolved organic matter in the Damariscotta Estuary, Maine, *Limnol. Oceanogr.*, 52, 32–45, 2007.
- Foster, P. L. and Morel, F. M. M.: Reversal of cadmium toxicity in a diatom: An interaction between cadmium activity and iron, *Limnol. Oceanogr.*, 27, 745–752, 1982.
- Franck, V. M., Bruland, K. W., Hutchins, D. A., and Brzezinski, M. A.: Iron and zinc effects on silicic acid and nitrate uptake kinetics in three high-nutrient, low-chlorophyll (HNLC) regions, *Mar. Ecol.-Prog. Ser.*, 252, 15–33, 2003.
- Frew, R. D., Hutchins, D. A., Nodder, S., Sañudo-Wilhelmy, S., Tovar-Sanchez, A., Leblanc, K., Hare, C. E., and Boyd, P. W.: Particulate iron dynamics during FeCycle in subantarctic waters southeast of New Zealand, *Global Biogeochemical Cycles*, 20, GB1S93, doi:10.1029/2005GB002558, 2006.
- Frogner, P., Gislason, S. R., and Oskarsson, N.: Fertilizing potential of volcanic ash in ocean surface water, *Geology*, 29, 487–490, 2001.
- Fu, F.-X., Mulholland, M. R., Garcia, N. S., Beck, A., Bernhardt, P. W., Warner, M. E., Sañudo-Wilhelmy, S. A., and Hutchins, D. A.: Interactions between changing pCO₂, N₂ fixation, and Fe limitation in the marine unicellular cyanobacterium *Crocosphaera*, *Limnol. Oceanogr.*, 53, 2472–2484, 2008.
- Furman, J. A. and Capone, D. G.: Possible biogeochemical consequences of ocean fertilization, *Limnol. Oceanogr.*, 36, 1951–1959, 1991.
- Gelting, J., Breitbarth, E., Stolpe, B., Hassellöv, M., and Ingri, J.: Fractionation of iron species and iron isotopes in the Baltic Sea euphotic zone, *Biogeosciences Discuss.*, 6, 6491–6537, 2009, <http://www.biogeosciences-discuss.net/6/6491/2009/>.
- Gerringa, L. J. A., Rijkenberg, M. J. A., Wolterbeek, H. T., Verburg, T. G., Boye, M., and de Baar, H. J. W.: Kinetic study reveals weak Fe-binding ligand, which affects the solubility of Fe in the Scheldt estuary, *Mar. Chem.*, 103, 30–45, 2007.
- Gervais, F., Riebesell, U., and Gorbunov, M. Y.: Changes in primary productivity and chlorophyll a in response to iron fertilization in the Southern Polar Frontal Zone, *Limnol. Oceanogr.*, 47, 1324–1335, 2002.
- Ginoux, P., Prospero, J. M., Torres, O., and Chin, M.: Long-term simulation of global dust distribution with the GOCART model: correlation with North Atlantic Oscillation, *Environmental Modelling & Software*, 19, 113–128, doi:10.1016/s1364-8152(03)00114-2, 2004.
- Gledhill, M. and van den Berg, C. M. G.: Determination of complexation of Fe(III) with natural organic complexing ligands in seawater using cathodic stripping voltammetry, *Mar. Chem.*, 47, 41–54, 1994.
- Gledhill, M.: The determination of heme b in marine phyto- and bacterioplankton, *Mar. Chem.*, 103, 393–403, doi:10.1016/j.marchem.2006.10.008, 2007.
- Gnanadesikan, A., Sarmiento, J. L., and Slater, R. D.: Effects of patchy ocean fertilization on atmospheric carbon dioxide and biological production, *Global Biogeochem. Cy.*, 17(17), 1050, doi:10.1029/2002gb001940, 2003.
- Gonzalez-Davila, M., Santana-Casiano, J. M., and Millero, F. J.: Competition between O₂ and H₂O₂ in the oxidation of Fe(II) in natural waters, *J. Solut. Chem.*, 35, 95–111, 2006.
- Guieu, C., Bonnet, S., Wagener, T., and Loye-Pilot, M. D.: Biomass burning as a source of dissolved iron to the open ocean?, *Geophys. Res. Lett.*, 32, L19608, doi:10.1029/2005GL022962, 2005.
- Gustafsson, O., Widerlund, A., Andersson, P. S., Ingri, J., Roos, P., and Ledin, A.: Colloid dynamics and transport of major elements through a boreal river – brackish bay mixing zone, *Mar. Chem.*, 71, 1–21, 2000.
- Han, Q., Moore, J. K., Zender, C., Measures, C., and Hydes, D.: Constraining oceanic dust deposition using surface ocean dissolved Al, *Global Biogeochem. Cy.*, 22, GB2003, doi:10.1029/2007gb002975, 2008.
- Hansard, S. P., Landing, W. M., Measures, C. I., and Voelker, B. M.: Dissolved iron(II) in the Pacific Ocean: Measurements from PO₂ and P16N Clivar/CO₂ repeat hydrography expeditions, *Deep Sea Res. I*, 56(7), 1117–1129, 2009.
- Hare, C. E., DiTullio, G. R., Trick, C. G., Wilhelm, S. W., Bruland, K. W., Rue, E. L., and Hutchins, D. A.: Phytoplankton commu-

- nity structure changes following simulated upwelled iron inputs in the Peru upwelling region, *Aquatic Microbial Ecology*, 38, 269–282, 2005.
- Hare, C. E., DiTullio, G. R., Riseman, S. F., Crossley, A. C., Popels, L. C., Sedwick, P. N., and Hutchins, D. A.: Effects of changing continuous iron input rates on a Southern Ocean algal assemblage, *Deep-Sea Research Part I-Oceanographic Research Papers*, 54, 732–746, doi:10.1016/j.dsr.2007.02.001, 2007a.
- Hare, C. E., Leblanc, K., DiTullio, G. R., Kudela, R. M., Zhang, Y., Lee, P. A., Riseman, S., and Hutchins, D. A.: Consequences of increased temperature and CO₂ for phytoplankton community structure in the Bering Sea, *Marine Ecology Progress Series*, 352, 9–16, doi:10.3354/meps07182, 2007b.
- Hassellöv, M. and von der Kammer, F.: Iron Oxides as Geochemical Nanovectors for Metal Transport in Soil-River Systems, *Elements*, 4, 401–406, doi:10.2113/gselements.4.6.401, 2008.
- Hassler, C. S., Slaveykova, V. I., and Wilkinson, K. J.: Discriminating between intra- and extracellular metals using chemical extractions, *Limnol. Oceanogr. Methods*, 2, 237–247, 2004.
- Hassler, C. S., Twiss, M. R., McKay, R. M. L., and Bullerjahn, G. S.: Optimization of iron-dependent cyanobacterial (*Synechococcus*, cyanophyceae) bioreporters to measure iron bioavailability, *J. Phycol.*, 42, 324–335, 2006.
- Hassler, C. S., Chafin, R. D., Klinger, M. B., and Twiss, M. R.: Application of the biotic ligand model to explain potassium interaction with thallium uptake and toxicity to plankton, *Environ. Toxicol. Chem.*, 26, 1139–1145, 2007.
- Hassler, C. S. and Schoemann, V.: Bioavailability of organically bound Fe to model phytoplankton of the Southern Ocean, *Biogeosciences*, 6, 2281–2296, 2009, <http://www.biogeosciences.net/6/2281/2009/>.
- Heldal, M., Fagerbakke, K. M., Tuomi, P., and Bratbak, G.: Abundant populations of iron and manganese sequestering bacteria in coastal water, *Aquatic Microbial Ecology*, 11, 127–133, 1996.
- Ho, T. Y., Quigg, A., Finkel, Z. V., Milligan, A. J., Wyman, K., Falkowski, P. G., and Morel, F. M. M.: The elemental composition of some marine phytoplankton, *J. Phycol.*, 39, 1145–1159, 2003.
- Hoagland, K. D., Rosowski, J. R., Gretz, M. R., and Roemer, S. C.: Diatom extracellular polymeric substances – function, fine-structure, chemistry, and physiology, *J. Phycol.*, 29, 537–566, 1993.
- Hoffmann, L. J., Peeken, I., and Lochte, K.: Iron, silicate, and light co-limitation of three Southern Ocean diatom species, *Polar Biol.*, 31, 1067–1080, doi:10.1007/s00300-008-0448-6, 2008.
- Holm-Hansen, O., Mitchell, B. G., Hewes, C. D., and Karl, D. M.: Phytoplankton blooms in the vicinity of Palmer Station, Antarctica, *Polar Biology*, 10, 49–57, 1989.
- Hopkinson, B. M. and Barbeau, K. A.: Organic and redox speciation of iron in the eastern tropical North Pacific suboxic zone, *Mar. Chem.*, 106, 2–17, 2007.
- Hopkinson, B. M., Roe, K. L., and Barbeau, K. A.: Heme uptake by *Microscilla marina* and evidence for heme uptake systems in the genomes of diverse marine bacteria, *Appl. Environ. Microbiol.*, 74, 6263–6270, doi:10.1128/aem.00964-08, 2008.
- Howell, K. A., Achterberg, E. P., Tappin, A. D., and Worsfold, P. J.: Colloidal metals in the tamar estuary and their influence on metal fractionation by membrane filtration, *Environ. Chem.*, 3, 199–207, doi:10.1071/en06004, 2006.
- Hudson, R. J. M. and Morel, F. M. M.: Iron transport in marine phytoplankton – kinetics of cellular and medium coordination reactions, *Limnol. Oceanogr.*, 35, 1002–1020, 1990.
- Hunter, K. A. and Boyd, P. W.: Iron-binding ligands and their role in the ocean biogeochemistry of iron, *Environ. Chem.*, 4, 221–232, doi:10.1071/en07012, 2007.
- Hurst, M. P. and Bruland, K. W.: An investigation into the exchange of iron and zinc between soluble, colloidal, and particulate size-fractions in shelf waters using low-abundance isotopes as tracers in shipboard incubation experiments, *Mar. Chem.*, 103, 211–226, doi:10.1016/j.marchem.2006.07.001, 2007.
- Hutchins, D. A. and Bruland, K. W.: Grazer-mediated regeneration and assimilation of Fe, Zn and Mn from planktonic prey, *Mar. Ecol. Progr. Series*, 110, 259–269, 1994.
- Hutchins, D. A., Wang, W. X., and Fisher, N. S.: Copepod grazing and the biogeochemical fate of diatom iron, *Limnol. Oceanogr.*, 40, 989–994, 1995.
- Hutchins, D. A. and Bruland, K. W.: Iron-limited diatom growth and Si:N uptake ratios in a coastal upwelling regime, *Nature*, 393, 561–564, 1998.
- Hutchins, D. A., Franck, V. M., Brzezinski, M. A., and Bruland, K. W.: Inducing phytoplankton iron limitation in iron-replete coastal waters with a strong chelating ligand, *Limnol. Oceanogr.*, 44, 1009–1018, 1999a.
- Hutchins, D. A., Witter, A. E., Butler, A., and Luther III, G. W.: Competition among marine phytoplankton for different chelated iron species, *Nature*, 400, 858–861, 1999b.
- Hutchins, D. A., Pustizzi, F., Hare, C. E., and DiTullio, G. R.: A shipboard natural community continuous culture system for ecologically relevant low-level nutrient enrichment experiments, *Limnol. Oceanogr.-Methods*, 1, 82–91, 2003.
- Ingri, J., Malinovsky, D., Rodushkin, I., Baxter, D. C., Widerlund, A., Andersson, P., Gustafsson, O., Forsling, W., and Ohlander, B.: Iron isotope fractionation in river colloidal matter, *Earth Planet. Sci. Lett.*, 245, 792–798, 2006.
- Jacobson, M. Z.: Studying ocean acidification with conservative, stable numerical schemes for nonequilibrium air-ocean exchange and ocean equilibrium chemistry, *J. Geophys. Res.-Atmos.*, 110, D07302, doi:10.01029/2004JD005220, 2005.
- Jickells, T. and Spokes, L. J.: Atmospheric iron inputs into the oceans, in: *The Biogeochemistry of Iron in Seawater*, edited by: Turner, D. R., and Hunter, K. A., John Wiley and Sons Ltd., West Sussex, England, 85–121, 2001.
- Jickells, T. D., Dorling, S., Deuser, W. G., Church, T. M., Arimoto, R., and Prospero, J. M.: Air-borne dust fluxes to a deep water sediment trap in the Sargasso Sea, *Global Biogeochem. Cy.*, 12, 311–320, 1998.
- Jickells, T. D., An, Z. S., Andersen, K. K., Baker, A. R., Bergametti, G., Brooks, N., Cao, J. J., Boyd, P. W., Duce, R. A., Hunter, K. A., Kawahata, H., Kubilay, N., laRoche, J., Liss, P. S., Mahowald, N., Prospero, J. M., Ridgwell, A. J., Tegen, I., and Torres, R.: Global Iron Connections Between Desert Dust, Ocean Biogeochemistry, and Climate, *Science*, 308, 67–71, 2005.
- Johnson, K. S.: Iron supply and demand in the upper ocean: Is extraterrestrial dust a significant source of bioavailable iron?, *Glob. Biogeochem. Cy.*, 15, 61–63, 2001.
- Johnson, K. S., Boyle, E., Bruland, K. W., Coale, K., Measures, C., Moffett, J., Aguilar-Islas, A., Barbeau, K., Bergquist, B., Bowie, A., Buck, K., Cai, Y., Chase, Z., Cullen, J., Doi, T., Elrod, V.,

- Fitzwater, S., Gordon, M., King, A., Laan, P., Laglera-Baquer, L., Landing, W., Lohan, M., Mendez, J., Milne, A., Obata, H., Osslander, L., Plant, J., Sarthou, G., Sedwick, P., Smith, G. J., Sohst, B., Tanner, S., van den Berg, S., and Wu, J.: The SAFE Iron Intercomparison Cruise: An International Collaboration to Develop Dissolved Iron in Seawater Standards, *EOS, Transactions of the American Geophysical Union*, 88, 131–132, 2007.
- Johnson, W. K., Miller, L. A., Sutherland, N. E., and Wong, C. S.: Iron transport by mesoscale Haida eddies in the Gulf of Alaska, *Deep-Sea Res.*, 52, 933–953, doi:10.1016/j.dsr.2.2004.08.017, 2005.
- Jones, M. T. and Gislason, S. R.: Rapid releases of metal salts and nutrients following the deposition of volcanic ash into aqueous environments, *Geochim. Cosmochim. Acta*, 72, 3661–3680, 2008.
- Journet, E., Desboeufs, K. V., Caquineau, S., and Colin, J. L.: Mineralogy as a critical factor of dust iron solubility, *Geophys. Res. Lett.*, 35, L07805, doi:10.1029/2007gl031589, 2008.
- Kintisch, E.: Carbon sequestration: Should oceanographers pump iron?, *Science*, 318, 1368–1370, 2007.
- Kondo, Y., Takeda, S., Nishioka, J., Obata, H., Furuya, K., Johnson, W. K., and Wong, C. S.: Organic iron(III) complexing ligands during an iron enrichment experiment in the western subarctic North Pacific, *Geophys. Res. Lett.*, 35, L12601, doi:10.1029/2008gl033354, 2008.
- Krachler, R., Jirsa, F., and Ayromlou, S.: Factors influencing the dissolved iron input by river water to the open ocean, *Biogeosciences*, 2, 311–315, 2005, <http://www.biogeosciences.net/2/311/2005/>.
- Lacan, F., Radic, A., Jeandel, C., Poitrasson, F., Sarthou, G., Pradoux, C., and Freyrier, R.: Measurement of the isotopic composition of dissolved iron in the open ocean, *Geophys. Res. Lett.*, 35, L24610, doi:10.1029/2008gl035841, 2008.
- Laës, A., Blain, S., Laan, P., Achterberg, E. P., Sarthou, G., and de Baar, H. J. W.: Deep dissolved iron profiles in the eastern North Atlantic in relation to water masses, *Geophys. Res. Lett.*, 30(4), 1902, doi:10.1029/2003gl017902, 2003.
- Laës, A., Blain, S., Laan, P., Ussher, S. J., Achterberg, E. P., Trguer, P., and de Baar, H. J. W.: Sources and transport of dissolved iron and manganese along the continental margin of the Bay of Biscay, *Biogeosciences*, 4, 181–194, 2007, <http://www.biogeosciences.net/4/181/2007/>.
- Laglera, L. M. and van den Berg, C. M. G.: Evidence for geochemical control of iron by humic substances in seawater, *Limnol. Oceanogr.*, 54, 610–619, 2009.
- Lam, C. K. S. C., Jickells, T. D., Richardson, D. J., and Russell, D. A.: Fluorescence-Based Siderophore Biosensor for the Determination of Bioavailable Iron in Oceanic Waters, *Anal. Chem.*, 78, 5040–5045, doi:10.1021/ac060223t, 2006.
- Lam, P. J. and Bishop, J. K. B.: The continental margin is a key source of iron to the HNLC North Pacific Ocean, *Geophys. Res. Lett.*, 35(5), L07608, doi:10.1029/2008gl033294, 2008.
- Lancelot, C., de Montety, A., Goosse, H., Becquevort, S., Schoemann, V., Pasquer, B., and Vancoppenolle, M.: Spatial distribution of the iron supply to phytoplankton in the Southern Ocean: a model study, *Biogeosciences*, 6, 2861–2878, 2009, <http://www.biogeosciences.net/6/2861/2009/>.
- Lannuzel, D., Schoemann, V., de Jong, J., Tison, J. L., and Chou, L.: Distribution and biogeochemical behaviour of iron in the East Antarctic sea ice, *Mar. Chem.*, 106, 18–32, doi:10.1016/j.marchem.2006.06.010, 2007.
- Lannuzel, D., Schoemann, V., de Jong, J., Chou, L., Delille, B., Becquevort, S., and Tison, J.-L.: Iron study during a time series in the western Weddell pack ice, *Mar. Chem.*, 108, 85–95, 2008.
- Liss, P., Chuck, A., Bakker, D., and Turner, S.: Ocean fertilization with iron: effects on climate and air quality, *Tellus Ser. B-Chem. Phys. Meteorol.*, 57, 269–271, 2005.
- Liu, X. W. and Millero, F. J.: The solubility of iron hydroxide in sodium chloride solutions, *Geochim. Cosmochim. Acta*, 63, 3487–3497, 1999.
- Liu, X. W. and Millero, F. J.: The solubility of iron in seawater, *Mar. Chem.*, 77, 43–54, 2002.
- Lohan, M. C. and Bruland, K. W.: Importance of vertical mixing for additional sources of nitrate and iron to surface waters of the Columbia River plume: Implications for biology, *Mar. Chem.*, 98, 260–273, doi:10.1016/j.marchem.2005.10.003, 2006.
- Loscher, B. M., DeBaar, H. J. W., DeJong, J. T. M., Veth, C., and Dehairs, F.: The distribution of Fe in the Antarctic Circumpolar Current, *Deep-Sea Res.*, 44, 143–187, 1997.
- Luo, C., Mahowald, N., Bond, T., Chuang, P. Y., Artaxo, P., Siefert, R., Chen, Y., and Schauer, J.: Combustion iron distribution and deposition, *Global Biogeochem. Cy.*, 22(17), GB1012, doi:10.1029/2007gb002964, 2008.
- Lyvén, B., Hasselov, M., Turner, D. R., Haraldsson, C., and Andersson, K.: Competition between iron- and carbon-based colloidal carriers for trace metals in a freshwater assessed using flow field-flow fractionation coupled to ICPMS, *Geochim. Cosmochim. Acta*, 67, 3791–3802, 2003.
- Mackey, D. J., O’Sullivan, J. E., and Watson, R. J.: Iron in the western Pacific: a riverine or hydrothermal source for iron in the Equatorial Undercurrent?, *Deep-Sea Res.*, 49, 877–893, 2002.
- Mahowald, N., Jickells, T. D., Baker, A. R., Artaxo, P., Benitez-Nelson, C. R., Bergametti, G., Bond, T. C., Chen, Y., Cohen, D. D., Herut, B., Kubilay, N., Losno, R., Luo, C., Maenhaut, W., McGee, K. A., Okin, G. S., Siefert, R. L., and Tsukuda, S.: Global distribution of atmospheric phosphorus sources, concentrations and deposition rates, and anthropogenic impacts, *Global Biogeochem. Cy.*, 22, GB4026, doi:10.1029/2008gb003240, 2008.
- Mahowald, N. M., Baker, A. R., Bergametti, G., Brooks, N., Duce, R. A., Jickells, T. D., Kubilay, N., Prospero, J. M., and Tegen, I.: Atmospheric global dust cycle and iron inputs to the ocean, *Global Biogeochem. Cy.*, 19, GB4025, doi:10.1029/2004gb002402, 2005.
- Maldonado, M. T., Boyd, P. W., Harrison, P. J., and Price, N. M.: Co-limitation of phytoplankton growth by light and Fe during winter in the NE subarctic Pacific Ocean, *Deep-Sea Res.*, 46, 2475–2485, 1999.
- Maldonado, M. T., Strzepek, R. F., Sander, S., and Boyd, P. W.: Acquisition of iron bound to strong organic complexes, with different Fe binding groups and photochemical reactivities, by plankton communities in Fe-limited subantarctic waters, *Global Biogeochem.*, 19, GB4S23, doi:10.1029/2005GB002481, 2005.
- Marchetti, A., Parker, M. S., Moccia, L. P., Lin, E. O., Arieta, A. L., Ribalet, F., Murphy, M. E. P., Maldonado, M. T., and Armbrust, E. V.: Ferritin is used for iron storage in bloom-forming marine pennate diatoms, *Nature*, 457, 467–470, doi:10.1038/nature07539, 2009.

- Martin, J. H.: Glacial-interglacial CO₂ change: the iron hypothesis, *Paleoceanography*, 5, 1–13, 1990.
- Mawji, E., Gledhill, M., Milton, J. A., Tarran, G. A., Ussher, S., Thompson, A., Wolff, G. A., Worsfold, P. J., and Achterberg, E. P.: Hydroxamate Siderophores: Occurrence and Importance in the Atlantic Ocean, *Environ. Sci. Technol.*, 42, 8675–8680, doi:10.1021/es801884r, 2008.
- McKay, R., Michael, L., la Roche, J., Yakunin, A., F. Durnford, D. G., and Geider, R. J.: Accumulation of Ferredoxin and Flavodoxin in a Marine Diatom in response to Fe, *J. Phycology*, 35, 510–519, 1999.
- Millero, F. J., Sotolongo, S., and Izaguirre, M.: The oxidation kinetics of Fe(II) in seawater, *Geochim. Cosmochim. Acta*, 51, 793–801, 1987.
- Millero, F. J. and Sotolongo, S.: The oxidation of Fe(II) with H₂O₂ in seawater, *Geochim. Cosmochim. Acta*, 53, 1867–1873, 1989.
- Mills, M. M., Ridame, C., Davey, M., La Roche, J., and Geider, R. J.: Iron and phosphorus co-limits nitrogen fixation in the eastern tropical North Atlantic, *Nature*, 429, 292–294, 2004.
- Moore, C. M., Mills, M. M., Milne, A., Langlois, R., Achterberg, E. P., Lochte, K., Geider, R. J., and La Roche, J.: Iron limits primary productivity during spring bloom development in the central North Atlantic, *Global Change Biol.*, 12, 626–634 doi:10.1111/j.1365-2486.2006.01122.x, 2006.
- Moore, C. M., Hickman, A. E., Poulton, A. J., Seeyave, S., and Lucas, M. I.: Iron-light interactions during the CROZet natural iron bloom and EXport experiment (CROZEX): II – Taxonomic responses and elemental stoichiometry, *Deep-Sea Res.*, 54, 2066–2084, doi:10.1016/j.dsr2.2007.06.015, 2007a.
- Moore, C. M., Seeyave, S., Hickman, A. E., Allen, J. T., Lucas, M. I., Planquette, H., Pollard, R. T., and Poulton, A. J.: Iron-light interactions during the CROZet natural iron bloom and EXport experiment (CROZEX) I: Phytoplankton growth and photophysiology, *Deep-Sea Res.*, 54, 2045–2065, doi:10.1016/j.dsr2.2007.06.011, 2007b.
- Moran, S. B., Yeats, P. A., and Balls, P. W.: On the role of colloids in trace metal solid-solution partitioning in continental shelf waters: A comparison of model results and field data, *Cont. Shelf Res.*, 16, 397–408, 1996.
- Morel, F. M. M. and Price, N. M.: The biogeochemical cycles of trace metals in the oceans, *Science*, 300, 944–947, 2003.
- Morel, F. M. M., Kustka, A. B., and Shaked, Y.: The role of unchelated Fe in the iron nutrition of phytoplankton, *Limnol. Oceanogr.*, 53, 400–404, 2008.
- Mosley, L. M., Hunter, K. A., and Ducker, W. A.: Forces between colloid particles in natural waters, *Environ. Sci. Technol.*, 37, 3303–3308, doi:10.1021/es026216d, 2003.
- Mylon, S. E., Chen, K. L., and Elimelech, M.: Influence of natural organic matter and ionic composition on the kinetics and structure of hematite colloid aggregation: Implications to iron depletion in estuaries, *Langmuir*, 20, 9000–9006, doi:10.1021/la049153g, 2004.
- Nowostawska, U., Kim, J. P., and Hunter, K. A.: Aggregation of riverine colloidal iron in estuaries: A new kinetic study using stopped-flow mixing, *Mar. Chem.*, 110, 205–210, 2008.
- Nunn, B. L. and Timperman, A. T.: Marine Proteomics, *Mar. Ecol. Progr. Series*, 332, 281–289, 2007.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G. K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M. F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.
- Öztürk, M., Steinnes, E., and Sakshaug, E.: Iron Speciation in the Trondheim Fjord from the Perspective of Iron Limitation for Phytoplankton, *Estuarine, Coastal Shelf Sci.*, 55, 197–212, 2002.
- Öztürk, M. and Bizsel, N.: Iron speciation and biogeochemistry in different nearshore waters, *Mar. Chem.*, 83, 145–156, 2003.
- Öztürk, M., Croot, P. L., Bertilsson, S., Abrahamsson, K., Karlson, B., David, R., Fransson, A., and Sakshaug, E.: Iron enrichment and photoreduction of iron under UV and PAR in the presence of hydroxycarboxylic acid: implications for phytoplankton growth in the Southern Ocean, *Deep Sea Res.*, 51, 2841–2856, 2004.
- Paytan, A., Mackey, K. R. M., Chen, Y., Lima, I. D., Doney, S. C., Mahowald, N., Labiosa, R., and Postf, A. F.: Toxicity of atmospheric aerosols on marine phytoplankton, *Proc. Natl. Acad. Sci. USA*, 106, 4601–4605, di:10.1073/pnas.0811486106, 2009.
- Peers, G. and Price, N. M.: Copper-containing plastocyanin used for electron transport by an oceanic diatom, *Nature*, 441, 341–344, 2006.
- Pickell, L. D., Wells, M. L., Trick, C. G., and Cochlan, W. P.: A sea-going continuous culture system for investigating phytoplankton community response to macro- and micro-nutrient manipulations, *Limnol. Oceanogr.-Methods*, 7, 21–32, 2009.
- Planquette, H., Statham, P. J., Fones, G. R., Charette, M. A., Moore, C. M., Salter, I., Nedelec, F. H., Taylor, S. L., French, M., Baker, A. R., Mahowald, N., and Jickells, T. D.: Dissolved iron in the vicinity of the Crozet Islands, Southern Ocean, *Deep-Sea Res.*, 54, 1999–2019, doi:10.1016/j.dsr2.2007.06.019, 2007.
- Pollard, R. T., Salter, I., Sanders, R. J., Lucas, M. I., Moore, C. M., Mills, R. A., Statham, P. J., Allen, J. T., Baker, A. R., Bakker, D. C. E., Charette, M. A., Fielding, S., Fones, G. R., French, M., Hickman, A. E., Holland, R. J., Hughes, J. A., Jickells, T. D., Lampitt, R. S., Morris, P. J., Nedelec, F. H., Nielsdottir, M., Planquette, H., Popova, E. E., Poulton, A. J., Read, J. F., Seeyave, S., Smith, T., Stinchcombe, M., Taylor, S., Thomalla, S., Venables, H. J., Williamson, R., and Zubkov, M. V.: Southern Ocean deep-water carbon export enhanced by natural iron fertilization, *Nature*, 457, 577–580, 2009.
- Powell, R. T. and Wilson-Finelli, A.: Importance of organic Fe complexing ligands in the Mississippi River plume, *Estuarine, Coastal Shelf Sci.*, 58, 757–763, 2003.
- Prospero, J. M., Ginoux, P., Torres, O., Nicholson, S. E., and Gill, T. E.: Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product, *Rev. Geophys.*, 40, 1002, doi:10.1029/2000RG000095, 2002.
- Pulido-Villena, E., Wagener, T., and Guieu, C.: Bacterial response to dust pulses in the western Mediterranean: Implications for carbon cycling in the oligotrophic ocean, *Global Biogeochem. Cy.*, 22(12), GB1020, doi:10.1029/2007GB003091, 2008.
- Quigg, A., Finkel, Z. V., Irwin, A. J., Rosenthal, Y., Ho, T. Y., Reinfelder, J. R., Schofield, O., Morel, F. M. M., and Falkowski, P. G.: The evolutionary inheritance of elemental stoichiometry in marine phytoplankton, *Nature*, 425, 291–294, 2003.

- Raiswell, R., Tranter, M., Benning, L. G., Siebert, M., De'ath, R., Huybrechts, P., and Payne, T.: Contributions from glacially derived sediment to the global iron (oxyhydr)oxide cycle: Implications for iron delivery to the oceans, *Geochim. Cosmochim. Acta*, 70, 2765–2780, doi:10.1016/j.gca.2005.12.027, 2006.
- Raiswell, R., Benning, L. G., Tranter, M., and Tulaczyk, S.: Bioavailable iron in the Southern Ocean: the significance of the iceberg conveyor belt, *Geochim. Trans.*, 9(7), doi:10.1186/1467-4866-9-7, 2008.
- Riebesell, U., Schulz, K. G., Bellerby, R. G. J., Botros, M., Fritsche, P., Meyerhofer, M., Neill, C., Nondal, G., Oschlies, A., Wohlers, J., and Zollner, E.: Enhanced biological carbon consumption in a high CO₂ ocean, *Nature*, 450, 545–510, doi:10.1038/nature06267, 2007.
- Rose, A. L.: Effect of Dissolved Natural Organic Matter on the Kinetics of Ferrous Iron Oxygenation in Seawater, *Environ. Sci. Technol.*, 37, 4877–4886, doi:10.1021/es034152g, 2003.
- Rose, J. M., Feng, Y., DiTullio, G. R., Dunbar, R. B., Hare, C. E., Lee, P. A., Lohan, M., Long, M., W. O. Smith Jr., Sohst, B., Tozzi, S., Zhang, Y., and Hutchins, D. A.: Synergistic effects of iron and temperature on Antarctic phytoplankton and microzooplankton assemblages, *Biogeosciences*, 6, 3131–3147, 2009, <http://www.biogeosciences.net/6/3131/2009/>.
- Roy, E. G., Jiang, C. H., Wells, M. L., and Tripp, C.: Determining subnanomolar iron concentrations in oceanic seawater using a siderophore-modified film analyzed by infrared spectroscopy, *Anal. Chem.*, 80, 4689–4695, doi:10.1021/ac800356p, 2008a.
- Roy, E. G., Wells, M. L., and King, D. W.: Persistence of iron(II) in surface waters of the western subarctic Pacific, *Limnol. Oceanogr.*, 53, 89–98, 2008b.
- Rue, E. L. and Bruland, K. W.: Complexation of iron(III) by natural organic ligands in the Central North Pacific as determined by a new competitive ligand equilibration/adsorptive cathodic stripping voltammetric method, *Mar. Chem.*, 50, 117–138, 1995.
- Rue, E. L. and Bruland, K. W.: The role of organic complexation on ambient iron chemistry in the equatorial Pacific Ocean and the response of a mesoscale iron addition experiment, *Limnol. Oceanogr.*, 42, 901–910, 1997.
- Rusch, D. B., Halpern, A. L., Sutton, G., Heidelberg, K. B., Williamson, S., Yooshep, S., Wu, D., Eisen, J. A., Hoffman, J. M., Remington, K., Beeson, K., Tran, B., Smith, H., Baden-Tillson, H., Stewart, C., Thorpe, J., Freeman, J., Andrews-Pfannkoch, C., Venter, J. E., Li, K., Kravitz, S., Heidelberg, J. F., Utterback, T., Rogers, Y. H., Falcón, L. I., Souza, V., Bonilla-Rosso, G., Eguiarte, L. E., Karl, D. M., Sathyendranath, S., Platt, T., Birmingham, E., Gallardo, V., Tamayo-Castillo, G., Ferrari, M. R., Strausberg, R. L., Nealson, K., Friedman, R., Frazier, M., and Venter, J. C.: The Sorcerer II Global Ocean Sampling expedition: northwest Atlantic through eastern tropical Pacific, *PLoS biology*, 5(e77), 398–431, 2007.
- Saito, M. A., Sigman, D. M., and Morel, F. M. M.: The bioinorganic chemistry of the ancient ocean: the co-evolution of cyanobacterial metal requirements and biogeochemical cycles at the Archean-Proterozoic boundary?, *Inorganic Chimica Acta*, 356, 308–318, 2003.
- Salmon, T. P., Rose, A. L., Neilan, B. A., and Waite, T. D.: The FeL model of iron acquisition: Nondissociative reduction of ferric complexes in the marine environment, *Limnol. Oceanogr.*, 51, 1744–1754, 2006.
- Sander, S., Mosley, L. M., and Hunter, K. A.: Investigation of interparticle forces in natural waters: Effects of adsorbed humic acids on iron oxide and alumina surface properties, *Environ. Sci. Technol.*, 38, 4791–4796, doi:10.1021/es049602z, 2004.
- Sander, S., Ginon, L., Anderson, B., and Hunter, K. A.: Comparative study of organic Cd and Zn complexation in lake waters – seasonality, depth and pH dependence, *Environ. Chem.*, 4, 410–423, 2007.
- Santana-Casiano, J. M., Gonzalez-Davila, M., and Millero, F. J.: Oxidation of Nanomolar Levels of Fe(II) with Oxygen in Natural Waters, *Environ. Sci. Technol.*, 39, 2073–2079, 2005.
- Santana-Casiano, J. M., Gonzalez-Davila, M., and Millero, F. J.: The role of Fe(II) species on the oxidation of Fe(II) in natural waters in the presence of O₂ and H₂O₂, *Mar. Chem.*, 99, 70–82, 2006.
- Santschi, P. H., Balnois, E., Wilkinson, K. J., Zhang, J. W., Buffle, J., and Guo, L. D.: Fibrillar polysaccharides in marine macromolecular organic matter as imaged by atomic force microscopy and transmission electron microscopy, *Limnol. Oceanogr.*, 43, 896–908, 1998.
- Sarmiento, J. L. and Orr, J. C.: 3-dimensional simulations of the impact of Southern-Ocean nutrient depletion on atmospheric CO₂ and ocean chemistry, *Symp on What Controls Phytoplankton Production in Nutrient-Rich Areas of the Open Sea*, San Marcos, Ca, 1991, ISI:A1991HR98100032, 1928–1950.
- Sarthou, G., Timmermans, K. R., Blain, S., and Treguer, P.: Growth physiology and fate of diatoms in the ocean: a review, *J. Sea Res.*, 53, 25–42, 2005.
- Sarthou, G., Vincent, D., Christaki, U., Obernosterer, I., Timmermans, K. R., and Brussaard, C. P. D.: The fate of biogenic iron during a phytoplankton bloom induced by natural fertilisation: Impact of copepod grazing, *Deep-Sea Res.*, 55, 734–751, doi:10.1016/j.dsr.2007.12.033, 2008.
- Sato, M., Takeda, S., and Furuya, K.: Iron regeneration and organic iron(III)-binding ligand production during in situ zooplankton grazing experiment, *Mar. Chem.*, 106, 471–488, 2007.
- Schmincke, H.-U.: *Volcanism*, Springer-Verlag, Berlin Heidelberg New York, 324 pp., 2004.
- Schoemann, V., Wollast, R., Chou, L., and Lancelot, C.: Effects of photosynthesis on the accumulation of Mn and Fe by *Phaeocystis* colonies, *Limnol. Oceanogr.*, 46, 1065–1076, 2001.
- Schoemann, V., De Jong, J. T. M., Lannuzel, D., Tison, J. L., Dellile, B., Chou, L., Lancelot, C., and Becquevort, S.: Microbiological control on the cycling Fe and its isotopes in Antarctic sea ice, 8th Annual V M Goldschmidt Conference, Vancouver, CANADA, ISI:000257301602198, A837–A837, 2008.
- Schroth, A. W., Crusius, J., Sholkovitz, E. R., and Bostick, B. C.: Iron solubility driven by speciation in dust sources to the ocean, *Nature Geosci.*, 2, 337–340, 2009.
- Schulz, K. G., Zondervan, I., Gerringa, L. J. A., Timmermans, K. R., Veldhuis, M. J. W., and Riebesell, U.: Effect of trace metal availability on coccolithophorid calcification, *Nature*, 430, 673–676, 2004.
- Sedwick, P. N., Sholkovitz, E. R., and Church, T. M.: Impact of anthropogenic combustion emissions on the fractional solubility of aerosol iron: Evidence from the Sargasso Sea, *Geochim. Geophys. Geosyst.*, 8, 21, Q10q06, doi:10.1029/2007gc001586, 2007.
- Shaked, Y., Kustka, A. B., Morel, F. M. M., and Erel, Y.: Simulta-

- neous determination of iron reduction and uptake by phytoplankton, *Limnol. Oceanogr.-Methods*, 2, 137–145, 2004.
- Shaked, Y., Kustka, A. B., and Morel, F. M. M.: A general kinetic model for iron acquisition by eukaryotic phytoplankton, *Limnol. Oceanogr.*, 50, 872–882, 2005.
- Sholkovitz, E. R.: Flocculation of dissolved Fe, Mn, Al, Cu, Ni, Co, and Cd during estuarine mixing, *Earth Planet. Sci. Lett.*, 41, 77–86, 1978.
- Sholkovitz, E. R., Boyle, E. A., and Price, N. B.: Removal of dissolved humic acids and iron during estuarine mixing, *Earth Planet. Sci. Lett.*, 40, 130–136, 1978.
- Sigman, D. M. and Boyle, E. A.: Glacial/interglacial variations in atmospheric carbon dioxide, *Nature*, 407, 859–869, 2000.
- Sillen, L. G. and Martell, A. E.: *Stability Constants*, Chemical Society, London, 1971.
- Smith, K. L., Robison, B. H., Helly, J. J., Kaufmann, R. S., Ruhl, H. A., Shaw, T. J., Twining, B. S., and Vernet, M.: Free-drifting icebergs: Hot spots of chemical and biological enrichment in the Weddell Sea, *Science*, 317, 478–482, doi:10.1126/science.1142834, 2007.
- Smith, W. O. and Nelson, D. M.: Phytoplankton bloom produced by a receding ice edge in the Ross Sea – spacial coherence with the density field, *Science*, 227, 163–166, 1985.
- Spokes, L. and Jickells, T. D.: Speciation of metals in the atmosphere, in: *Chemical Speciation in the Environment*, edited by: Ure, A. and Davidson, C., Blackwell, Malden, 159–187, 2002.
- Statham, P. J., Skidmore, M., and Tranter, M.: Inputs of glacially derived dissolved and colloidal iron to the coastal ocean and implications for primary productivity, *Global Biogeochem. Cy.*, 22, GB3013, doi:10.1029/2007gb003106, 2008.
- Steigenberger, S., Statham, P. J., Völker, C., and Passow, U.: The role of polysaccharides and diatom exudates in the redox cycling of Fe and the photoproduction of hydrogen peroxide in coastal seawaters, *Biogeosciences*, 7, 109–119, 2010, <http://www.biogeosciences.net/7/109/2010/>.
- Stolpe, B., Hassellöv, M., Andersson, K., and Turner, D. R.: High resolution ICPMS as an on-line detector for flow field-flow fractionation; multi-element determination of colloidal size distributions in a natural water sample, *Analytica Chimica Acta*, 535, 109–121, 2005.
- Stolpe, B. and Hassellöv, M.: Changes in size distribution of fresh water nanoscale colloidal matter and associated elements on mixing with seawater, *Geochim. Cosmochim. Acta*, 71, 3292–3301, 2007.
- Stolpe, B., and Hassellöv, M.: Colloidal biopolymers binding iron, copper, silver, lanthanum and lead in coastal seawater – significance for the seasonal and spatial variations in element size distributions, *Limnol. Oceanogr.*, 55(1), 187–202, 2010.
- Straub, S. M. and Schmincke, H. U.: Evaluating the tephra input into Pacific Ocean sediments: distribution in space and time, *Geologische Rundschau*, 87, 461–476, 1998.
- Strzepek, R. F. and Harrison, P. J.: Photosynthetic architecture differs in coastal and oceanic diatoms, *Nature*, 431, 689–692, 2004.
- Strzepek, R. F., Maldonado, M. T., Higgins, J. L., Hall, J., Safi, K., Wilhelm, S. W., and Boyd, P. W.: Spinning the “Ferrous Wheel”: The importance of the microbial community in an iron budget during the FeCycle experiment, *Global Biogeochem. Cy.*, 19, GB4S26, doi:10.1029/2005GB002490, 2005.
- Sunda, W. and Huntsman, S.: Effect of pH, light, and temperature on Fe-EDTA chelation and Fe hydrolysis in seawater, *Mar. Chem.*, 84, 35–47, 2003.
- Sunda, W. G. and Huntsman, S. A.: Effect of competitive interactions between manganese and copper on cellular manganese and growth in estuarine and oceanic species of the diatom *Thalassiosira*, *Limnol. Oceanogr.*, 28, 924–934, 1983.
- Sunda, W. G. and Huntsman, S. A.: Iron uptake and growth limitation in oceanic and coastal phytoplankton, *Mar. Chem.*, 50, 189–206, 1995.
- Sunda, W. G. and Huntsman, S. A.: Processes regulating cellular metal accumulation and physiological effects: Phytoplankton as a model system, *The Sciences of the Total Environment*, 219, 165–181, 1998.
- Sunda, W. G. and Huntsman, S. A.: Effect of Zn, Mn, and Fe on Cd accumulation in phytoplankton: Implications for oceanic Cd cycling, *Limnol. Oceanogr.*, 45, 1501–1516, 2000.
- Takeda, S.: Influence of iron availability on nutrient consumption ratio of diatoms in oceanic waters, *Nature*, 393, 774–777, 1998.
- Timmermans, K. R., van der Wagt, B., Veldhuis, M. J. W., Maatman, A., and de Baar, H. J. W.: Physiological responses of three species of marine pico-phytoplankton to ammonium, phosphate, iron and light limitation, *J. Sea Res.*, 53, 109–120, 2005.
- Toner, B. M., Fakra, S. C., Manganini, S. J., Santelli, C. M., Marcus, M. A., Moffett, J., Rouxel, O., German, C. R., and Edwards, K. J.: Preservation of iron(II) by carbon-rich matrices in a hydrothermal plume, *Nature Geosci.*, 2, 197–201, doi:10.1038/ngeo433, 2009.
- Tovar-Sanchez, A., Sanudo-Wilhelmy, S. A., Kustka, A. B., Agusti, S., Dachs, J., Hutchins, D. A., Capone, D. G., and Duarte, C. M.: Effects of dust deposition and river discharges on trace metal composition of *Trichodesmium* spp. in the tropical and subtropical North Atlantic Ocean, *Limnol. Oceanogr.*, 51, 1755–1761, 2006.
- Tovar-Sanchez, A., Duarte, C. M., Hernández-León, S., and Sañudo-Wilhelmy, S. A.: Krill as a central node for iron cycling in the Southern Ocean, *Geophys. Res. Lett.*, 34, L11601, doi:10.1029/2006GL029096, 2007.
- Trapp, J. M., Millero, F. J., and Prospero, J. M.: Trends in the solubility of iron in dust-dominated aerosols in the Equatorial Atlantic Trade Winds: The importance of iron speciation and sources, *Geochem. Geophys. Geosyst.*, 11(1), in press, 2010.
- Tsuda, A., Takeda, S., Saito, H., Nishioka, J., Nojiri, Y., Kudo, I., Kiyosawa, H., Shiimoto, A., Imai, K., Ono, T., Shimamoto, A., Tsumune, D., Yoshimura, T., Aono, T., Hinuma, A., Kinugasa, M., Suzuki, K., Sohrin, Y., Noiri, Y., Tani, H., Deguchi, Y., Tsurushima, N., Ogawa, H., Fukami, K., Kuma, K., and Saino, T.: A mesoscale iron enrichment in the western Subarctic Pacific induces a large centric diatom bloom, *Science*, 300, 958–961, 2003.
- Turner, D. R. and Hunter, K. A.: *The Biogeochemistry of Iron in Seawater*, IUPAC Series on Analytical and Physical Chemistry of Environmental Systems, John Wiley and Sons Ltd., West Sussex, England, 396 pp., 2001.
- Twining, B. S., Baines, S. B., Fisher, N. S., and Landry, M. R.: Cellular iron contents of plankton during the Southern Ocean Iron Experiment (SOFEX), *Deep-Sea Res.*, 51, 1827–1850, doi:10.1016/j.dsr.2004.08.007, 2004.
- Ussher, S. J., Worsfold, P. J., Achterberg, E. P., Laes, A., Blain, S., Laan, P., and de Baar, H. J. W.: Distribution and redox speciation

- of dissolved iron on the European continental margin, *Limnol. Oceanogr.*, 52, 2530–2539, 2007.
- Vasconcelos, M., Leal, M. F. C., and van den Berg, C. M. G.: Influence of the nature of the exudates released by different marine algae on the growth, trace metal uptake, and exudation of *Emiliana huxleyi* in natural seawater, *Mar. Chem.*, 77, 187–210, 2002.
- Venter, J. C., Remington, K., Heidelberg, J. F., Halpern, A. L., Rusch, D., Eisen, J. A., Wu, D. Y., Paulsen, I., Nelson, K. E., Nelson, W., Fouts, D. E., Levy, S., Knap, A. H., Lomas, M. W., Nealson, K., White, O., Peterson, J., Hoffman, J., Parsons, R., Baden-Tillson, H., Pfannkoch, C., Rogers, Y. H., and Smith, H. O.: Environmental genome shotgun sequencing of the Sargasso Sea, *Science*, 304, 66–74, doi:10.1126/science.1093857, 2004.
- Verdugo, P., Alldredge, A. L., Azam, F., Kirchman, D. L., Passow, U., and Santschi, P. H.: The oceanic gel phase: a bridge in the DOM-POM continuum, *Mar. Chem.*, 92, 67–85, 2004.
- Vigenault, B. and Campbell, P. G. C.: Uptake of cadmium by freshwater green algae: Effects of pH and aquatic humic substances, *J. Phycol.*, 41, 55–61, 2005.
- Vong, L., Laës, A., and Blain, S.: Determination of iron-porphyrin-like complexes at nanomolar levels in seawater, *Analytica Chimica Acta*, 588, 237–244, doi:10.1016/j.aca.2007.1002.1007 2007.
- Vraspir, J. and Butler, A.: Chemistry of marine ligands and siderophores, *Ann. Rev. Mar. Sci.*, 1, 43–63, 2009.
- Wagener, T., Guieu, C., Losno, R., Bonnet, S., and Mahowald, N.: Revisiting atmospheric dust export to the Southern Hemisphere ocean: Biogeochemical implications, *Global Biogeochem. Cy.*, 22, GB2006, doi:10.1029/2007gb002984, 2008.
- Wang, W. X. and Dei, R. C. H.: Bioavailability of iron complexed with organic colloids to the cyanobacteria *Synechococcus* and *Trichodesmium*, *Aquatic Microbial Ecology*, 33, 247–259, 2003.
- Watson, A. J.: Iron limitation in the oceans, in: *The biogeochemistry of iron in seawater*, edited by: Turner, D. R., and Hunter, K. A., John Wiley & Sons Ltd., Chichester, 9–33, 2001.
- Weber, L., Völker, C., Oschlies, A., and Burchard, H.: Iron profiles and speciation of the upper water column at the Bermuda Atlantic Time-series Study site: a model based sensitivity study, *Biogeosciences*, 4, 689–706, 2007, <http://www.biogeosciences.net/4/689/2007/>.
- Wells, M. L.: A neglected dimension, *Nature*, 391, 530–531, 1998.
- Wells, M. L., Trick, C. G., Cochlan, W. P., Hughes, M. P., and Trainer, V. L.: Domoic acid: The synergy of iron, copper, and the toxicity of diatoms, *Limnol. Oceanogr.*, 50, 1908–1917, 2005.
- Willey, J. D., Kieber, R. J., Seaton, P. J., and Miller, C.: Rainwater as a source of Fe(II)-stabilizing ligands to seawater, *Limnol. Oceanogr.*, 53, 1678–1684, 2008.
- Wolff, E. W., Fischer, H., Fundel, F., Ruth, U., Twarloh, B., Littot, G. C., Mulvaney, R., Rothlisberger, R., de Angelis, M., Boutron, C. F., Hansson, M., Jonsell, U., Hutterli, M. A., Lambert, F., Kaufmann, P., Stauffer, B., Stocker, T. F., Steffensen, J. P., Bigler, M., Siggaard-Andersen, M. L., Udisti, R., Becagli, S., Castellano, E., Severi, M., Wagenbach, D., Barbante, C., Gabrielli, P., and Gaspari, V.: Southern Ocean sea-ice extent, productivity and iron flux over the past eight glacial cycles, *Nature*, 440, 491–496, doi:10.1038/nature04614, 2006.
- Worms, I., Simon, D. F., Hassler, C. S., and Wilkinson, K. J.: Bioavailability of trace metals to aquatic microorganisms: importance of chemical, biological and physical processes on biouptake, *Biochimie*, 88, 1721–1731, doi:10.1016/j.biochi.2006.09.008, 2006.
- Wu, J. F., Boyle, E., Sunda, W., and Wen, L. S.: Soluble and colloidal iron in the oligotrophic North Atlantic and North Pacific, *Science*, 293, 847–849, 2001.
- Wu, J. F., Chung, S. W., Wen, L. S., Liu, K. K., Chen, Y. L. L., Chen, H. Y., and Karl, D. M.: Dissolved inorganic phosphorus, dissolved iron, and Trichodesmium in the oligotrophic South China Sea, *Global Biogeochem. Cy.*, 17, doi:10.1029/2002GB001924, 2003.
- Ye, Y., Völker, C., and Wolf-Gladrow, D. A.: A model of Fe speciation and biogeochemistry at the Tropical Eastern North Atlantic Time-Series Observatory site, *Biogeosciences*, 6, 2041–2061, 2009, <http://www.biogeosciences.net/6/2041/2009/>.
- Yoosuf, S., Sutton, G., Rusch, D. B., Halpern, A. L., Williamson, S. J., Remington, K., Eisen, J. A., Heidelberg, K. B., Manning, G., Li, W. Z., Jaroszewski, L., Cieplak, P., Miller, C. S., Li, H. Y., Mashiyama, S. T., Joachimiak, M. P., van Belle, C., Chandonia, J. M., Soergel, D. A., Zhai, Y. F., Natarajan, K., Lee, S., Raphael, B. J., Bafna, V., Friedman, R., Brenner, S. E., Godzik, A., Eisenberg, D., Dixon, J. E., Taylor, S. S., Strausberg, R. L., Frazier, M., and Venter, J. C.: The Sorcerer II Global Ocean Sampling expedition: Expanding the universe of protein families, *PLoS Biol.*, 5, 432–466, doi:10.1371/journal.pbio.0050016, 2007.
- Yoshida, T., Hayashi, K., and Ohmoto, H.: Dissolution of iron hydroxides by marine bacterial siderophore, *Chem. Geol.*, 184, 1–9, 2002.
- Zhang, W. and Wang, W.-X.: Colloidal organic carbon and trace metal (Cd, Fe, and Zn) releases by diatom exudation and copepod grazing, *Journal of Experimental Marine Biology and Ecology*, 307, 17–34, 2004.