Atmosphere-ocean changes in the Pacific Southern Ocean over the past 1 Million years and implications for global climate

Frank Lamy¹, Helge Arz², Oliver Esper¹, Rainer Gersonde¹, Ian Hall³, Naomi Harada⁴, Andrea Jaeschke¹, Jerome Kaiser², Rolf Kilian⁵, Gerhard Kuhn¹, Fabrice Lambert⁶, Carina Lange⁷, Lester Lembke-Jene¹, Alfredo Martinez-Garcia⁸, Ralf Tiedemann¹, Gisela Winckler⁹, Shuzhuang Wu¹

¹AWI, Bremerhaven, Germany, Frank.Lamy@awi.de;
²Leibniz Institute for Baltic Sea Research, Rostock-Warnemünde, Germany;
³School of Earth and Ocean Sciences, Cardiff University, Cardiff, CF10 3AT, UK;
⁴Japan Agency for Marine-Earth Science and Technology, Yokosuka 237-0061, Japan;
⁵Universität Trier, Trier, Germany;
⁶Department of Physical Geography, Pontifical Catholic University of Chile, Chile;
⁷COPAS/IDEAL, University of Concepción, Concepción, Chile;
⁸Max-Planck-Institut für Chemie, Mainz, Germany;
⁹Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA.

<u>Abstract</u>

Atmosphere-ocean interactions play an important role for understanding processes and feedbacks in the Southern Ocean (SO) and are relevant for changes in Antarctic ice-sheets and atmospheric CO₂ concentrations. The most important atmospheric forcing at high and mid-latitudes of the Southern Hemisphere is the westerly wind belt (SWW), which strongly affects the strength and extension of the Antarctic Circumpolar Current (ACC), upwelling of deep-water masses, and controls the back-flow of intermediate waters to the tropics. In order to address orbital and millennial-scale changes of the SWW and the ACC, we present sediment proxy records from the Pacific SO including the Chilean Margin and the Drake Passage.

The Drake Passage (DP) represents the most important oceanic gateway along the ACC. Based on grain-size and geochemical properties of sediment records from the southernmost continental margin of South America, we reconstruct changes in DP throughflow dynamics over the past 65,000 years. In combination with published sediment records from the Scotia Sea and preliminary sediment records from the central Drake Passage (Polarstern cruise PS97, 2016), we argue for a considerable total reduction of DP transport and reveal an up to ~40% decrease in flow speed along the northernmost ACC pathway entering the DP during glacial times. Superimposed on this long-term decrease are high-amplitude millennial-scale variations, which parallel Southern Ocean and Antarctic temperature patterns. The glacial intervals of strong weakening of the ACC entering the DP imply a reduced Pacific-Atlantic exchange via the DP ("cold-water route").

The reduced Drake Passage glacial throughflow was accompanied by a pronounced northward extension of the Antarctic cold-water sphere in the Southeast Pacific sector and stronger export of northern ACC water into the South Pacific gyre. These oceanographic changes are consistent with reduced SWW within the modern maximum wind strength zone over the subantarctic ACC and reduced wind forcing due to extended sea-ice further south. Despite this reduction in winds in the core of the westerlies, we observe 3-fold higher dust deposition during glacial periods in

Past Antarctic Ice Sheet Dynamics (PAIS) Conference

September 10-15th 2017, Trieste - Italy

the Pacific Southern Ocean (SO). This observation may be explained by a combination of factors including more expanded arid dust source areas in Australia and a northward extent or enhancement of the SWW over Southeast Australia during glacials that would plausibly increase the dust uptake and export into the Pacific SO. Such scenario would imply stronger SWW at the present northernmost margin of the wind belt coeval with weaker core westerlies in the south and reduced ACC strength, including Drake Passage throughflow during glacials.

We conclude that changes in DP throughflow play a critical role for the global meridional overturning circulation and interbasin exchange in the Southern Ocean, most likely regulated by variations in the westerly wind field and changes in Antarctic sea-ice extent.

Keywords: Pelagic Southern Ocean, Antarctic Circumpolar Current, Southern Westerlies, Teleconnections.