

1 **Global phylogeography of *Oithona similis* s.l. (Crustacea, Copepoda, Oithonidae) - a**
2 **cosmopolitan plankton species or a complex of cryptic lineages?**

3

4 Astrid Cornils*, Britta Wend-Heckmann, Christoph Held

5

6 Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Am Handelshafen
7 12, 27570 Bremerhaven, Germany

8

9 *Corresponding author: Astrid Cornils

10 E-mail: astrid.cornils@awi.de

11 Phone: +4947148312077

12

13

14 **Abstract**

15 Traditionally, many small-sized copepod species are considered to be widespread, bipolar or
16 cosmopolitan. However, these large-scale distribution patterns need to be re-examined in
17 view of increasing evidence of cryptic and pseudo-cryptic speciation in pelagic copepods.

18 Here, we present a phylogeographic study of *Oithona similis* s.l. populations from the Arctic
19 Ocean, the Southern Ocean and its northern boundaries, the North Atlantic and the
20 Mediterranean Sea. *O. similis* s.l. is considered as one of the most abundant species in
21 temperate to polar oceans and acts as an important link in the trophic network between the
22 microbial loop and higher trophic levels such as fish larvae. Two gene fragments were
23 analysed: the mitochondrial cytochrome oxidase c subunit I (COI), and the nuclear ribosomal
24 28S genetic marker. Seven distinct, geographically delimited, mitochondrial lineages could
25 be identified, with divergences among the lineages ranging from 8 to 24 %, thus representing
26 most likely cryptic or pseudocryptic species within *O. similis* s.l. Four lineages were identified
27 within or close to the borders of the Southern Ocean, one lineage in the Arctic Ocean and
28 two lineages in the temperate Northern hemisphere. Surprisingly the Arctic lineage was more
29 closely related to lineages from the Southern hemisphere than to the other lineages from the
30 Northern hemisphere, suggesting that geographic proximity is a rather poor predictor of how
31 closely related the clades are on a genetic level.

32

33 **Keywords:** *Oithona*, Copepoda, species delimitation, biogeography, evolutionary history

34 **Introduction**

35 The existence of widespread or cosmopolitan species is widely accepted in marine plankton
36 due to the lack of apparent geographic barriers and high dispersal potentials. Thus, it has
37 been assumed that allopatric speciation may be less important in pelagic species than
38 sympatric or parapatric speciation (Norris, 2000). During the past years molecular studies
39 have revealed that many supposedly widespread species are mosaics of several cryptic or
40 pseudocryptic species (Andrews et al., 2014; Burridge et al., 2015; Cornils and Held, 2014;
41 Darling et al., 2007; Halbert et al., 2013; Hunt et al., 2010; Miyamoto et al., 2012). The mode
42 of speciation in the pelagic is largely unknown but several features have been suggested as
43 possible barriers to gene flow in the oceans: landmasses or continents (Blanco-Bercial et al.,
44 2011a), oceanic gyres (Andrews et al., 2014), frontal systems (Stupnikova et al., 2013) or
45 temperature and salinity ranges (Chen and Hare, 2011). Also vertical partitioning of cryptic
46 species has been observed (Miyamoto et al., 2010; Weiner et al., 2012). The presence of
47 cryptic or pseudocryptic species within nominal pelagic species provide evidence that
48 traditional species concepts based on morphological identification may have greatly
49 underestimated species richness in the oceans as suggested previously (Norris, 2000).

50 The cyclopoid copepod *Oithona similis* s.l. Claus, 1866 is accepted as a cosmopolitan
51 species, presumably occurring in all oceanic zones (Razouls et al., 2016) and presumably
52 the most abundant copepod species worldwide, although its importance has long been
53 overlooked due to the use of too coarse mesh sizes for plankton hauls (Gallienne and
54 Robins, 2001; Turner, 2004). Previous investigations have stressed that *O. similis* s.l. is a
55 prominent link between the microbial loop and higher trophic levels (Atkinson, 1995;
56 Castellani et al., 2005; Zamora-Terol et al., 2013). Its preferred habitats are polar and
57 temperate epipelagic marine waters where it is often more abundant than calanoid copepods
58 (Atkinson, 1998; Hopcroft et al., 2010; Pinkerton et al., 2010; Sabatini and Kiørboe, 1994;
59 Zamora-Terol et al., 2014). Furthermore, it has also been found in subtropical and tropical
60 waters (Cepeda et al., 2012; Nishida, 1985), although some of the reported findings seem
61 questionable (Nishida, 1985). Morphological differences have not been detected between *O.*
62 *similis* s.l. populations across latitudes, except for morphometric differences in the Northern
63 hemisphere, where prosome length and cephalon shape differed significantly among
64 populations from North Atlantic and Arctic water masses (Shuvalov, 1972). He suspected
65 that *O. similis* s.l. is “a polytypic species which comprises various subspecies or intraspecific
66 forms and groups with distinct ecological requirements regarding temperature throughout the
67 life cycle.” A first attempt to investigate the population structure of *O. similis* s.l., using the
68 nuclear molecular marker 28S, revealed low levels of intraspecific variety between North and
69 South Atlantic populations (Cepeda et al., 2012). However, the polar habitats of *O. similis* s.l.
70 were not included in that study.

71

72 In polar epipelagic zooplankton communities *O. similis* s.l. is the only dominant copepod
73 species with a presumed cosmopolitan distribution. All other dominant species are restricted
74 to polar and subpolar waters of the respective hemisphere. In the Arctic Ocean, predominant
75 epipelagic species are *Calanus glacialis* Jaschnov, 1955, *Calanus hyperboreus* Kröyer,
76 1838, *Pseudocalanus* spp. or *Triconia borealis* Sars, 1918 (Ashjian et al., 2003; Hunt et al.,
77 2014; Kosobokova and Hirche, 2000; Madsen et al., 2008), while in the Southern Ocean and
78 adjacent waters *Calanoides acutus* Giesbrecht, 1902, *Calanus propinquus* Brady, 1883,
79 *Calanus simillimus* Giesbrecht, 1892, *Rhincalanus gigas* Brady, 1883 and *Ctenocalanus citer*
80 Heron and Bowman, 1971 account for the majority of abundance and biomass (Atkinson,
81 1998; Hunt and Hosie, 2006a; 2006b; Park and Ferrari, 2009).

82 The present study aims at a global assessment of evolutionary lineages within *Oithona*
83 *similis* s.l. using sequences of two gene fragments: the mitochondrial cytochrome oxidase *c*
84 subunit I (COI), and the nuclear large subunit ribosomal gene (28S). We analysed specimens
85 from both the Southern Ocean and the Arctic Ocean, from different localities in the Atlantic
86 Ocean, and from the Bay of Villefranche in the northwestern Mediterranean Sea near the
87 type locality of *O. similis* s.l. (Claus 1866). The data set was complemented with previously
88 published sequences from the Atlantic and Pacific Ocean. Using several independent
89 species delimitation methods we estimated the existence and number of cryptic or
90 pseudocryptic species within *O. similis* s.l. Additionally, specimens from co-occurring *Oithona*
91 species in the polar systems (*Oithona frigida* Giesbrecht, 1902, *Oithona atlantica* Farran,
92 1908) and the temperate to subtropical regions (*Oithona nana* Giesbrecht, 1893, *Oithona*
93 *plumifera* Baird, 1843, *Oithona davisae* Ferrari and Orsi, 1984) were obtained to examine the
94 interspecific variability in the COI and 28S gene.

95

96 **Material and Methods**

97 **Sampling, and preparation for molecular analysis**

98 For the present study *Oithona* specimens were collected on various research cruises from
99 2006 to 2013 (Fig. 1, Appendix 1). In the polar regions stratified mesozooplankton samples
100 were collected with a multinet (HydroBios MultiNet Type Midi, 0.25 m² aperture), equipped
101 with five nets of 55 to 200 µm mesh-size. During the Polarstern cruises ANT XXIV/2, ARK
102 XXIII/ and ARK XXV/1, specimens were sorted alive under a stereomicroscope immediately
103 after the haul and were transferred to pure ethanol (undenatured 96% EtOH) for genetic
104 analysis. The remaining sample was preserved in 4% formaldehyde seawater solution. All
105 other Antarctic and Arctic specimens used in this study were sorted from mesozooplankton
106 samples preserved in pure ethanol. The ethanol in all samples was changed after 24 hours
107 to ensure optimal preservation for genetic analyses. For comparison, specimens from other

108 regions were included in the study, e.g. specimens from Villefranche sur Mer (Ligurian Sea)
109 close to the type locality of *O. similis* (Claus, 1866), and from Helgoland Roads (North Sea),
110 the type locality for *Oithona helgolandica* (Claus, 1863). The latter has often been classified
111 as synonymous with *O. similis* (for review see Razouls et al. 2016), but the original
112 description of a male specimen of *O. helgolandica* (Claus 1863) probably refers to *Oithona*
113 *nana* (Sars, 1918). In this study, we focused on female *O. similis* s.l. in order to eliminate the
114 influence of sexual dimorphisms of body length and morphological characters.

115

116 **Morphological characterization**

117 Prior to DNA extraction the specimens were placed in distilled water for up to five minutes at
118 room temperature and were identified morphologically under a stereomicroscope under low
119 light conditions. In addition, total length and Prosome:Urosome (P:U) ratios of 50 specimens,
120 representing each of the molecular defined groups, were measured (Appendix 1). To
121 evaluate whether cryptic or pseudocryptic species are found within *O. similis* s.l., three to ten
122 female specimens were set aside for further morphological analysis. These females were
123 dissected and the structure and ornamentation of the swimming legs was studied.

124

125 **DNA extraction, amplification and sequencing**

126 DNA was extracted from single individuals following the protocol for DNA purification from
127 tissues of the QIAamp DNA Mini Kit (Qiagen) with the following modifications: (1) The
128 individuals were placed in a tube containing ATL buffer and glass beads (diameter: 425 –
129 600 μm ; Sigma G8772), vortexed at maximum speed for five minutes and centrifuged for 10
130 min at 13,200 rpm (revolutions per minute), before the proteinase K was added. (2) Then the
131 tubes were incubated in a thermo-shaker at 56°C and 550 rpm overnight. (3) 1 μl carrier
132 RNA was applied with the AL buffer. (4) The DNA was eluted in 50 μl AE buffer. DNA
133 samples were stored at -20°C until further analysis.

134 The mitochondrial protein coding gene COI was amplified using the primer sets LCO1490
135 and HCO2198 (Folmer et al., 1994). Due to difficulties with the amplification in some
136 specimens a versatile primer set based on a highly conservative mitochondrial DNA region
137 was used (L1384-COI and H2612-COI; (Machida et al., 2004). Polymerase Chain Reaction
138 (PCR) amplifications were performed in 25 μl reaction volumes. For COI the reaction volume
139 included 5 μl of 5x KAPA2G Buffer B (KAPABiosystems), 5 μl of 5x Enhancer 1
140 (KAPABiosystems), 0.125 μl of 100 μM of each primer, 0.5 μl of 10 mM KAPA dNTP Mix,
141 0.15 μl of 5 U/ μl KAPA2G Robust DNA Polymerase (KAPABiosystems) and 3 μl of DNA
142 template solution. The PCR for COI consisted of 35 cycles of denaturation at 95°C for 20 s,
143 annealing at 49°C for 20 s and extension at 72°C for 20 s. An 800 bp fragment of the large
144 subunit (28S) ribosomal DNA was amplified using the primer set 28SF1 and 28SR1 (Ortman,

145 2008). PCR amplifications were carried out according to the protocols of (Cepeda et al.,
146 2012).
147 PCR products were run on a 2% agarose/TBE gel and afterwards stained with ethidium
148 bromide for band characterization. For COI positive results were either purified with the
149 QIAquick PCR purification kit (Qiagen) or with ExoSap-IT (0.25 µl exo, 1 µl SAP).
150 Subsequently they were used for cycle sequencing with Big Dye Terminator Ver. 3.1 (Applied
151 Biosystems Inc., ABI) and the same primers as for the PCR amplifications. The sequences
152 were run on an ABI 3130XL DNA sequencer (Applied Biosystems). In case of 28S positive
153 PCR results were sent for purification and sequencing to Eurofins Genomics
154 (<http://www.eurofins.de>) for purification with the Agencourt AMPure XP systems (Beckman
155 Coulter) and sequencing on ABI 3730XL DNA sequencer (Applied Biosystems).

156

157 **Sequence editing and additional sequences**

158 In total, 223 COI and 70 28S sequences were obtained. In CodonCode Aligner Vers. 3.7.1.1
159 (CodonCode Corporation) both strands were assembled into consensus sequences, aligned
160 and manually checked for sequencing errors. Twenty-one COI sequences labelled *Oithona*
161 *similis* were added from GenBank and twelve additional 28S sequences under the name of
162 *Oithona helgolandica* (Table 1), resulting in 244 COI and 82 28S sequences for further
163 analyses. Additional COI GenBank sequences labelled "*Oithona similis*" (KC136272 –
164 KC136284) clustered in a preliminary neighbor-joining tree with sequences of *Oithona*
165 *brevicornis* (GenBank accession numbers AB604190, AB604189) instead of the *O. similis* s.l.
166 and were consequently removed from the dataset.

167 The aligned sequences were cut to a length of 579 bp (28S) and 438 bp (COI). All 28S
168 sequences had the same length, but 27 sequences of the 244 COI sequences were shorter
169 than 400bp. Additionally, the 28S sequences were aligned with MAFFT v7.157b using the L-
170 INS-I option (Kato and Standley, 2013). For the phylogenetic analysis and the haplotype
171 networks, the sequences were collapsed to haplotypes with jMOTU (Jones et al., 2011) with
172 zero base cut-off resulting in 41 (COI) and 15 (28S) unique haplotypes.

173 The amino acid sequence of the final COI alignment did not reveal any stop codons
174 indicating that pseudogenes probably were not present. Following the advice from (Calvignac
175 et al., 2011) the genetic diversity of each codon position was noted separately revealing that
176 the diversity was highest at the third codon position (1st codon position: 0.0528 ± 0.0109
177 standard error (S.E.), 2nd codon position: 0.0001 ± 0.0001 S.E. and 3rd codon position:
178 0.3434 ± 0.0183 S.E.). The new COI and 28S sequences were deposited in GenBank under
179 the accession numbers KU982646 - KU982960. Alignments, models and tree data for Figs. 2
180 - 4 are available at: <https://doi.org/10.1594/PANGAEA.862554>.

181

182

183 **Phylogenetic analysis**

184 To study the interspecific diversity and assess the monophyly of *O. similis* s.l. sequences of
185 other *Oithona* species were added (Table 1, 2; Fig. 2, 3). For the interspecific diversity
186 uncorrected *p*-distances were calculated with MEGA 6.06 (Tamura et al., 2013). For the
187 phylogenetic analysis sequences were analysed both concatenated and separated by gene
188 fragment. The best-fitting partitioning scheme and nucleotide substitution model were
189 determined with PartitionFinder v1.1.0 (Lanfear et al., 2012). For COI each codon position
190 was analysed separately. The selection criterion for the nucleotide substitution model was
191 the corrected Akaike information criterion AICc for small sample sizes. Only the models
192 available in the programs RaxML v 8.0 (Stamatakis, 2014), MrBayes 3.2 (Ronquist et al.,
193 2012) and BEAST 1.8.1 (Drummond et al., 2012) were allowed, resulting in the same
194 partitioning scheme: 28S, COI 1st codon position, COI 2nd codon position, COI 3rd codon
195 position. The best-fitting nucleotide substitution models for RAXML were GTR+I+G for 28S,
196 COI 1st and 2nd codon position, and GTR+G for COI 3rd codon position. For MrBayes the
197 following substitution models were chosen: 28S: GTR+I+G, COI 1st codon position:
198 GTR+I+G, COI 2nd codon position: GTR+I, and COI 3rd codon position: HKY+G. The
199 resulting partitioning schemes and substitution models were then implemented in RAXML
200 and MrBayes to carry out Maximum Likelihood (ML) and Bayesian Inference (BI) analyses,
201 respectively. RAXML was run under the option GTRGAMMAI and a complete random starting
202 tree of 10,000 bootstrap replicates. Best-known likelihood tree search was performed under
203 GTRGAMMAI and a completely random starting tree. The final tree topology was evaluated
204 under GTRGAMMAI to yield stable likelihood values. In MrBayes MCMC analysis was
205 conducted on two independent runs of 10,000,000 generations, each with four chains (1
206 cold, 3 hot), with a sample frequency of 1000 generations. To check for stationary distribution
207 the average standard deviation of the split frequencies were examined for convergence, i.e.
208 the two independent runs converge when the standard deviation of the split frequencies is
209 approaching zero (< 0.01, Ronquist et al., 2011), and the first 25 % of the sampled trees
210 were discarded as “burn-in”. Only Bayesian posterior probability (BPP) values > 0.90 were
211 displayed to show only strongly supported clades. As outgroup sequences of *Mesocyclops*
212 *leuckartii* Claus, 1857 from the most closely related family with published sequences for both
213 genes were chosen (GenBank Accession numbers: KF153692, KF357729).

214

215 **Species delimitation**

216 To test the hypothesis that *O. similis* s.l. is a complex of reproductively isolated species,
217 three independent species delimitation methods were applied to the sequences of both
218 genes: the General Mixed Yule Coalescent Model (GMYC; Pons et al., 2006), Automated

219 Barcoding Gap Discovery (ABGD; Puillandre et al., 2012) and Rosenberg's probability of
220 reciprocal monophyly (Rosenberg, 2007). GMYC is a maximum-likelihood approach, which
221 measures the transition from intra- to inter-species branching patterns to identify species
222 (Pons et al., 2006). The analysis was conducted with a Bayesian implementation of the
223 GMYC model (bGMYC) (Reid and Carstens, 2012) in the statistical computing language R
224 (R Development Core Team, 2015). This model differs from the original GMYC model as it
225 allows uncertainties in phylogenies and implements flexible prior distributions. Prior to the
226 analysis, ultrametric consensus trees were generated with BEAST 1.8.1. The program was
227 run using the nucleotide substitution model TrN+I+G and the trees were analysed without
228 partitioning the data as suggested by PartitionFinder and a strict molecular clock with a fixed
229 mean substitution rate (1.0). All other tree priors were left at their default values. Two
230 independent analyses of 10,000,000 generations were run in BEAST and sampled every
231 1,000 generations. The first 1,000 generations were discarded as burn-in. Tracer 1.6 was
232 used to check for a sufficient effective sampling size (ESS) and good mixing. The two runs
233 were then combined with Logcombiner from the BEAST package with a lower sampling
234 frequency to 98 trees for the multi-threshold bGMYC analysis (Monaghan et al., 2009). The
235 analysis was run with multiple thresholds under default settings. bGMYC returns a graph with
236 an ultrametric tree and a heatmap-like figure showing the posterior probabilities of
237 conspecificity (Reid and Carstens, 2012).

238 ABGD sorts the sequences into groups based on pairwise distances (Puillandre et al., 2012).
239 It is an iterative process and detects significant differences between intra- and interspecific
240 variations (i.e. barcoding gap). The COI alignment was uploaded to the website
241 <http://wwabi.snv.jussieu.fr/public/abgd/abgdweb.html> and run with the default settings (P
242 (prior limit to intraspecific diversity) ranged between 0.001 and 0.1; X (gap widths) = 1 using
243 the available models JC86 (Jukes-Cantor) and K80 (Kimura).

244 Rosenberg's probability of reciprocal monophyly was used to determine if the data set is
245 suitable to detect robust monophyletic lineages (Rosenberg, 2007). This method is
246 implemented as "rosenberg" in the R package SPIDER (Brown et al., 2012) and marks
247 dichotomous nodes where the probability of monophyly is significant.

248 In general, DNA barcoding is used to identify species within a standard gene region, in this
249 case COI (Hebert et al., 2003). New sequences are compared to a database of known
250 sequences. To test if the lineages identified with GMYC and ABGD can be correctly identified
251 via DNA barcoding the functions "nearNeighbour", "bestCloseMatch" and "threshID" of the R
252 package SPIDER were used. For this, the optimal threshold for species identification was
253 determined with the function "localMinima" from the R package SPIDER, and all sequences
254 were labeled according to the affiliation in the results of GMYC and ABGD. The function
255 "nearNeighbour" identifies the closest specimen and returns the score "true" if the species

256 label is the same, or “false” if the labels differ. The “bestCloseMatch” (best close match
257 analysis; (Meier et al., 2006) searches for the closest specimens and determines if it is within
258 a given threshold. If it is, then the outcome is scored as “correct”, if it is outside the threshold
259 the outcome is “no ID” (this is also true for singletons). If the closest match includes more
260 than one species the identification is “ambiguous”. When all the matches within the threshold
261 have different species labels the identification is “incorrect”. The third analysis (“threshID”) is
262 threshold-based with a given threshold genetic distance of 1%. This method resembles the
263 BOLD species identification systems
264 (http://www.boldsystems.org/index.php/IDS_OpenIdEngine). The results of this analysis are
265 the same as in the best close match analysis.
266 Haplotype networks are a common method to visualize genealogical relationships among *O.*
267 *similis* s.l. TCS haplotype networks (Clement et al., 2000) were created for both genes using
268 POPART (Leigh and Bryant, 2015). For this the 27 sequences shorter than 400bp were
269 removed to gain a better resolution of the relationships between the lineages.

270 **Divergence time estimation**

271 The program DAMBE5 (Xia, 2013) was used to check whether a molecular clock could be
272 used to date some of the nodes in our COI sequences. Using the test of Xia et al. (2003) the
273 program can measure the degree of substitution saturation. High levels of saturation are not
274 suitable for molecular clock estimation (Xia and Lemey, 2009). The program BEAST (settings
275 as above) was used to estimate the time of the most recent common ancestors (tmrca).
276 Fossils of copepods are rare (Harvey and Pedder, 2013; Selden et al., 2010) and therefore to
277 our knowledge no copepod-specific estimates of mutation rates exist (Blanco-Bercial et al.,
278 2011a; Marrone et al., 2013; Thum and Harrison, 2009). Often, tmrcas in copepods were
279 estimated from known sequences divergence of the decapod *Alpheus* spp. with 1.4% per
280 million years (Knowlton and Weigt, 1998). Although this rate has been widely used to
281 estimate molecular evolution in marine invertebrates, there are newer estimates in marine
282 invertebrates showing that substitution rates could also be considerably higher (e.g. Calvo et
283 al., 2015). At present however, we have refrained from using external information to calibrate
284 the tree and only provide estimates of relative divergence times with the 95% highest
285 posterior intervals (HPD) and not absolute node ages.

286

287 **Results**

288 Both the COI and the 28S phylogenetic trees yielded the specimens morphologically
289 identified as *O. similis* s.l. as a clearly monophyletic group (Fig. 2, 3). While the genetic
290 variation in the nuclear 28S gene fragment was consistently lower within *O. similis* s.l. (0 –
291 0.010) compared to the overlap with the distances to other *Oithona* species (0.023 – 0.210).
292 In COI genetic distances between *O. similis* s.l. and other *Oithona* species (0.192 – 0.331)

293 overlapped with the genetic variation within *O. similis* s.l. (0 - 0.235), thus suggesting that *O.*
294 *similis* s.l. is more likely a species complex than a panmictic species.

295

296 **Species delimitation**

297 A total of 244 COI sequences worldwide of *O. similis* s.l. were used in the present study.
298 Seven cryptic mitochondrial lineages and two singletons were discovered with the species
299 delimitation methods ABDG and bGMYC (Appendix 4). They were well-supported (posterior
300 probability (PP) > 0.95. For all nodes Rosenberg's probability of reciprocal monophyly was
301 significant (Appendix 4). The three species identification methods (nearest neighbor, best
302 close match and threshold ID) assigned 242 of the 244 COI sequences correctly, using a
303 genetic distance threshold of 0.053. The two singleton sequences from the Chukchi Sea
304 (ARK2 and ARK3) were scored as "no ID" thus indicating that they could not be assigned to
305 any of the discovered species. The maximum genetic distance (uncorrected *p*-distance)
306 within each lineage was always lower than 1%. Genetic divergence among the seven
307 mitochondrial lineages within the *O. similis* species complex ranged between 8.3 and 23.7%
308 (Table 3) and thus they were all more divergent than 8%, the proposed threshold in COI for
309 distinct species in copepods (Bucklin et al., 1999, Hill et al., 2001). Furthermore, the genetic
310 distances among lineages satisfied the 4x rule (Birky et al., 2005), as they were at least 4x
311 higher than genetic distances within lineages. Thus, most probably the seven independent
312 genetic lineages represent cryptic or pseudocryptic species within *O. similis* s.l.

313 The two species delimitation methods bGMYC and ABGD were also applied to the 82
314 sequences of the nuclear 28S gene fragment. For bGMYC no significant lineages were
315 found, while with ABGD three lineages were recovered. The lineage ARK1 is identical with
316 the mitochondrial lineage ARK1. Clade 2 includes all other 28S sequences, except for the
317 two sequences grouped in Clade 3. The latter belong to two specimens from the Polar Front.

318

319 **Phylogeography**

320 Although the sampling in this study may not be high enough to conclusively rule out the
321 presence of lineages or species in an area in which we did not sample it, the pattern
322 emerging from analysing our presence/pseudo-absence data shows seven mitochondrial
323 lineages with regional distributions rather than a single cosmopolitan species (Fig. 4,
324 Appendix 4). Three lineages and the two singletons are localized in the Northern
325 hemisphere. The lineage ARK1 was only found in the Arctic Ocean and its adjacent seas
326 (Chukchi Sea, Beaufort Sea, Canadian Basin, Central Arctic Ocean, Greenland Sea,
327 Norwegian Sea). Two mitochondrial lineages were identified from the North Atlantic (NA) and
328 the North Pacific (NP) respectively, the latter is based only on published sequences from
329 GenBank. The two singletons ARK2 and ARK3 were collected from the Chukchi Sea in the

330 Pacific part of the Arctic. In the Southern hemisphere four mitochondrial lineages could be
331 recovered with the species delimitation methods. The lineage ANT was found in Antarctic
332 waters (Weddell Gyre, the Coastal Current, the Amundsen Sea and the Polar Front). PF was
333 identified only from samples at the Polar Front of Atlantic and Indian Antarctic sectors. AS
334 was only found in the Amundsen Sea, and BC occurred in the Beagle Channel in Tierra del
335 Fuego (Fig. 1, 4).

336 An overlap between the lineages PF and ANT near the Polar Front (Atlantic sector; Appendix
337 2) revealed evidence for vertical partitioning of the two lineages (Fig. 5). While PF was
338 mainly found in the upper 100 m, ANT occurred mainly deeper than 100 m. Further south, in
339 the Weddell Gyre and the Coastal Current, PF was absent and ANT was distributed from the
340 surface to 250 m depth.

341 The haplotype network also showed that only the cryptic mitochondrial lineage from the
342 Arctic Ocean (ARK1) is recognized in the nuclear gene fragment 28S. All Arctic 28S shared
343 the same allele. However, most 28S sequences from the Southern Ocean and its boundaries
344 form another 28S allele (ANT, PF, AS, BC, Fig. 4) and sequences from the Northern Atlantic
345 (NA) and from the Southwest Atlantic (published in GenBank) form a third group with several
346 haplotypes.

347

348 **Divergence Time estimation**

349 The test of Xia et al. (2003) was applied to the COI data set with the proportion of invariable
350 sites as suggested by DAMBE5 (COI: 0.55864). Only fully resolved sites were considered.

351 The index of substitution saturation (ISS) was significantly lower than the critical ISS and the
352 degree of substitution saturation can be considered to be low in the data set. Thus, the data
353 set is suitable for further molecular clock analysis. The effective sampling size (ESS) was
354 higher than 1000 for most parameters, indicating convergence and good mixing of the
355 MCMC chain. The tmrcas were recovered with the 95% HPD. The phylogenetic analysis
356 revealed a strong separation between clades of the Northern hemisphere (NA, NP, ARK2,
357 ARK3) from the polar (ANT, ARK1) and subpolar clades (BC, PF, AS; Fig. 6). Conservative
358 estimates based on mitochondrial substitution rates from other crustaceans (Knowlton and
359 Weigt, 1998) suggest that this separation may have occurred prior to the middle Miocene
360 climate transition, about 21.5 - 13.4 million years ago (mya). Separation of the subpolar
361 clades and the polar groups occurred later in time. The separation of the two polar clades
362 ARK1 and ANT occurred last. Divergence time estimate from other crustaceans (see above)
363 date this event between 5.9 – 11.2 mya. This estimate for the last common ancestor of the
364 two polar lineages is a conservative estimate, if higher substitution rates are assumed (e.g.
365 5.16%, Calvo et al., 2015) the timing of this split shifts to even more recent estimates (around

366 2 mya (95% HPD: 1.6 - 3.0 mya)). The lineages in the Northern hemisphere (NP, ARK2,
367 ARK3) and (NA) lineages separated prior to those of the southern hemisphere.

368

369 **Morphology**

370 During the detailed morphological analysis it could be excluded that the genetic groups found
371 within *O. similis* s.l. were misidentified representatives of the closely related species *Oithona*
372 *fallax* or *Oithona decipiens*. All analysed individuals only had one outer spine at the outer
373 margin of the third exopod segment (Exp3) of swimming leg two (P2) and no inner marginal
374 spine of the first exopod segment (Exp1) of all four swimming legs (P1 – P4). In *O. fallax* two
375 spines on P2 Exp3 and one spine on P1-P4 Exp1 would be expected. All analysed
376 individuals had one outer marginal spine on the second exopod segment (Exp2) of the first
377 swimming leg (P1), which is missing in *O. decipiens*. Between the genetic groups no
378 differences in swimming leg ornamentation of *O. similis* s.l. could be detected. However, total
379 length measurements prior to DNA extraction of 51 specimens revealed that individuals from
380 lineage ANT were considerably larger than all others groups while the individuals from the
381 northeastern Atlantic were smallest (Appendix 3).

382

383 **Discussion**

384 The present study provides molecular evidence that *O. similis* s.l. is not a single species with
385 a cosmopolitan distribution, but instead a complex of cryptic or pseudocryptic genetic
386 lineages with much smaller, regional distributions. These lineages are confined to different
387 climate zones (Antarctic, Arctic, Polar Front, temperate Atlantic or Pacific). Genetic
388 divergence has also been reported from other widespread planktonic organisms (Burrige et
389 al., 2015; Casteleyn et al., 2010; Cornils and Held, 2014; Darling et al., 2007; Halbert et al.,
390 2013; Hunt et al., 2010; Kulagin et al., 2013). A recent global plankton metabarcoding survey
391 in the photic zone of the oceans even revealed that only 0.35% of OTUs (operational
392 taxonomic units) are cosmopolitan (de Vargas et al., 2015). These recent findings oppose
393 the assumption that global panmixia is common in planktonic species (e.g. Palumbi, 1992,
394 Norris, 2000).

395

396 **Taxonomic and phylogenetic implications**

397 It is noteworthy that the all mitochondrial as well as nuclear lineages of *Oithona similis* s.l.
398 form a well-supported monophylum and do not mix with other congeners. This means that
399 according to our present knowledge the species outside *O. similis* s.l. group can be
400 considered as outgroup, but do not have to be included in a reconstruction of evolutionary
401 history inside the group. Our results confirm the close phylogenetic relationships between *O.*
402 *similis* s.l. and the *O. atlantica* group (Nishida, 1985). Due to the lack of samples the close

403 relationship between *O. similis* s.l. and *O. decipiens* or *O. fallax* could not be investigated.
404 However, their presence within the *O. similis* s.l. group could be excluded morphologically.
405 The species delimitation methods GMYC and ABGD detected seven cryptic mitochondrial
406 lineages of *O. similis* s.l., but only three clearly distinct nuclear lineages could be recognized
407 using the 28S ribosomal gene. The two genes agreed in the discovery of a cryptic Arctic
408 lineage (ARK1), specimens representing the other six cryptic *O. similis* lineages were poorly
409 resolved and shared seven 28S alleles, allowing no further differentiation. These results
410 indicate that the speciation process in *O. similis* s.l. is probably too recent for the mutation
411 rate within the nuclear 28S gene fragment to be detectable.

412 Alternatively, the different patterns of divergence in the mitochondrial and nuclear gene
413 fragments, for example in the Southern hemisphere (ANT, PF, AS, BC) might also be
414 explained by a temporary separation and diversification of *O. similis* populations followed by
415 a period of hybridization. While hybridization has led to the homogenization of ephemeral
416 genetic differences in the nuclear genome, the lack of recombination has led to a persistent
417 record of this period of separation in the mitochondrial genes. According to this interpretation,
418 the coexisting mitochondrial lineages, although not indicating the existence of a second
419 species, would nevertheless count as a record of previous genetic isolation (see Dietz et al.,
420 2015).

421 Although ribosomal nuclear genes are known to have a low resolution within copepod
422 species (Machida and Tsuda, 2010; Marrone et al., 2013) and are mainly applied to
423 investigate monophyly of higher ranks (e.g. Blanco-Bercial et al., 2011b; Cornils and Blanco-
424 Bercial, 2013), they may include highly variable segments that differentiate between species
425 (e.g. Held, 2000) and the 28S gene fragment has been successfully applied to distinguish
426 between morphotypes of *Oithona dissimilis* Lindberg, 1940 (Ueda et al., 2011) and to identify
427 *Oithona* species, including *O. similis* s.l., in the Atlantic Ocean (Cepeda et al., 2012). In our
428 case however, the genetic differences of 28S were lower than in *O. dissimilis* (*O. dissimilis*:
429 1.84 – 2.21%; *O. similis* s.l.: 0 – 1.0%). Thus, to investigate the possibility of hybridization
430 nuclear genes with faster mutation rates, such as ITS (internal transcribed spacers) or
431 microsatellites, are needed (e.g. Makino and Tanabe, 2009, Parent et al., 2012).

432

433 **Identifying taxonomically valid species in *Oithona similis* s.l.**

434 Matching clades in our molecular phylogeny with available taxonomic species is a task in
435 itself and logically separate from demonstrating that the clades are genetically isolated from
436 one another. In this paper, *O. similis* s.l. specimens were taken near the type locality off the
437 coast of Nice (Claus, 1866, Fig. 1), the type locality of *O. similis* s.str. There has however,
438 been a long debate whether an earlier description by the same author of *O. helgolandica*
439 from Helgoland in the North Sea (Claus, 1863) is synonymous with the description from the

440 Mediterranean (for review see Cepeda et al., 2016). However, both descriptions are very
441 incomplete and the description from the North Sea could also match with *O. nana* (Sars,
442 1918). Our data show that the specimens from the Mediterranean (Villefranche), Helgoland
443 and other regions in the NE and NW Atlantic form a distinct mitochondrial lineage and may
444 thus all belong to *O. similis* s.str. This separation is also visible in the haplotype network of
445 the ribosomal gene fragment 28S and includes also sequences from the SW Atlantic, taken
446 from Cepeda et al. (2012). The placement of the specimens from the SW Atlantic within *O.*
447 *similis* s.str. remains unclear until a comparison with COI sequences is possible. Supporting
448 evidence that the Mediterranean and the North Atlantic population may be the same species
449 also comes from ecological research (Castellani et al., 2016) as they respond similarly to
450 temperature. Although a full taxonomic revision of the *Oithona similis* species complex is
451 beyond the scope of this paper but in view of the mounting evidence that it comprises many
452 species we believe that the best way to handle the situation currently is to use *O. similis* s.l.
453 together with a reference to the geographic origin as an interim identification. In order to
454 reconcile these records with a future taxonomic revision of *O. similis* it is advisable to keep
455 voucher samples suitable for molecular analysis.

456

457 **Biogeography and evolutionary history**

458 Although it is unclear if the clades characterized by diverged COI but uniform 28S sequences
459 have to be interpreted as fully diverged species or not, the differentiation of the mitochondrial
460 haplotypes is nevertheless indicative of historical isolation (Dietz et al., 2015). The
461 distribution of seven mitochondrial lineages in *O. similis* s.l. allows inferences about the
462 processes that have shaped the evolutionary history of this planktonic group that have been
463 obscured previously by treating it as a single evolutionary unit.

464 The distribution area of each of the newly delineated species or clades in *O. similis* s.l. is
465 considerably smaller but surprisingly geographic proximity is a rather poor predictor of how
466 closely related the clades are on a genetic level. The deepest split in our samples includes a
467 well-supported group of lineages exclusively from the Northern hemisphere. Within it, the
468 Northern Atlantic (NA) lineage is sister to a group of *O. similis* s.l. from the North Pacific (NP)
469 and the Pacific Arctic (ARK2 and ARK3), indicating that the speciation is taking place on an
470 ocean basin scale. The node leading to ARK2 and NP is poorly supported and thus, it is
471 difficult to decide if (1) the Arctic was colonized from the Atlantic and subsequently the North
472 Pacific or (2) the Atlantic and Pacific lineage existed in parallel, in which case the NP lineage
473 would be expected to have given rise to the two Arctic *O. similis* s.l. lineages. There is,
474 however, evidence that planktonic organisms in the North Pacific enter the Pacific Arctic (e.g.
475 Nelson et al., 2009). The two Arctic lineages ARK2 and ARK3 are however, singletons, and it
476 is not known if these lineages do not appear in the subarctic North Pacific as well.

477 One of the best-supported results is that clearly the Arctic was colonized twice by two very
478 distantly related lineages of *Oithona*. In addition to the two Arctic lineages in the northern
479 hemisphere group (ARK2, ARK3), a third Arctic *Oithona* species candidate (ARK1) is nested
480 deeply within a well-supported group of southern hemisphere group of *O. similis* species
481 candidates. In the nuclear 28S the two singletons share the same allele as the specimens
482 from the Southern hemisphere and not with the other Arctic (ARK1) and the NA assemblage.
483 A close relationship between subpolar North Pacific populations with populations from the
484 Southern hemisphere has also been observed in foraminiferans (Darling et al., 2007).
485 The lack of lower latitude *O. similis* s.l. in this analysis precludes a definitive statement at this
486 time, but several lines of evidence suggest that the group ARK1 indeed evolved from a cold-
487 adapted ancestor close to the Antarctic and separated from it once it managed the transit to
488 the Arctic Ocean. The relative divergence estimates (Fig. 6) also suggest that the Antarctic
489 and Arctic lineages are the youngest in the tree. This has also been observed in other polar
490 organisms (Held, 2000). If the sequence divergence estimate of 1.4% per mya (*Alpheus* sp.,
491 Knowlton and Weigt, 1988) is roughly correct, the split between ARK1 and ANT occurred
492 after the worldwide latitudinal climate gradient had steepened in the upper Miocene with the
493 Arctic and the Southern Ocean being already polar (Zachos et al., 2001). Although this steep
494 climate gradient must have made the dispersal to the other polar ocean difficult, it likely
495 made the chances of survival and successful colonization higher once it arrived there
496 because ARK1 likely inherited adaptations to polar conditions from the ancestral species it
497 shares with ANT.

498 This long-distance dispersal may be less unlikely than it may appear at first sight because at
499 about the same time two more independent splits occurred between a species pair that today
500 comprises a polar specialist and their closest relative that is geographically and climatically
501 distant, namely (AS+(PF+BC)) and (ARK2 and NP). It is reasonable to assume that the *O.*
502 *similis* s.l. species in the upper Miocene were more widely distributed than today making long
503 distance dispersal between climatic zones as well as across hemispheres rare but not
504 impossible before the climatic gradient steepened to the extent seen today.

505 Our data clearly show that the Antarctic (ANT) and the Arctic (ARK1) lineages are highly
506 divergent, which has also been shown for other common polar epipelagic species, e.g. the
507 pteropod *Limacina helicina* (Hunt et al., 2010) and foraminifera (Darling et al., 2007), and can
508 also be seen in the morphologically similar polar species of *Calanus* spp. (Kozol et al., 2012).
509 These results emphasize that cosmopolitan or bipolar distribution in epipelagic planktonic
510 species may be much less common than previously thought, possibly due to dietary or
511 reproductive requirements. Within the epi- to mesopelagic genus *Microcalanus* bipolar and
512 even cosmopolitan cryptic mitochondrial lineages have been found (Cornils, unpublished
513 data). In tropical regions these species show a meso- to bathypelagic distribution. Thus,

514 gene flow between the polar ecosystems may only be feasible if species are able to
515 submerge to deeper (colder) water layers in the tropics (Allcock, 2014).
516 Geographic overlaps of the lineages occurred only at the Antarctic Polar Front (ANT – PF,
517 ANT - AS) and in the Chukchi Sea (ARK1 – ARK2, ARK1 – ARK3). Our data indicate that the
518 apparent co-occurrence of the lineages ANT and PF is possibly a product of the encounter of
519 two different water masses, the colder waters of the East-Wind-Drift (Antarctic Intermediate
520 Water) and the warmer surface waters of the Antarctic Circumpolar Current (ACC). The
521 lineage PF may be associated with the warmer water masses as it was mainly found in the
522 surface waters, while ANT occurred below 100 m depth and thus, could be associated with
523 the deeper Antarctic Intermediate Water (Fig. 5). The latter is also known from the other
524 dominant epipelagic Antarctic copepod species: *Calanoides acutus*, *Rhincalanus gigas* or
525 *Calanus propinquus* can also be found as far north as the Polar Front, but they appear to be
526 associated with the deeper Antarctic Intermediate Water (Park and Ferrari, 2009). These
527 results reinforce the notion that the Antarctic Polar Front is an important ecological barrier for
528 zooplankton species (Marin, 1987). The same mechanism of different vertical distributions
529 may also account for the seemingly co-occurring mitochondrial lineages ANT and AS in the
530 Amundsen Sea.

531 In the Northern Atlantic our mitochondrial sequences did not reveal any geographic overlap
532 between the lineages ARK1 and NA. However, early morphological observations on *O.*
533 *similis* s.l. in the Northern hemisphere revealed considerable differences in size and in the
534 curvature of the cephalosome between Arctic specimens and boreal forms from the North
535 Atlantic and North Pacific (Shuvalov, 1972). Shuvalov observed that the Atlantic *O. similis* s.l.
536 form could be found as far north as Svalbard or the Barents Sea. These findings indicate that
537 the transport of the lineage NA via the North Atlantic Current to the Arctic Ocean would be
538 possible. This phenomenon is also known for other boreal species such as *Calanus*
539 *finmarchicus* (Hirche and Kosobokova, 2007). 28S sequences of *O. similis* s.l. from the
540 Irminger Current (Cepeda et al., 2012) and from the Norwegian Sea (present study) were
541 placed in the variable Atlantic group of the haplotype network indicating that the NA lineage
542 may also occur at higher latitudes.

543

544 **Ecological implications and response to environmental change**

545 In polar and temperate habitats *O. similis* s.l. is a prominent link between the microbial loop
546 and higher trophic levels such as fish larvae (Checkley, 1982; Paradis et al., 2012; Turner,
547 2004). Although we assume that the ecological role in the different pelagic communities
548 remains similar among the cryptic lineages there may be behavioral, ecological or
549 physiological differences between the lineages as it has been observed in marine protists
550 (Darling et al., 2007). Shuvalov (1972) found many dead *O. similis* s.l. specimens in abruptly

551 changing water masses at the interface of the Arctic and the North Atlantic Ocean and
552 assumed that temperature tolerance differs between the polar and the subpolar/temperate
553 specimens, since the seasonal cycle of water temperature varies greatly between high and
554 lower latitudes. Generally it has been observed that temperature and food availability
555 influence the fecundity of *O. similis* s.l. (Castellani et al., 2005; Ward and Hirst, 2007).
556 Unfortunately, large-scale studies comparing temperature tolerance of *O. similis* s.l. from
557 different latitudes are lacking so far. Thus, we will use the intensely studied globally
558 distributed *Calanus* species complex as a model for the *O. similis* s.l. complex. The *Calanus*
559 species are morphological very similar, but they exhibit differences e.g. in their reproductive
560 cycles (Niehoff et al., 2002). In the northern Atlantic latitudinal shifts within the *Calanus* spp.
561 have been observed during the past 40 years, probably associated with ocean warming
562 (Beaugrand et al., 2003, Chust et al., 2014). Temperature has been identified as the critical
563 factor that makes marine systems vulnerable to global warming and the zooplankton
564 composition, abundance and trophic efficiency is highly linked to it (Richardson, 2008).
565 We assume that the lineages associated with temperate water masses have a higher
566 tolerance to changing temperatures due to the pronounced seasonality compared to the
567 stenotherm polar lineages ARK1 and ANT. Future warming of the polar oceans could thus
568 initiate a poleward shift of the subpolar lineages of *O. similis* s.l. to the high Antarctic or an
569 invasion of the temperate Atlantic or Pacific lineages to the Arctic. These invasions could
570 cause severe changes in the seasonal availability of *Oithona* in the polar oceans and thus,
571 alter the food availability for higher trophic levels such as fish larvae. In the North Sea the
572 boreal *Calanus finmarchicus* is gradually replaced by the warm-temperate *Calanus*
573 *helgolandicus* which has lead to severe consequences for the recruitment of Atlantic cod
574 (Beaugrand et al., 2003). Therefore, understanding of the present environmental factors and
575 historical events that shape the biogeography of widespread zooplankton organisms is
576 critical to analyse the impact of climate change on higher tropic levels.
577 In conclusion, *O. similis* s.l. is not a single species with a circumglobal distribution. Instead, it
578 consists of at least seven distinct mitochondrial lineages that most likely represent cryptic or
579 pseudocryptic species. This study provides further evidence that widespread epipelagic
580 copepod species are often a complex of cryptic species with deep divergences in
581 mitochondrial genes.

582

583 **Acknowledgements**

584 We are grateful for the support of Andrea Eschbach in the molecular laboratory and to the
585 colleagues who provided samples for the present study: S. Gasparini provided samples from
586 Villefranche sur Mer. U. Tilmann took samples from the Northeastern Atlantic, the crew of the
587 vessel "Aade" took samples from Helgoland Roads. Other zooplankton samples were

588 provided by Holger Auel, Ann Bucklin, Barbara Niehoff, Sigrid Schnack-Schiel and Steve
589 Wickham. This work was supported by the Deutsche Forschungsgemeinschaft (DFG) in the
590 framework of the priority programme "Antarctic Research with comparative investigations in
591 Arctic ice areas" by the grants CO 706/2-1 (A. Cornils) and SCHM 352/43-1 (B. Wend-
592 Heckmann, C. Held).

593

594 **References**

- 595 Allcock, A.L., 2014. Chapter 10.4. Bipolarity, in: de Broyer, C., Koubbi, P. (Eds.),
596 Biogeographic Atlas of the Southern Ocean. The Scientific Committee on Antarctic
597 Research, Scott Polar Research Institute, Lensfield Road, Cambridge, CB2 1ER, United
598 Kingdom (www.scar.org)., pp. 1–9.
- 600 Andrews, K.R., Norton, E.L., Fernandez-Silva, I., Portner, E., Goetze, E., 2014. Multilocus
601 evidence for globally distributed cryptic species and distinct populations across ocean gyres
602 in a mesopelagic copepod. *Mol. Ecol.* 23, 5462–5479. doi:10.1111/mec.12950
603
- 604 Ashjian, C.J., Campbell, R.G., Welch, H.E., Butler, M., Van Keuren, D., 2003. Annual cycle in
605 abundance, distribution, and size in relation to hydrography of important copepod species in
606 the western Arctic Ocean. *Deep-Sea Res. I* 50, 1235–1261. doi:10.1016/S0967-
607 0637(03)00129-8
608
- 609 Atkinson, A., 1995. Omnivory and feeding selectivity in five copepod species during spring in
610 the Bellingshausen Sea, Antarctica. *ICES J. Mar. Sci.* 52, 385–396.
611
- 612 Atkinson, A., 1998. Life cycle strategies of epipelagic copepods in the Southern Ocean. *J.*
613 *Mar. Syst.* 15, 289–311.
614
- 615 Beaugrand, G., Brander, K.M., Lindley, J.A., Souissi, S., Reid, P.C., 2003. Plankton effect on
616 cod recruitment in the North Sea. *Nature* 426, 661–664.
617
- 618 Birky, C., Wolf, C., Maughan, H., Herbertson, L., Henry E., 2005. Speciation and selection
619 without sex. *Hydrobiologia* 546, 29–45.
620
- 621 Blanco-Bercial, L., Álvarez-Marqués, F., Bucklin, A., 2011a. Comparative phylogeography
622 and connectivity of sibling species of the marine copepod *Clausocalanus* (Calanoida). *J. Exp.*
623 *Mar. Biol. Ecol.* 404, 108–115. doi:10.1016/j.jembe.2011.05.011
624
- 625 Blanco-Bercial, L., Bradford-Grieve, J.M., Bucklin, A., 2011b. Molecular phylogeny of the
626 Calanoida (Crustacea: Copepoda). *Mol. Phylogenet. Evol.* 59, 103–113.
627
- 628 Brown, S.D.J., Collins, R.A., Boyer, S., Lefort, M.-C., Malumbres-Olarte, J., Vink, C.J.,
629 Cruickshank, R.H., 2012. Spider: An R package for the analysis of species identity and
630 evolution, with particular reference to DNA barcoding. *Mol. Ecol. Resour.* 12, 562–565.
631 doi:10.1111/j.1755-0998.2011.03108.x
632
- 633 Bucklin, A., Guarneri, M., Hill, R.S., Bentley, A., Kaartvedt, S., 1999. Taxonomic and
634 systematic assessment of planktonic copepods using mitochondrial COI sequence variation
635 and competitive, species-specific PCR. *Hydrobiologia* 401, 239–254.
636
- 637 Burridge, A.K., Goetze, E., Raes, N., Huisman, J., Peijnenburg, K.T.C.A., 2015. Global
638 biogeography and evolution of *Cuvierina* pteropods. *BMC Evol. Biol.* 15, 39.
639 doi:10.1186/s12862-015-0310-8
640

641 Calvignac, S., Konecny, L., Malard, F., Douady, C.J., 2011. Preventing the pollution of
642 mitochondrial datasets with nuclear mitochondrial paralogs (numts). *Mitochondrion* 11, 246–
643 254. doi:10.1016/j.mito.2010.10.004
644
645 Calvo, M., Alda, F., Oliverio, M., Templado, J., Machordom, A., 2015. Surviving the
646 Messinian Salinity Crisis? Divergence patterns in the genus *Dendropoma* (Gastropoda:
647 Vermetidae) in the Mediterranean Sea. *Mol. Phylogenet. Evol.* 91, 17–26.
648
649 Casteleyn, G., Leliaert, F., Backeljau, T., Debeer, A.-E., Kotaki, Y., Rhodes, L., Lundholm,
650 N., Sabbe, K., Vyverman, W., 2010. Limits to gene flow in a cosmopolitan marine planktonic
651 diatom. *Proc. Natl. Acad. Sci.* 107, 12952–12957. doi:10.1073/pnas.1001380107
652
653 Castellani C., Licandro, P., Fileman, E., di Capua, I., Mazzocchi, M.G., 2016. *Oithona similis*
654 likes it cool: evidence from two long-term time series. *J. Plankton Res.* 38, 703-717.
655 doi:10.1093/plankt/fbv104
656
657 Castellani, C., Robinson, C., Smith, T., Lampitt, R.S., 2005. Temperature affects respiration
658 rate of *Oithona similis*. *Mar. Ecol. Progr. Ser.* 285, 129–135. doi:10.3354/meps285129
659
660 Cepeda, G.D., Blanco-Bercial, L., Bucklin, A., Berón, C.M., Viñas, M.D., 2012. Molecular
661 Systematic of Three Species of *Oithona* (Copepoda, Cyclopoida) from the Atlantic Ocean:
662 Comparative Analysis Using 28S rDNA. *PLoS ONE* 7, e35861.
663 doi:10.1371/journal.pone.0035861.s001
664
665 Cepeda, G.D., Sabatini, M.E., Scioscia, C.L., Ramírez, F.C., Viñas, M.D., 2016. On the
666 uncertainty beneath the name *Oithona similis* Claus, 1866 (Copepoda, Cyclopoida). *ZooKeys*
667 552, 1-15.
668
669 Checkley, D.M., 1982. Selective feeding by Atlantic herring (*Clupea harengus*) larvae on
670 zooplankton in natural assemblages. *Mar. Ecol. Progr. Ser.* 9, 245–253.
671
672 Chen, G., Hare, M.P., 2011. Cryptic diversity and comparative phylogeography of the
673 estuarine copepod *Acartia tonsa* on the US Atlantic coast. *Mol. Ecol.* 20, 2425–2441.
674 doi:10.1111/j.1365-294X.2011.05079.x
675
676 Chust, G., Castellani, C., Licandro, P., Ibaibarriaga, L., Sagarminaga, Y., Irigoien, X., 2014.
677 Are *Calanus* spp. shifting poleward in the North Atlantic? A habitat modelling approach. *ICES*
678 *J. Mar. Sci.* 71, 241–253.
679
680 Claus, C.F.W., 1866. Die Copepoden-Fauna von Nizza, ein Beitrag zur Charakteristik der
681 Formen und deren Abänderungen. Elwert'sche Universitäts-Buchhandlung, Marburg &
682 Leipzig.
683
684 Claus, C.F.W., 1863. Die frei lebenden Copepoden: Mit besonderer Berücksichtigung der
685 Fauna Deutschlands, der Nordsee und des Mittelmeeres. W. Engelmann.
686
687 Clement, M.J., Posada, D., Crandall, K.A. 2000. TCS: A computer program to estimate gene
688 genealogies. *Mol. Ecol.* 9, 1657-1659.
689
690 Cornils, A., Blanco-Bercial, L., 2013. Phylogeny of the Paracalanidae Giesbrecht, 1888
691 (Crustacea: Copepoda: Calanoida). *Mol. Phylogenet. Evol.* 69, 861–872.
692
693 Cornils, A., Held, C., 2014. Evidence of cryptic and pseudocryptic speciation in the
694 *Paracalanus parvus* species complex (Crustacea, Copepoda, Calanoida). *Front. Zool.* 11, 1–
695 17. doi:10.1186/1742-9994-11-19
696

697 Cornils A., Wend-Heckmann, B., 2015. First report of the planktonic copepod *Oithona*
698 *davisae* in the northern Wadden Sea (North Sea): Evidence for recent invasion? Helgoland
699 Mar. Res. 69, 243-248.
700

701 Darling, K.F., Kucera, M., Wade, C.M., 2007. Global molecular phylogeography reveals
702 persistent Arctic circumpolar isolation in a marine planktonic protist. Proc. Natl. Acad. Sci.
703 104, 5002–5007.
704

705 de Vargas, C., Audic, S., Henry, N., Decelle, J., Mahé, F., Logares, R., Lara, E., Berney, C.,
706 Le Bescot, N., Probert, I., Carmichael, M., Poulain, J., Romac, S., Colin, S., Aury, J.M.,
707 Bittner, L., Chaffron, S., Dunthorn, M., Engelen, S., Flegontova, O., Guidi, L., Horak, A.,
708 Jaillon, O., Lima-Mendez, G., Luke, J., Malviya, S., Morard, R., Mulo, M., Scalco, E., Siano,
709 R., Vincent, F., Zingone, A., Dimier, C., Picheral, M., Searson, S., Kandels-Lewis, S., Tara
710 Oceans Coordinators, Acinas, S.G., Bork, P., Bowler, C., Gorsky, G., Grimsley, N., Hingamp,
711 P., Iudicone, D., Not, F., Ogata, H., Pesant, S., Raes, J., Sieracki, M.E., Speich, S.,
712 Stemmann, L., Sunagawa, S., Weissenbach, J., Wincker, P., Karsenti, E., Boss, E., Follows,
713 M., Karp-Boss, L., Krzic, U., Reynaud, E.G., Sardet, C., Sullivan, M.B., Velayoudon, D.,
714 2015. Eukaryotic plankton diversity in the sunlit ocean. Science 348, 1261605–1261605.
715 doi:10.1126/science.1261605
716

717 Dietz L., Arango, C.P., Dömel, J.S., Halanynch, K.M., Harder, A.M., Held, C., Mahon, A.R.,
718 Mayer, C., Melzer, R.R., Rouse, G.W., Weis, A., Wilson, N.G., Leese, F., 2015. Regional
719 differentiation and extensive hybridization between mitochondrial clades of the Southern
720 Ocean giant sea spider *Colossendeis megalonyx*. R. Soc. open sci. 2: 140424.
721

722 Drummond, A.J., Suchard, M.A., Xie, D., Rambaut, A., 2012. Bayesian Phylogenetics with
723 BEAUti and the BEAST 1.7. Mol. Biol. Evol. 29, 1969–1973. doi:10.1093/molbev/mss075
724

725 Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., 1994. DNA primers for
726 amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan
727 invertebrates. Mol. Mar. Biol. Biotechnol. 3, 294–299.
728

729 Gallienne, C.P., Robins, D.B., 2001. Is *Oithona* the most important copepod in the world's
730 oceans? J. Plankton Res. 23, 1421–1432.
731

732 Halbert, K.M.K., Goetze, E., Carlon, D.B., 2013. High cryptic diversity across the global
733 range of the migratory planktonic copepods *Pleuromamma piseki* and *P. gracilis*. PLoS ONE
734 8, e77011. doi:10.1371/journal.pone.0077011.s006
735

736 Harvey, T.H.P., Pedder, B.E., 2013. Copepod Mandible Palynomorphs From the Nolichucky
737 Shale (Cambrian, Tennessee): Implications for the Taphonomy and Recovery of Small
738 Carbonaceous Fossils. PALAIOS 28, 278–284. doi:10.2110/palo.2012.p12-124r
739

740 Hebert, P.D.N., Cywinska, A., Ball, S.L., deWaard, J.R., 2003. Biological identifications
741 through DNA barcodes. Proc. R. Soc. B 270, 313–321. doi:10.1098/rspb.2002.2218
742

743 Held, C., 2000. Phylogeny and biogeography of serolid isopods (Crustacea, Isopoda,
744 Serolidae) and the use of ribosomal expansion segments in molecular systematics. Mol.
745 Phylogenet. Evol. 15, 165–178.
746

747 Hill R., Allen, L.D., Bucklin, A., 2001. Multiplexed species-specific PCR protocol to
748 discriminate four N. Atlantic *Calanus* species, with an mtCOI gene tree for ten *Calanus*
749 species. Mar. Biol. 139, 279–287.
750

751 Hirche, H.-J., Kosobokova, K., 2007. Distribution of *Calanus finmarchicus* in the northern
752 North Atlantic and Arctic Ocean—Expatriation and potential colonization. Deep-Sea Res. II

753 54, 2729–2747. doi:10.1016/j.dsr2.2007.08.006
754
755 Hopcroft, R.R., Kosobokova, K.N., Pinchuk, A.I., 2010. Zooplankton community patterns in
756 the Chukchi Sea during summer 2004. *Deep-Sea Res. II* 57, 27–39.
757 doi:10.1016/j.dsr2.2009.08.003
758
759 Hunt, B.P.V., Hosie, G.W., 2006a. The seasonal succession of zooplankton in the Southern
760 Ocean south of Australia, part I: The seasonal ice zone. *Deep-Sea Res. I* 53, 1182–1202.
761 doi:10.1016/j.dsr.2006.05.001
762
763 Hunt, B.P.V., Hosie, G.W., 2006b. The seasonal succession of zooplankton in the Southern
764 Ocean south of Australia, part II: The Sub-Antarctic to Polar Frontal Zones. *Deep-Sea Res. I*
765 53, 1203–1223. doi:10.1016/j.dsr.2006.05.002
766
767 Hunt, B.P.V., Nelson, R.J., Williams, B., McLaughlin, F.A., Young, K.V., Brown, K.A., Vagle,
768 S., Carmack, E.C., 2014. Zooplankton community structure and dynamics in the Arctic
769 Canada Basin during a period of intense environmental change (2004–2009). *J. Geophys.*
770 *Res.: Oceans* 119, 2518–2538. doi:10.1002/2013JC009156
771
772 Hunt, B.P.V., Strugnell, J., Bednarsek, N., Linse, K., Nelson, R.J., Pakhomov, E., Seibel, B.,
773 Steinke, D., Würzberg, L., 2010. Poles apart: The “bipolar” pteropod species *Limacina*
774 *helicina* is genetically distinct between the Arctic and Antarctic oceans. *PLoS ONE* 5, e9835.
775 doi:10.1371/journal.pone.0009835.g001
776
777 Jones, M., Ghoorah, A., Blaxter, M.L., 2011. jMOTU and Taxonator: Turning DNA barcode
778 sequences into annotated operational taxonomic units. *PLoS ONE* 6, e19259.
779 doi:10.1371/journal.pone.0019259.t002
780
781 Jungbluth M.J., Lenz, P.H., 2013. Copepod diversity in a subtropical bay based on a
782 fragment of the mitochondrial COI gene. *J. Plankton Res.* 35, 630–643.
783
784 Katoh, K., Standley, D.M., 2013. MAFFT Multiple Sequence Alignment Software Version 7:
785 Improvements in Performance and Usability. *Mol. Biol. Evol.* 30, 772–780.
786 doi:10.1093/molbev/mst010
787
788 Kiesling T.L., Wilkinson, E., Rabalais, J., Ortner, P.B., McCabe, M.M., Fell, J.W., 2002. Rapid
789 identification of adult and naupliar stages of copepods using DNA hybridization methodology.
790 *Mar. Biotechnol.* 4, 30–39.
791
792 Knowlton, N., Weigt, L.A., 1998. New dates and new rates for divergence across the Isthmus
793 of Panama. *Proc. R. Soc. B* 265, 2257–2263. doi:10.1098/rspb.1998.0568
794
795 Kosobokova, K., Hirche, H.J., 2000. Zooplankton distribution across the Lomonosov Ridge,
796 Arctic Ocean: species inventory, biomass and vertical structure. *Deep-Sea Res. I* 47, 2029–
797 2060.
798
799 Kozol, R., Blanco-Bercial, L., Bucklin, A., 2012. Multi-gene analysis reveals a lack of genetic
800 divergence between *Calanus agulhensis* and *C. sinicus* (Copepoda; Calanoida). *PLoS ONE*
801 7, e45710. doi:10.1371/journal.pone.0045710
802
803 Kulagin, D.N., Stupnikova, A.N., Neretina, T.V., Mogue, N.S., 2013. Spatial genetic
804 heterogeneity of the cosmopolitan chaetognath *Eukrohnia hamata* (Möbius, 1875) revealed
805 by mitochondrial DNA. *Hydrobiologia* 721, 197–207. doi:10.1007/s10750-013-1661-z
806
807 Lanfear, R., Calcott, B., Ho, S.Y.W., Guindon, S., 2012. PartitionFinder: Combined Selection
808 of Partitioning Schemes and Substitution Models for Phylogenetic Analyses. *Mol. Biol. Evol.*

809 29, 1695–1701. doi:10.1093/molbev/mss020
810
811 Leigh, J.W., Bryant, D. 2015. popart: full-feature software for haplotype network construction.
812 Methods Ecol. Evol. 6, 1110–1116.
813
814 Machida, R.J., Miya, M.U., Nishida, M., Nishida, S., 2004. Large-scale gene rearrangements
815 in the mitochondrial genomes of two calanoid copepods *Eucalanus bungii* and *Neocalanus*
816 *cristatus* (Crustacea), with notes on new versatile primers for the srRNA and COI genes.
817 Gene 332, 71–78. doi:10.1016/j.gene.2004.01.019
818
819 Machida, R.J., Tsuda, A., 2010. Dissimilarity of species and forms of planktonic *Neocalanus*
820 copepods using mitochondrial COI, 12S, nuclear ITS, and 28S gene sequences. PLoS ONE
821 5: e10278.
822
823 Madsen, S.D., Nielsen, T.G., Hansen, B.W., 2008. Annual population development and
824 production by small copepods in Disko Bay, western Greenland. Mar. Biol. 155, 63–77.
825 doi:10.1007/s00227-008-1007-y
826
827 Makino W., Tanabe, A.S., 2009. Extreme population genetic differentiation and secondary
828 contact in the freshwater copepod *Acanthodiaptomus pacificus* in the Japanese Archipelago.
829 Mol. Ecol. 18, 3699–3713.
830
831 Marin, V., 1987. The oceanographic structure of the eastern Scotia Sea—IV. Distribution of
832 copepod species in relation to hydrography in 1981. Deep-Sea Res. A 34, 105–121.
833
834 Marrone, F., Brutto, Lo, S., Hundsdoerfer, A.K., Arculeo, M., 2013. Overlooked cryptic
835 endemism in copepods: Systematics and natural history of the calanoid subgenus
836 *Occidodiaptomus* Borutzky 1991 (Copepoda, Calanoida, Diaptomidae). Mol. Phylogenet.
837 Evol. 66, 190–202. doi:10.1016/j.ympev.2012.09.016
838
839 Meier, R., Shiyang, K., Vaidya, G., Ng, P.K.L., 2006. DNA barcoding and taxonomy in
840 Diptera: a tale of high intraspecific variability and low identification success. Syst. Biol. 55,
841 715–728.
842
843 Miyamoto, H., Machida, R.J., Nishida, S., 2010. Genetic diversity and cryptic speciation of
844 the deep sea chaetognath *Caecosagitta macrocephala* (Fowler, 1904). Deep-Sea Res. II 57,
845 2211–2219. doi:10.1016/j.dsr2.2010.09.023
846
847 Miyamoto, H., Machida, R.J., Nishida, S., 2012. Global phylogeography of the deep-sea
848 pelagic chaetognath *Eukrohnia hamata*. Prog. Oceanogr. 104, 99–109.
849 doi:10.1016/j.pocean.2012.06.003
850
851 Monaghan, M.T., Wild, R., Elliot, M., Fujisawa, T., Balke, M., Inward, D.J.G., Lees, D.C.,
852 Ranaivosolo, R., Eggleton, P., Barraclough, T.G., Vogler, A.P., 2009. Accelerated Species
853 Inventory on Madagascar Using Coalescent-Based Models of Species Delineation. Syst.
854 Biol. 58, 298–311. doi:10.1093/sysbio/syp027
855
856 Nelson, R.J., Carmack, E.C., McLaughlin, F.A., Cooper, G.A., 2009. Penetration of Pacific
857 zooplankton into the western Arctic Ocean tracked with molecular population genetics. Mar.
858 Ecol. Progr. Ser. 381, 129–138.
859
860 Niehoff B., Madsen, S.D., Hansen, B.W., Nielsen, T.G., 2002. Reproductive cycles of three
861 dominant *Calanus* species in Disko Bay, West Greenland. Mar. Biol. 140, 567–576.
862
863 Nishida, S., 1985. Taxonomy and distribution of the family Oithonidae (Copepoda,
864 Cyclopoida) in the Pacific and Indian Oceans. Bull. Ocean Res. Inst., Tokyo 20, 1–167.

865
866 Norris, R.D., 2000. Pelagic species diversity, biogeography, and evolution. *Paleobiology* 26,
867 236–258.
868
869 Ortman, B.D., 2008. DNA barcoding the medusozoa and ctenophora. Ann Arbor: University
870 of Connecticut, pp. 1-122. <http://digitalcommons.uconn.edu/dissertations/AAI3345202>.
871
872 Palumbi, S.R., 1992. Marine speciation on a small planet. *Trends Ecol. Evol.* 7, 114-118.
873
874 Paradis, V., Sirois, P., Castonguay, M., Plourde, S., 2012. Spatial variability in zooplankton
875 and feeding of larval Atlantic mackerel (*Scomber scombrus*) in the southern Gulf of St.
876 Lawrence. *J. Plankton Res.* 34, 1064–1077. doi:10.1093/plankt/fbs063
877
878 Parent G.J., Plourde, S., Turgeon, J., 2012. Natural hybridization between *Calanus*
879 *finmarchicus* and *C. glacialis* (Copepoda) in the Arctic and Northwest Atlantic. *Limnol.*
880 *Oceanogr.* 57, 1057–1066.
881
882 Park, E.T., Ferrari, F.D., 2009. Species diversity and distributions of pelagic calanoid
883 copepods from the Southern Ocean, in: Krupnik, I., Lang, M.A., Miller, S.E. (Eds),
884 Smithsonian at the Poles: contributions to International Polar Year Science, pp. 143–180.
885
886 Pinkerton, M.H., Smith, A.N.H., Raymond, B., Hosie, G.W., Sharp, B., Leathwick, J.R.,
887 Bradford-Grieve, J.M., 2010. Spatial and seasonal distribution of adult *Oithona similis* in the
888 Southern Ocean: Predictions using boosted regression trees. *Deep-Sea Res. I* 57, 469–485.
889 doi:10.1016/j.dsr.2009.12.010
890
891 Pons, J., Barraclough, T., Gomez-Zurita, J., Cardoso, A., Duran, D., Hazell, S., Kamoun, S.,
892 Sumlin, W., Vogler, A.P., 2006. Sequence-based species delimitation for the DNA taxonomy
893 of undescribed insects. *Syst. Biol.* 55, 595–609. doi:10.1080/10635150600852011
894
895 Puillandre, N., Lambert, A., Brouillet, S., Achaz, G., 2012. ABGD, Automatic Barcode Gap
896 Discovery for primary species delimitation. *Mol. Ecol.* 21, 1864–1877. doi:10.1111/j.1365-
897 294X.2011.05239.x
898
899 R Development Core Team, 2015. R: A language and environment for statistical computing.
900 R Foundation for Statistical Computing, Vienna Austria. <http://www.R-project.org>.
901
902 Razouls, C., de Bovée, F., Kouwenberg, J., Desreumaux, N., 2016. Diversity and geographic
903 distribution of marine planktonic copepods. <http://copepodes.obs-banyuls.fr>
904
905 Reid, N.M., Carstens, B.C., 2012. Phylogenetic estimation error can decrease the accuracy
906 of species delimitation: a Bayesian implementation of the general mixed Yule-coalescent
907 model. *BMC Evol. Biol.* 12, 196. doi:10.1186/1471-2148-12-196
908
909 Richardson, A.J., 2008. In hot water: zooplankton and climate change. *ICES J. Mar. Sci.* 65,
910 279–295.
911
912 Ronquist F., Huelsenbeck J.P., Teslenko M. 2011. MrBayes version 3.2 Manual: Tutorials
913 and Model Summaries, pp 1–103. <http://mrbayes.sourceforge.net/download.php>
914
915 Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D.L., Darling, A., Höhna, S., Larget, B.,
916 Liu, L., Suchard, M.A., Huelsenbeck, J.P., 2012. MrBayes 3.2: Efficient bayesian
917 phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 539–
918 542. doi:10.1093/sysbio/sys029
919
920 Rosenberg, N.A., 2007. Statistical tests for taxonomic distinctiveness from observations of

921 monophyly. *Evolution* 61, 317–323. doi:10.1111/j.1558-5646.2007.00023.x

922

923 Sabatini, M., Kiørboe, T., 1994. Egg production, growth and development of the cyclopoid

924 copepod *Oithona similis*. *J. Plankton Res.* 16, 1329–1351.

925

926 Sars, G.O., 1918. An account of the Crustacea of Norway with short descriptions and figures

927 of all species. Volume VI: Copepoda. Cyclopoida (118 plates) pp. 1–776.

928

929 Selden, P.A., Huys, R., Stephenson, M.H., Heward, A.P., Taylor, P.N., 2010. Crustaceans

930 from bitumen clast in Carboniferous glacial diamictite extend fossil record of copepods. *Nat.*

931 *Comms.* 1, 1–6. doi:10.1038/ncomms1049

932

933 Shuvalov, V.S., 1972. Geographical variability of some species of the family Oithonidae

934 (Copepoda, Cyclopoida), in.: Zvereva, Z.A. (Ed.), Geographic and seasonal variability of

935 marine plankton. *Stud. mar. Fauna* 12, 169–186.

936

937 Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

938 large phylogenies. *Bioinformatics Applications Note* 30, 1312–1313.

939 doi:10.1093/bioinformatics/btu033/-/DC1

940

941 Strass, V. H. 2010. Physical oceanography during POLARSTERN cruise ANT-XXIV/2

942 (SCACE). Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research,

943 Bremerhaven, doi:10.1594/PANGAEA.733665

944

945 Stupnikova, A.N., Molodtsova, T.N., Mugue, N.S., Neretina, T.V., 2013. Genetic variability of

946 the *Metridia lucens* complex (Copepoda) in the Southern Ocean. *J. Mar. Syst.* 128, 175–184.

947 doi:10.1016/j.jmarsys.2013.04.016

948

949 Tamura, K., Stecher, G., Peterson, D., FilipSKI, A., Kumar, S., 2013. MEGA6: Molecular

950 Evolutionary Genetics Analysis Version 6.0. *Mol. Biol. Evol.* 30, 2725–2729.

951 doi:10.1093/molbev/mst197

952

953 Thum, R.A., Harrison, R.G., 2009. Deep genetic divergences among morphologically similar

954 and parapatric *Skistodiatomus* (Copepoda: Calanoida: Diaptomidae) challenge the

955 hypothesis of Pleistocene speciation. *Biol. J. Linnean Soc.* 96, 150–165.

956

957 Turner, J.T., 2004. The importance of small planktonic copepods and their roles in pelagic

958 marine food webs. *Zool. Stud.* 43, 255–266.

959

960 Ueda, H., Yamaguchi, A., Saitoh, S.-I., Sakaguchi, S.O., Tachihara, K., 2011. Speciation of

961 two salinity-associated size forms of *Oithona dissimilis* (Copepoda: Cyclopoida) in estuaries.

962 *J. Nat. Hist.* 45, 2069–2079.

963

964 Ward, P., Hirst, A.G., 2007. *Oithona similis* in a high latitude ecosystem: abundance,

965 distribution and temperature limitation of fecundity rates in a sac spawning copepod. *Mar.*

966 *Biol.* 151, 1099–1110. doi:10.1007/s00227-006-0548-1

967

968 Weiner, A., Aurahs, R., Kurasawa, A., Kitazato, H., Kucera, M., 2012. Vertical niche

969 partitioning between cryptic sibling species of a cosmopolitan marine planktonic protist. *Mol.*

970 *Ecol.* 21, 4063–4073. doi:10.1111/j.1365-294X.2012.05686.x

971

972 Xia, X., 2013. DAMBE5: a comprehensive software package for data analysis in molecular

973 biology and evolution. *Mol. Biol. Evol.* 30, 1720–1728. doi:10.1093/molbev/mst064/-/DC1

974

975 Xia, X., Lemey, P., 2009. Assessing substitution saturation with DAMBE, in: Lemey, P.,

976 Salemi, M., Vandamme, A.-M. (Eds.) *The phylogenetic handbook: a practical approach to*

- 977 DNA and protein phylogeny. Cambridge University Press, Cambridge, pp. 615–630.
978
- 979 Zachos, J., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and
980 aberrations in global climate 65 Ma to present. *Science* 292, 686–693.
981
- 982 Zamora-Terol, S., Kjellerup, S., Swalethorp, R., Saiz, E., Nielsen, T.G., 2014. Population
983 dynamics and production of the small copepod *Oithona* spp. in a subarctic fjord of West
984 Greenland. *Polar Biol.* 37, 953–965. doi:10.1007/s00300-014-1493-y
985
- 986 Zamora-Terol, S., Nielsen, T., Saiz, E., 2013. Plankton community structure and role of
987 *Oithona similis* on the western coast of Greenland during the winter-spring transition. *Mar.*
988 *Ecol. Progr. Ser.* 483, 85–102. doi:10.3354/meps10288

Figure captions

Figure 1: Station map for *Oithona similis* sampling locations. The samples from the type localities of *Oithona similis* (Mediterranean Sea) and *Oithona helgolandica* (North Sea) are marked in bold letters. Yellow dots (North Pacific Ocean) and the purple dot (Polarfront) south of Australia are based on location derived from GenBank COI sequences (Table 1). The geographic information of the 28S sequences from GenBank are not included, but can be viewed in Cepeda et al. (2012).

Figure 2: Bayesian phylogenetic tree of COI haplotypes for the genus *Oithona*. Bayesian posterior probabilities (BPP) and maximum likelihood bootstrap values (BS) are placed at the above and below nodes. Only values above 0.9 (BPP) and 50 (BS) are displayed. Colour bars represent the sampling localities of *Oithona similis* s.l. with the number of sequences (n). *O. similis* s.l. specimens from the type localities are marked in bold letters.

Figure 3: Bayesian phylogenetic tree of 28S alleles for the genus *Oithona*. Bayesian posterior probabilities (BPP) and maximum likelihood bootstrap values (BS) are placed at the above and below nodes. Only values above 0.9 (BPP) and 50 (BS) are displayed. Colour bars represent the sampling localities of *Oithona similis* s.l. with the number of sequences (n). *O. similis* s.l. specimens from the type localities are marked in bold letters.

Figure 4: TCS haplotype networks of COI and 28S for *O. similis* s.l., generated using PopArt. Numbers represent mutational steps that separate haplotypes. Circle size of the haplotypes indicate relative haplotype frequency. Colours represent the sampling locations. The number of total haplotypes differs from Table 3 due to PopArt's exclusion of nucleotide positions with missing and ambiguous data. 27 COI sequences with less than 400 bp and eight 28S sequences with ambiguities were removed to get a higher resolution of the phylogeography of the discovered lineages.

Figure 5: Vertical distribution of the specimens of the cryptic species PF and ANT at the station of the Polar Front in the Atlantic sector. Red lines show the vertical temperature profiles of the sampled stations (from Strass, 2010, see Supplementary Table for stations).

Figure 6: Relative timing of the diversification of *O. similis* s.l. based on COI with terminal lineage names and 95% HPD node bars. Arctic lineages (ARK) are marked in bold letters.

Figure 1

