



# The effect of cosmic dust on Southern Ocean biogeochemistry is small but non-negligible

C. Völker<sup>5</sup>, J. Plane<sup>1</sup>, S. Dhomse<sup>1</sup>, J.-F. Lamarque<sup>2</sup>, M. Long<sup>2</sup>, A. Sáiz Lopez<sup>3</sup>, and A. Tagliabue<sup>4</sup>

1:Univ. Leeds, UK, 2:NCAR, USA, 3:CSIC, Spain, 4:Univ. Liverpool, UK, 5:AWI, Germany

## 1. Introduction



Lithogenic dust deposition flux in  $\text{g m}^{-2} \text{yr}^{-1}$  (Jickells et al. 2005)

Dust deposition over the ocean varies over more than five orders of magnitude, with high deposition rates downwind of deserts, but extremely low deposition over the South Pacific and the Southern Ocean. Although total cosmogenic dust deposition is much smaller, it is more evenly distributed and may be an important Fe source in these regions (Johnson, 2001). We estimated Fe input from a new model climatology of cosmic dust deposition and estimated its effect on marine biogeochemical cycles with global models.

## 2. Methods


Modeled deposition flux for meteoric smoke particles from Dhomse et al., 2013 was converted to Fe flux, assuming 100% solubility, and added to the Fe flux from lithogenic dust. Several global biogeochemical models of the ocean have then been integrated over at least 1000 years with and without the cosmic dust flux to study the reaction of the marine Fe cycle and biological productivity.

The models are REcoM-MITgcm (Ye and Völker, 2017), CESM-BEC (Moore et al., 2013) and ORCA-PISCES (Aumont et al., 2015)

The models differ in several aspects of their iron cycle, e.g. the strength of the sedimentary iron source or the existence of an explicit description of iron in sea-ice. Results shown here are from REcoM


## 3. Results

### 1) Comparing Fe flux from cosmic and terrestrial dust




The iron flux from meteoric smoke particles (MSPs), albeit on average smaller, is distributed more homogeneously than the flux from terrestrial dust (note the different scales). In most regions it is only a small addition, but in the South Pacific and the Indo-Pacific parts of the Southern Ocean, it can actually dominate over terrestrial Fe flux. Dust flux from Albani et al, 2014, has been converted to iron flux, assuming a constant solubility of 2% and crustal Fe abundance of 3.5%.

### 3) Effect on phytoplankton productivity



The increased dFe concentrations lead only to slight enhancement of phytoplankton net primary production (NPP) in iron-limited regions, and small reductions in NPP in the nitrogen-limited South Atlantic subtropical gyre, driven by macronutrient uptake in the Fe-limited regions. Production changes are largest near the boundaries between N- and Fe-limited regions. Overall, the Fe input from MSPs leads to an increase of global NPP by 2% and of global export by 1%.

### 2) Effect on dissolved iron



The additional flux of Fe from MSPs results in an increase of dissolved iron (dFe) that is mostly smaller than 0.1 nM. Although relatively, the MPS iron flux is largest in the Southern Ocean, the absolute changes in dFe are largest in the subtropics.

	Without cosmic dust particles	With cosmic dust particles
total dFe deposition $10^9 \text{ g Fe year}^{-1}$	317	329
sediment dFe source $10^9 \text{ g Fe year}^{-1}$	271	
hydrothermal dFe $10^9 \text{ g Fe year}^{-1}$	50	
residence time years	69	
global net primary production $\text{Pg C year}^{-1}$	44.33	45.38
global export production $\text{Pg C year}^{-1}$	9.71	9.80

## 4. Discussion

The additional iron flux to the ocean from the deposition of meteoric smoke particles is a significant part of the total deposition flux in the South Pacific and parts of the iron-limited Southern Ocean.

Nevertheless, the effect on biological productivity, both locally and globally is relatively modest. This is probably because in the regions where MSPs contribute disproportionately, the deposition flux of iron is less important than the flux of iron from below, through upwelling and the annual mixed layer cycle.

The magnitude of other important iron sources to the ocean (sediment diagenesis and hydrothermalism) is still unclear, which causes different biogeochemical models to have widely different Fe residence times (Tagliabue et al., 2016).

It is therefore important to analyse the robustness of our results with a suite of different ocean biogeochemical models. This is currently underway.

## 5. References

- Aumont, O., Ethé, C., Tagliabue, A., Bopp, L., & Gehlen, M. (2015). PISCES-v2: An ocean biogeochemical model for carbon and ecosystem studies. *Geoscientific Model Development*, 8(8), 2465–2513. <https://doi.org/10.5194/gmd-8-2465-2015>
- Dhomse, S. S., R. W. Saunders, W. Tian, M. P. Chipperfield, and J. M. C. Plane (2013). Plutonium-238 observations as a test of modeled transport and surface deposition of meteoric smoke particles. *Geophysical Research Letters*, 40(16), 4454–4458. doi: 10.1002/grl.50840
- Johnson, K. S. (2001). Iron supply and demand in the upper ocean: Is extraterrestrial dust a significant source of bioavailable iron? *Global Biogeochemical Cycles*, 15(1), 61–63. doi:10.1029/2000GB001295
- Moore, J. K., Lindsay, K., Doney, S. C., Long, M. C., & Misumi, K. (2013). Marine ecosystem dynamics and biogeochemical cycling in the Community Earth System Model [CESM1 (BGC)]: Comparison of the 1990s with the 2090s under the RCP4.5 and RCP8.5 scenarios. *Journal of Climate*, 26(23), 9291–9312.
- Plane, J. M. C. (2012). Cosmic dust in the earth's atmosphere. *Chemical Society Reviews*, 41(19), 6507. doi: 10.1039/c2cs35132c
- Tagliabue, A., O. Aumont, R. DeAth, J. P. Dunne, S. Dutkiewicz, E. Galbraith, K. Misumi, J. K. Moore, A. J. Ridgwell, E. Sherman, C. Stock, M. Vichi, C. Völker, and A. Yool (2016). How well do global ocean biogeochemistry models simulate dissolved iron distributions? *Global Biogeochemical Cycles*, 30. doi: 10.1002/2015GB005289
- Ye, Y., and C. Völker (2017). On the Role of Dust-Deposited Lithogenic Particles for Iron Cycling in the Tropical and Subtropical Atlantic. *Global Biogeochemical Cycles*, 31(10), 1543–1558. doi: 10.1002/2017GB005663.