

Minireview

Methyl/alkyl-coenzyme M reductase-based anaerobic alkane oxidation in archaea

Yinzhaio Wang ^{1,2} Gunter Wegener ^{3,4}S. Emil Ruff^{5,6} and Fengping Wang^{1,7,8*}

¹State Key Laboratory of Microbial Metabolism, School of Life Sciences and Biotechnology, Shanghai Jiao Tong University, Shanghai, 200240, China.

²State Key Laboratory of Ocean Engineering, School of Naval Architecture, Ocean & Civil Engineering, Shanghai Jiao Tong University, Shanghai, 200240, China.

³Max Planck Institute for Marine Microbiology, Bremen, Germany.

⁴MARUM, Center for Marine Environmental Sciences, University of Bremen, Bremen, Germany.

⁵Ecosystems Center, Marine Biological Laboratory, Woods Hole, MA, USA.

⁶J. Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological Laboratory, Woods Hole, MA, USA.

⁷School of Oceanography, Shanghai Jiao Tong University, Shanghai, 200240, China.

⁸Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), Zhuhai, Guangdong, China.

Summary

Methyl-coenzyme M reductase (MCR) has been originally identified to catalyse the final step of the methanogenesis pathway. About 20 years ago anaerobic methane-oxidizing archaea (ANME) were discovered that use MCR enzymes to activate methane. ANME thrive at the thermodynamic limit of life, are slow-growing, and in most cases form syntrophic consortia with sulfate-reducing bacteria. Recently, archaea that have the ability to anaerobically oxidize non-methane multi-carbon alkanes such as ethane and *n*-butane were described in both enrichment cultures and environmental samples. These anaerobic multi-carbon alkane-oxidizing archaea (ANKA) use enzymes

homologous to MCR named alkyl-coenzyme M reductase (ACR). Here we review the recent progresses on the diversity, distribution and functioning of both ANME and ANKA by presenting a detailed MCR/ACR-based phylogeny, compare their metabolic pathways and discuss the gaps in our knowledge of physiology of these organisms. To improve our understanding of alkane oxidation in archaea, we identified three directions for future research: (i) expanding cultivation attempts to validate omics-based metabolic models of yet-uncultured organisms, (ii) performing biochemical and structural analyses of key enzymes to understand thermodynamic and steric constraints and (iii) investigating the evolution of anaerobic alkane metabolisms and their impact on biogeochemical cycles.

Introduction

Methane (CH₄) is the simplest and most abundant hydrocarbon on Earth (Wahlen, 1993) and a greenhouse gas that substantially influences the Earth's climate (Conrad, 2009; Yvon-Durocher *et al.*, 2014). Most of the methane in natural environments, such as marine sediments, cold seeps, wetlands, freshwater lakes and hot springs, is produced by methanogenic archaea (Reeburgh, 2007; Thauer, 2019). Methane is also a component of subsurface-derived natural gas originating from deep marine and terrestrial reservoirs. These natural gases are a diverse mix of methane and short-chain alkanes including ethane, propane, butane, pentane and hexane produced by thermocatalytic decomposition of organic matters in geothermally heated sediments (Stolper *et al.*, 2014).

Both methane and other alkanes can be utilized by microorganisms as energy and carbon sources. Under oxic conditions, microorganisms activate alkanes using methane or alkane monooxygenases that produce methanol or other alcohols as primary intermediates (Wang *et al.*, 2017). Such monooxygenases are key enzymes present in numerous bacterial lineages, and strains of almost 30 genera have been cultured that oxidize methane or other alkanes (Kallistova *et al.*, 2017; Dedysh and

Received 18 December, 2019; revised 28 April, 2020; accepted 30 April, 2020. **For correspondence. E-mail fengpingw@sjtu.edu.cn; Telephone: 021-34204503; Fax: 021-34207205

Knief, 2018). These aerobic alkane oxidizers are the dominant alkanotrophs in oxic and hypoxic soils, surface sediments and aquatic environments (Kallistova *et al.*, 2017). However, alkanes are also consumed in anoxic habitats such as marine or freshwater sediments, cold seeps and subsurface ecosystems (reviewed in Evans *et al.*, 2019; Bhattarai *et al.*, 2019). For instance, it was estimated that the anaerobic oxidation of methane (AOM) removes around 80% of the methane in marine sediments (Reeburgh, 2007; Knittel and Boetius, 2009; Boetius and Wenzhöfer, 2013). To our current knowledge, the anaerobic oxidation of methane and ethane is performed exclusively by archaea (Chen *et al.*, 2019; reviewed by Thauer, 2019), while the anaerobic oxidation of propane and butane can be performed by both archaea (Laso-Pérez *et al.*, 2016; Wang *et al.*, 2019a) and bacteria (Kniemeyer *et al.*, 2007; Jaekel *et al.*, 2013). The anaerobic oxidation of longer chain alkanes has only been described for bacteria (Aeckersberg *et al.*, 1991; Coates *et al.*, 1997), yet metagenomic surveys indicate a large unexplored potential for alkane degradation in the archaeal domain (Borrel *et al.*, 2019; Laso-Pérez *et al.*, 2019; Wang *et al.*, 2019b). In this review we mainly present an updated diversity of anaerobic alkane-oxidizing archaea, illustrate their global distribution using public metagenomic data sets from natural environments and summarize the biochemistry of anaerobic methane and multi-carbon alkane-oxidizing pathways.

Diversity and distribution of anaerobic alkane-oxidizing archaea

All so far cultured anaerobic methane, ethane, propane and *n*-butane-oxidizing archaea belong to the phylum Euryarchaeota (Table 1, Nauhaus *et al.*, 2007; Holler *et al.*, 2011; Laso-Pérez *et al.*, 2016; Chen *et al.*, 2019; Hahn *et al.*, 2020). They are closely related to methanogens and hence contain most genes of the methanogenesis pathway. Yet, anaerobic alkane-oxidizing archaea use the methanogenesis pathway in a reverse direction, thereby consuming alkanes instead of producing them. The key enzyme of the reverse methanogenesis pathway is a methyl-coenzyme M reductase (MCR) or alkyl-coenzyme M reductase (ACR) that activates the methane or other alkanes as methyl-CoM or alkyl-CoM respectively (Thauer, 2019). Global surveys based on 16S rRNA gene sequences showed that anaerobic methane-oxidizing archaea (ANME) are widely distributed in marine cold seep environments (Knittel *et al.*, 2005; Ruff *et al.*, 2015). In most ANME and anaerobic multi-carbon alkane-oxidizing archaea (here abbreviated as ANKA), the MCR or ACR-encoding genes are well conserved and can be used as phylogenetic markers (Friedrich, 2005; Evans *et al.*, 2019). We

screened public metagenomic data sets for MCR and ACR sequences and found that both ANME and ANKA are widely distributed across the globe, being particularly abundant at methane- and alkane-rich ecosystems (Fig. 1). We show that different types of anaerobic alkane-oxidizing archaea appear in distinct ecosystems or distinct niches in the same ecosystem.

Diversity of anaerobic methane-oxidizing archaea

ANME belong to two orders in the phylum Euryarchaeota, i.e., *Ca.* Methanophagales (ANME-1) and Methanosarcinales (including ANME-2a/b/c/d, ANME-3; Fig. 2). ANME-1 were originally described at cold seeps (Hinrichs *et al.*, 1999; Michaelis *et al.*, 2002), but they thrive in a wide range of environments particularly in marine and lacustrine sediments (Fig. 1). In marine sediments, ANME-1 often dominate deeper parts of sulfate–methane transition zones (SMTZ). Additionally, thermophilic ANME-1 subtypes occur in hydrothermally heated sediments of the Guaymas Basin (Teske *et al.*, 2002; Holler *et al.*, 2011; Wang *et al.*, 2019a). In most cases, ANME-1 form consortia with the sulfate-reducing *Deltaproteobacteria* of the SEEP-SRB1/2 clade or with *Ca.* Desulfofervidus, the only isolated partner bacterium involved in AOM to date (Holler *et al.*, 2011; Krukenberg *et al.*, 2016). Occasionally, ANME-1 occur without obvious partners (Orphan *et al.*, 2002; Ruff *et al.*, 2016), and in sulfate-depleted horizons below SMTZ. These observations indicate that ANME are capable to use alternative electron acceptors or thrive as methanogens (Lloyd *et al.*, 2011; Bertram *et al.*, 2013; Niu *et al.*, 2017; Beulig *et al.*, 2019). Yet, to date ANME-1 with alternative lifestyles have not been cultured.

Archaea of the ANME-2a/b clade dominate most marine cold seeps (Fig. 1; Orphan *et al.*, 2002; Mills *et al.*, 2003; Wegener *et al.*, 2008; Ruff *et al.*, 2015). ANME-2a/b preferentially occur in cold or moderately temperate environments and in enrichment cultures, they frequently outcompete other ANME clades (Zhang *et al.*, 2011; Wegener *et al.*, 2016). The MCR from ANME-2c is closely related to that of ANME-2a/b (Fig. 2), and the two ANME clades occupy similar ecological niches (Felden *et al.*, 2014; Krukenberg *et al.*, 2018). They also form dense consortia with sulfate-reducing SEEP-SRB1 or SEEP-SRB2 partner bacteria (Schreiber *et al.*, 2010; Kleindienst *et al.*, 2012).

Ca. Methanoperedenaceae (ANME-2d) are closely related to ANME-2a/b on the McrA phylogenetic tree (Fig. 2), yet are capable of thriving without partner bacteria (Haroon *et al.*, 2013). All cultivated members of this group and most environmental sequences derive from freshwater settings (Fig. 1; Cui *et al.*, 2015; Welte *et al.*, 2016). *Ca.* Methanoperedenens nitroreducens encodes

Table 1 MCR/ACR-based anaerobic alkane-oxidizing archaea.

Organism ^a	MCR/ACR function	Electron acceptor	References
ANME-1 ^b	Methane activation	SO ₄ ²⁻	Hinrichs <i>et al.</i> , 1999; Boetius <i>et al.</i> , 2000
ANME-2a/b		SO ₄ ²⁻	Orphan <i>et al.</i> , 2002; Wang <i>et al.</i> , 2014
ANME-2c		SO ₄ ²⁻	Kruenberg <i>et al.</i> , 2018; Wang <i>et al.</i> , 2019a
ANME-2d ^c		NO ₃ ⁻ , Fe ³⁺ , Mn ⁴⁺	Haroon <i>et al.</i> , 2013; Arshad <i>et al.</i> , 2015; Cai <i>et al.</i> , 2018; Leu <i>et al.</i> , 2020
ANME-3	<i>n</i> -Butane/propane activation	SO ₄ ²⁻	Niemann <i>et al.</i> , 2006; Omoregie <i>et al.</i> , 2008
<i>Ca.</i> Syntrophoarchaeum		SO ₄ ²⁻	Laso-Pérez <i>et al.</i> , 2016; Wang <i>et al.</i> , 2019a
<i>Ca.</i> Argoarchaeum		SO ₄ ²⁻	Chen <i>et al.</i> , 2019
<i>Ca.</i> Ethanoperedens	Ethane activation	Unknown	Hahn <i>et al.</i> , 2020
<i>Ca.</i> Verstraetearchaeota ^d			Unknown
<i>Ca.</i> Nezharchaeota	Unknown, potentially in long-chain alkane metabolism	Unknown	Wang <i>et al.</i> , 2019b; Hua <i>et al.</i> , 2019
<i>Ca.</i> Korarchaeota			Wang <i>et al.</i> , 2019b; McKay <i>et al.</i> , 2019
Thaumarchaeota			Hua <i>et al.</i> , 2019
Archaeoglobi ^e			Wang <i>et al.</i> , 2019b; Colman <i>et al.</i> , 2019
<i>Ca.</i> Bathyarchaeota ^d			Evans <i>et al.</i> , 2015
<i>Ca.</i> Hadesarchaeota			Wang <i>et al.</i> , 2019b; Hua <i>et al.</i> , 2019
<i>Ca.</i> Helarchaeota			Seitz <i>et al.</i> , 2019
Archaeoglobi ^e			Wang <i>et al.</i> , 2019b; Boyd <i>et al.</i> , 2019
<i>Ca.</i> Methanoliparia			Borrel <i>et al.</i> , 2019; Laso-Pérez <i>et al.</i> , 2019

^aReported at different phylogenetic levels.

^bANME-1 is now classified as the novel order *Candidatus* (*Ca.*) Methanophagales.


^cANME-2d is now classified as the family *Ca.* Methanoperedenaceae.

^d*Ca.* Verstraetearchaeota and *Ca.* Bathyarchaeota were the first discovered organisms with these MCR-encoding genes outside the Euryarchaeota phylum (Evans *et al.*, 2015; Vanwonterghem *et al.*, 2016). Due to unknown function, we refer to the Verstraetearchaeota-type and Bathyarchaeota-type MCRs (see Fig. 2).

^eDifferent MAGs of Archaeoglobi contain either Verstraetearchaeota-type MCR or Bathyarchaeota-type MCR-encoding genes.

nitrate reductases (*nar*) and couples methane oxidation to the reduction of nitrate to nitrite. Hence, it does not require a partner bacterium, but it benefits from partnerships with nitrite-reducing partners (Haroon *et al.*, 2013;

Arshad *et al.*, 2015). Recently it has been discovered that other *Ca.* Methanoperedens strains coupled AOM to the reduction of iron oxides and other metal oxides (Cai *et al.*, 2018; Liang *et al.* 2019; Leu *et al.*, 2020).

**Fig 1** Abundances of gene sequences affiliating with eleven types of McrA/AcrA encoding genes.

The sequences derived from public metagenomes in the NCBI SRA database. They originated from hydrocarbon seeps ($n = 17$), marine sediments ($n = 10$), freshwater sediments ($n = 10$) and hot springs ($n = 8$). For comparison, all abundance values are normalized, i.e., the abundance of each McrA/AcrA type in a metagenome was divided by the arithmetical mean of this type across all studied metagenomes. Sequence abundance is shown on a pseudo-log scale to include zeros, i.e., data sets/ecosystems in which certain McrA/AcrA gene sequences were not detected. Many potential McrA sequences have best hits with AcrA or Bathyarchaeota-type McrA sequences, but with <60% amino acid identities. Hence, they are classified as Bathyarchaeota-like or AcrA-like sequences respectively.




Fig 2 McrA/AcrA phylogenetic tree. The tree was constructed based on alignments of McrA/AcrA sequences generated using MAFFT and then filtered with trimAl. The tree was built by the IQ-Tree method with model LG + C60 + F + G using 1000 bootstrap iterations. Black circles, triangles or squares show experimentally confirmed anaerobic methane, ethane and *n*-butane-oxidizing archaea respectively. Note that the two cultured *Ca. Syntrophoarchaeum* strains encode four different ACRs.

Members of the ANME-3 clade dominate methane-rich arctic mud volcanoes (Niemann *et al.*, 2006; Lee *et al.*, 2019; Ruff *et al.*, 2019), cold seep sediments in the Eastern Mediterranean (Omorigie *et al.*, 2008) as well as sediments of the Marine Lake Grevelingen (Cassarini *et al.*, 2019). ANME-3 usually forms consortia with bacteria related to the genus *Desulfobulbus*. ANME-3 has not been cultured, and genomic information for this clade is lacking. The clade is closely related to methanogens of the genus *Methanococcoides* (Fig. 2) indicating that the capability to reverse methanogenesis may have evolved independently in several archaeal lineages.

Diversity of anaerobic multi-carbon alkane-oxidizing archaea

More than 15years after the discovery of ANME, a thermophilic enrichment culture was established that

contained consortia of syntrophic archaea and bacteria mediating the anaerobic oxidation of *n*-butane (Laso-Pérez *et al.*, 2016). The archaea were identified as *Ca. Syntrophoarchaeum* (formerly GoM-Arc87 clade) in the new order *Ca. Syntropharchaeales* (Adam *et al.*, 2017). Their partner bacterium is *Ca. Desulfofervidus*, which also form consortia with thermophilic ANME-1. *Ca. Syntrophoarchaeum* contains four *mcr*-like gene operons that are highly divergent from those of methanogens and methanotrophs based on amino acid identity. The encoded enzyme enables organisms from this genus and the later discovered *Ca. Argoarchaeum* (Chen *et al.*, 2019) to activate non-methane alkanes such as *n*-butane, propane and ethane. Hence these enzymes were called alkyl-coenzyme M reductases (ACR, Chen *et al.*, 2019). ACR sequences are mostly found in marine hydrocarbon seep environments, oil reservoirs or hydrothermal environments that are rich in multi-carbon

alkanes (Fig. 2; Wang *et al.*, 2019a). *Ca.* Syntrophoarchaeum express four highly divergent *acr* gene operons, yet it is not known which of the encoded enzymes is required to activate specific alkanes such as *n*-butane (Laso-Pérez *et al.*, 2016). A study on hydrothermal sediments in Guaymas Basin showed that these archaea were also active in natural environments (Wang *et al.*, 2019a). *Ca.* Argoarchaeum and its recently discovered related genus *Ca.* Ethanoperedens contain one *acr* gene operon that encode enzymes optimized for the activation of ethane (Fig. 2; Chen *et al.*, 2019; Hahn *et al.*, 2020). These archaea belong to the order Methanosarcinales and are closely related to ANME-2d. Physiological experiments confirmed that these archaea oxidize ethane coupled to sulfate reduction (Chen *et al.*, 2019; Hahn *et al.*, 2020). Biochemical and structural characterization of these enzymes is needed to resolve the functional details and specificity of this alkane-oxidizing ACR family.

Diversity of MCR/ACR-containing archaea with unknown function

Until recently, *mcr* genes were only found in members of the phylum Euryarchaeota with methane as single substrate or reaction product. However, Evans *et al.* (2015) described two metagenome-assembled genomes (MAGs) affiliating with the candidate phylum Bathyarchaeota that contained *mcr* genes. The catalytic (alpha) subunits of MCRs from Bathyarchaeota have low amino acid similarity (i.e., < 60%) to alpha subunits of MCRs of the known ANMEs and methanogens. To date, members of the phylum Bathyarchaeota have defied cultivation and the exact function of the Bathyarchaeota-type MCR is still unknown. The Bathyarchaeota-type MCR sequences were detected in many marine and terrestrial environments (Fig. 1). This MCR type appears in MAGs from a variety of different lineages including *Ca.* Hadesarchaeota (Hua *et al.*, 2019; Wang *et al.*, 2019b), the candidate phylum Helarchaeota from the Asgard superphylum (Seitz *et al.*, 2019), *Ca.* Methanoliparia (Borrel *et al.*, 2019; Laso-Pérez *et al.*, 2019) and Archaeoglobi in the Euryarchaeota phylum (Boyd *et al.*, 2019; Wang *et al.*, 2019b), suggesting horizontal gene transfers. In MAGs of *Ca.* Methanoliparia, the Bathyarchaeota-type *mcr* gene operon appears together with a canonical *mcr* gene operon. Based on the co-occurrence of these apparently functionally different MCR types, *Ca.* Methanoliparia may couple alkane degradation to the formation of methane (Borrel *et al.*, 2019; Laso-Pérez *et al.*, 2019).

A different type of *mcr* gene was discovered in MAGs of the candidate phylum Verstraetearchaeota in the TACK superphylum (Fig. 2; Vanwonterghem *et al.*, 2016;

Berghuis *et al.*, 2019). The Verstraetearchaeota-type *mcr* genes additionally appear in MAGs of the candidate phylum Korarchaeota (Borrel *et al.*, 2019; McKay *et al.*, 2019; Wang *et al.*, 2019b), *Ca.* Nezharchaeota (Hua *et al.*, 2019; Wang *et al.*, 2019b), Thaumarchaeota (Hua *et al.*, 2019) and Archaeoglobi (Colman *et al.*, 2019; Hua *et al.*, 2019; Wang *et al.*, 2019b). The Verstraetearchaeota-type MCR were mostly found in hot springs (Fig. 1). Interestingly, some archaeal lineages belonging to Archaeoglobi and *Ca.* Korarchaeota possess both MCR and dissimilatory sulfate reductase-encoding genes, suggesting that these organisms might be able to couple methane oxidation and sulfate reduction without involving syntrophic partners (Colman *et al.*, 2019; McKay *et al.*, 2019; Wang *et al.*, 2019b). Organisms with Verstraetearchaeota-type MCR have not yet been cultured; hence the function of this MCR family is unknown. The large diversity of archaeal MAGs with *mcr* gene operons but unknown functioning underlines the great demand for cultivation in this field of research.

Metabolism of anaerobic alkane-oxidizing archaea

The AOM process is catalysed by enzymes similar to those of the methanogenesis pathway. They allow the activation of methane and the complete oxidation of methyl groups to CO₂. For the anaerobic multi-carbon alkane oxidation, ANKA activate alkanes via ACR, analogous to the activity of MCR in ANME, yet they also combine a fatty acid degradation and the Wood-Ljungdahl pathway. Additionally, ANKA require a machinery for the transformation of alkyl-CoM to acyl-CoA, yet so far, enzymes catalysing these reactions have not been identified. Electrons released in anaerobic alkane oxidation are subsequently transferred via electron carriers to partner bacteria or directly to terminal electron acceptors.

Activation of methane and other alkanes via MCR/ACR enzymes

In all so far described anaerobic alkane-oxidizing archaea, the first step for alkane oxidation is the transformation of methane or multi-carbon alkanes to methyl- or alkyl-CoM via MCR or ACR respectively (Fig. 3A, Krüger *et al.*, 2003; Shima *et al.*, 2012; Laso-Pérez *et al.*, 2016; Chen *et al.*, 2019). The MCR sequences of methane oxidizers are highly similar to those of methanogens (Fig. 2), and their phylogeny suggests that methane-activating MCRs may have evolved from multiple methanogenic MCR lineages. The three-dimensional structure and functioning of methanogenic and methane-activating MCRs were resolved by protein crystallization and biochemical approaches (Ermler *et al.*, 1997; Shima *et al.*, 2012; Wongnate *et al.*, 2016; also reviewed in Thauer, 2019).




Fig 3 Pathways of alkanes oxidation and electron flows in anaerobic alkane-oxidizing archaea.

A. Schematic of the pathways for anaerobic methane, ethane and multi-carbon alkanes oxidation, including the activation of alkanes to alkyl-CoM (Part I); the transfer of methyl- or alkyl-CoM to methyl-H₄MPT or acyl-CoA, respectively (Part II); the beta-oxidation and Wood-Ljungdahl pathway for complete oxidation of methyl-H₄MPT or acyl-CoA, respectively (Part III); in C_nH_{2n+2}, n > 3. W-L: Wood-Ljungdahl pathway; β-ox: beta-oxidation pathway. Remarks: ¹The genomes of ANME-1 and *Ca. Syntrophoarchaeum* do not encode a methylenetetrahydromethanopterin (H₄MPT) reductase (Mer). Mer might be replaced by methylenetetrahydrofolate (H₄MF) reductase (MetF). ²Some of the published MAGs of *Ca. Argoarchaeum* do not code for a formylmethanofuran tetrahydromethanopterin formyltransferase (Ftr); it could be replaced by methylenetetrahydrofolate dehydrogenase (FolD). ³The even- and odd-carbon numbered alkane degradation may finally yield an acetyl-CoA or a propionyl-CoA, respectively, and the metabolism of propionyl-CoA in ANKA is still unknown.

These MCRs are highly similar, and their active centres are identical. Yet the MCR of ANME-1 contains cysteine-rich patches near the active centre and several post-translational amino acid modifications that are not found in the MCR of methanogens. Furthermore, in ANME-1, the cofactor F430 contains an additional methylthiol that is not observed in ANME-2 and ANME-3 or methanogens (Mayr *et al.*, 2008). Based on simulated three-dimensional structures inferred from amino acid sequences, ACRs largely resemble typical MCRs. Proposed modifications of ACR include a replacement of aromatic by aliphatic amino acids near or at the active pocket (Borrel *et al.*, 2019; Chen *et al.*, 2019), which might be needed to sterically accommodate multi-carbon substrates. However, only enzyme crystallization will allow resolving the true structure of these enzymes.

Transfer of methyl- or alkyl-CoM to methyl-H₄MPT or acyl-CoA respectively

ANMEs encode a tetrahydromethanopterin S-methyltransferase (MTR) to transfer the methyl group from CoM to tetrahydromethanopterin to allow further reactions that are required for complete oxidation (Fig. 3A, Hallam *et al.*, 2004; Timmers *et al.*, 2017). Ethane oxidizers *Ca. Argoarchaeum* and *Ca. Ethanoperedens* also encode for MTR (Chen *et al.*, 2019; Hahn *et al.*, 2020). Chen *et al.* (2019) proposed that the MTR of *Ca. Argoarchaeum* catalyses the ethyl transfer to a yet unknown intermediate (Fig. 3A, Chen *et al.*, 2019). However, MTR is not known to catalyse multi-carbon compound transformations. Alternatively, MTR may recycle small quantities of methyl-CoM that form as a by-product in ACR (Hahn *et al.*, 2020). Some of the required transformation steps of ethyl-CoM to acetyl-CoA might be catalysed by aldehyde ferredoxin oxidoreductases (Hahn *et al.*, 2020). The propane and *n*-butane-oxidizing archaea *Ca. Syntrophoarchaeum* do not have MTR, indicating that this enzyme is not essential for multi-carbon alkane metabolism in archaea. Laso-Pérez *et al.* (2016) and Wang *et al.* (2019a) suggested that the transfer between butyl-CoM to butyryl-CoA might be catalysed by homologues of methylcobamide:CoM methyltransferase that are found in all *Ca. Syntrophoarchaeum* genomes (Laso-Pérez *et al.*, 2016; Wang *et al.*, 2019a). Yet this reaction cannot explain the alkyl oxidation and transfer to CoA. Extensive metabolite analyses of the ANKA cultures are required to clarify the role of MTR in ethane oxidizers and to resolve how CoM-bound alkyl units are transferred to CoA-bound

fatty acids. Further insights into the role of the above-mentioned enzymes could be retrieved by heterologous gene expressions followed by *in vitro* enzyme experiments.

Beta-oxidation and Wood–Ljungdahl pathway for complete oxidation of alkanes

In ANMEs, methyl-H₄MPT can be subsequently oxidized to CO₂ by the C1 branch of Wood–Ljungdahl pathway. Ethane-oxidizing *Ca. Argoarchaeum* and *Ca. Ethanoperedens* also use the complete Wood–Ljungdahl pathway for acetyl-CoA degradation. In *Ca. Syntrophoarchaeum* and potential other long-chain alkane degraders, the formed acyl-CoA is first subjected to beta-oxidation, then the produced acetyl-CoA is oxidized via the Wood–Ljungdahl pathway (Fig. 3A, Laso-Pérez *et al.*, 2016, 2019; Wang *et al.*, 2019a). The beta-oxidation pathway is widespread in all domains of life, and it can also be used in the generation of fatty acid (Dibrova *et al.*, 2014). The pathway involves at least four enzymes: acyl-CoA dehydrogenase (FadE), crotonase (FadB1), 3-hydroxyacyl-CoA dehydrogenase (FadB2) and acetyl-CoA acetyltransferase (AtoB). *Ca. Syntrophoarchaeum* possess and express several of these genes (Laso-Pérez *et al.*, 2016; Wang *et al.*, 2019a). Interestingly, ANME-1 and ANME-2d also contain genes encoding beta-oxidation (Wang *et al.*, 2019a). However ANME show only low expression of these genes (Krukenberg *et al.*, 2018); hence, the encoded enzymes should not have catabolic functions. Instead they may be involved in the production or degradation of rare long-chain fatty acids (Jagersma *et al.*, 2012).


In ANMEs, the complete Wood–Ljungdahl pathway with the acetyl-CoA synthase:CO dehydrogenase complex has a central anabolic role as it produces acetyl-CoA for biomass production. Interestingly, isotope-labelling studies revealed that many ANME strains assimilate inorganic carbon (DIC) and only small amounts of methane when forming lipids and total biomass, and hence they should be considered as autotrophs (Kellermann *et al.*, 2012; Wegener *et al.*, 2016). This is puzzling because the Wood–Ljungdahl pathway is able to catalyse both DIC assimilation and methane oxidation. To regulate the extent of carbon fixation, both processes might be temporally separated. Alternatively, ANMEs may have a yet unidentified additional enzymatic route for carbon fixation or methyl group oxidation. For ANKA, *Ca. Argoarchaeum* and *Ca. Syntrophoarchaeum* produce acetyl-CoA as an

B. Electron transfer and coenzymes cycling with emphasis on CoB-S-S-CoM regeneration, electron carriers (coloured in red), electron transfer (coloured in blue) and proton or sodium ion translocation (coloured in yellow). Particularly, the oxidation of CoB-SH and CoM-SH deliver two electrons; the oxidation of FADH₂ and NADH deliver four electrons; the oxidation of F₄₂₀H₂ deliver two electrons. Fqo¹ (Fqo/FpoAHJKLMN), Fqo² (Fqo/FpoBCDFIO).

intermediate in the degradation of their substrate (Fig. 3A, Laso-Pérez *et al.*, 2016; Chen *et al.*, 2019). Hence, these organisms should be heterotrophs. To confirm this hypothesis, isotope-labelling experiments need to be performed.

Electron transfer and coenzyme cycling

The oxidation of methane and other alkanes according to the general catabolic reaction (1)



releases 8 (methane), 14 (ethane) or 26 (butane) electrons. These electrons are transferred to terminal electron acceptors via a number of electron-shuttling coenzymes generating transmembrane proton or sodium gradients for ATP synthesis. The first step of anaerobic alkane-oxidizing archaea is the oxidation of alkanes by MCR/ACR. The electron acceptor of this reaction is CoM-S-S-CoB, which is reduced to alkyl-S-CoM, now carrying the covalently bound alkane chain, and HS-CoB. The subsequent regeneration of CoM-S-S-CoB releases two electrons that are transferred to cytoplasmic electron carriers or directly to the membrane. For CoM-S-S-CoB regeneration, ANME-2a/b/c/d and *Ca. Argoarchaeum* have a cytoplasmic (HdrABC) and a membrane-bound (HdrDE) heterodisulfide reductase. HdrDE catalyses CoM-S-S-CoB regeneration but requires the inflow of two protons (Wang *et al.*, 2014; Arshad *et al.*, 2015; Krukenberg *et al.*, 2018; Chen *et al.*, 2019). In contrast, ANME-1 and *Ca. Syntrophoarchaeum* possess only the cytoplasmic HdrABC (Stokke *et al.*, 2012; Laso-Pérez *et al.*, 2016; Wang *et al.*, 2019a). Unlike HdrDE, HdrABC alone cannot catalyse the regeneration of CoM-S-S-CoB (Wagner *et al.*, 2017). Arshad *et al.* (2015) suggested that an enzyme complex consisting of HdrABC and a coenzyme F_{420} hydrogenase (FrhB) couples CoM-S-S-CoB regeneration and ferredoxin oxidation to the reduction of two molecules of F_{420} (Fig. 3B, Arshad *et al.*, 2015).

In ANME, the transfer of the methyl group from CoM to H_4MPT is catalysed by MTR with the inflow of two sodium ions. The further oxidation of CH_3-H_4MPT via the Wood-Ljungdahl pathway involves the reduction of $F_{420}/F_{420}H_2$. In total, four electrons are transferred from CH_3-H_4MPT to two molecules of F_{420} by forming two $F_{420}H_2$ and $CH-H_4MPT$ in two reactions. ANME-2a/b/c/d and *Ca. Argoarchaeum* can reoxidize $F_{420}H_2$ with membrane-bound $F_{420}H_2$:phenazine oxidoreductase (Fpo, Wang *et al.*, 2014; Arshad *et al.*, 2015; Krukenberg *et al.*, 2018; Chen *et al.*, 2019). In contrast, ANME-1 and *Ca. Syntrophoarchaeum* use $F_{420}H_2$:quinone oxidoreductases (Fqo) for this function (Fig. 3B, Meyerdierks *et al.*, 2010; Krukenberg *et al.*, 2018; Laso-Pérez *et*

al., 2016; Wang *et al.*, 2019a). Through methanophenazine (MP) or menaquinone (MQ), electrons are transferred to cytochromes and external sinks, a process that is coupled to proton translocations (Fig. 3B). During the last step of the Wood-Ljungdahl pathway, two electrons from formyl group oxidation are transferred to ferredoxin, which can be replenished by the previously described reaction catalysed by HdrABC in all ANMEs. ANME-2a/b and ANME-2c additionally encode a membrane-integral, sodium-motive ferredoxin:NAD⁺ oxidoreductase complex (Rnf). The reaction catalysed by Rnf involves the translocation of two sodium ions (Fig. 3B, Wang *et al.*, 2014; Krukenberg *et al.*, 2018; Wang *et al.*, 2019a).

In ANKA, the transformation of alkyl-CoM to acyl-CoA should release four electrons to yet unknown electron carriers. Anaerobic ethane oxidizers, similar to ANMEs, use the Wood-Ljungdahl pathway to degrade acetyl-CoA, which releases another eight electrons if completely oxidized to CO_2 (Fig. 3A). *Ca. Syntrophoarchaeum* uses the beta-oxidation pathway for acyl-CoA degradation. For instance, the degradation of butyryl-CoA into two units of acetyl-CoA, releases eight electrons that are transferred to two units of FAD and NAD⁺ (Fig. 3B). These electron carriers are reoxidized by enzymes of the respiratory chain translocating around six protons per acetyl-CoA released (Villanueva *et al.*, 2017).

Most ANME and ANKA transfer the electrons from alkane oxidation to partner bacteria, while specific strains of ANME-2d deliver electrons to nitrate or metal oxides. MAGs of ANME and ANKA contain many genes encoding cytochromes and archaeum-like proteins. These genes are highly expressed, and microscopic analyses revealed high abundance of extracellular cytochromes partly coupled to the S-layer of archaea (McGlynn *et al.*, 2015; Wegener *et al.*, 2015; Laso-Pérez *et al.*, 2016; Krukenberg *et al.*, 2018). Moreover, the partner bacteria produce cytochromes and pili-based nanowires. Together these cell structures may allow direct interspecies electron transfer from the archaea to their partner bacteria. Heterologous expression experiments in model organisms as shown for pili-based nanowires (Walker *et al.*, 2018) will help to assess unresolved pathways in anaerobic alkane oxidation and the coupling to electron acceptors.

In AOM, the combined membrane-coupled reactions yield a surplus of about two protons in the periplasm per molecule methane oxidized depending on the different types of ANME. Based on the required inflow of five protons per ATP (Thauer, 2011), or the free energy released from the methane oxidation ($\Delta G^{\prime m} = -35 \text{ kJ mol}^{-1}$, McGlynn, 2017), this translates to a yield of only 0.3–0.4 mol ATP per molecule methane oxidized. In ANKA some catabolic reaction steps and their energy

requirements are unknown; hence, their proton translocations cannot be budgeted. However, thermodynamic considerations for the net reaction of multi-carbon alkane oxidation suggest more proton translocations per mole substrate oxidized, which would explain the higher growth yields of these organisms compared with ANME (Hahn *et al.*, 2020).

Conclusion and perspectives

The physiology and ecology of ANME have been studied intensively for 20 years. In contrast, the first ANKA have been identified about 4 years ago. Hence, to date, only three genera of non-methane alkane oxidizers have been cultured, and all belong to the class Methanomicrobia in the Euryarchaeota phylum. Based on metagenomic studies, an enormous diversity of ACR-containing archaea awaits their cultivation. A larger collection of cultures will help to understand the physiology, biochemistry, evolution and ecological roles of the fascinating alkane-metabolizing archaea. Structural and functional features of MCR/ACR enzymes that determine the preference for methane or multi-carbon alkane substrates needs to be resolved, ideally by crystallization of their native forms. Required biochemical characterizations include substrate spectra and kinetic analyses. Also, the transformation of CoM-bound alkyl units to CoA-bound fatty acids is not understood. As these reactions lack analogues in other organisms, extended metabolomics analyses are required. To resolve the role of alkane-metabolizing archaea in the global carbon cycle, metagenomic information needs to be combined with environmental information including alkane, sulfate/sulfide, nitrate/nitrite and metal ions fluxes. This environmental research should include understudied environments including the terrestrial subsurface, deep-sea ecosystems and hydrocarbon reservoirs.

Acknowledgements

We acknowledge funding by the State Key R&D Project of China (grant number 2018YFC0310800, 2016YFA0601102), COMRA Project DY135-B2-12, the National Nature Science Foundation of China (grant numbers 41525011, 41902313, 91751205), the State Key Laboratory of Ocean Engineering Foundation (grant number GKZD010075), and the DFG Cluster of Excellence 2077 'The Ocean Floor—Earth's Uncharted Interface' at MARUM, University of Bremen. This is a contribution to the International Center for Deep-Life Investigation (IC-DLI).

References

Adam, P.S., Borrel, G., Brochier-Armanet, C., and Gribaldo, S. (2017) The growing tree of archaea: new

- perspectives on their diversity, evolution and ecology. *ISME J* **11**: 2407–2425.
- Aeckersberg, F., Bak, F., and Widdel, F. (1991) Anaerobic oxidation of saturated hydrocarbons to CO₂ by a new type of sulfate-reducing bacterium. *Arch Microbiol* **156**: 5–14.
- Arshad, A., Speth, D.R., de Graaf, R.M., Op den Camp, H.J., Jetten, M.S., and Welte, C.U. (2015) A metagenomics-based metabolic model of nitrate-dependent anaerobic oxidation of methane by Methanoperedens-like archaea. *Front Microbiol* **6**: 1423.
- Berghuis, B.A., Yu, F.B., Schulz, F., Blainey, P.C., Woyke, T., and Quake, S.R. (2019) Hydrogenotrophic methanogenesis in archaeal phylum Verstraetearchaeota reveals the shared ancestry of all methanogens. *Proc Natl Acad Sci U S A* **116**: 5037–5044.
- Bertram, S., Blumenberg, M., Michaelis, W., Siebert, M., Krüger, M., and Seifert, R. (2013) Methanogenic capabilities of ANME-archaea deduced from ¹³C-labelling approaches. *Environ Microbiol* **15**: 2384–2393.
- Beulig, F., Røy, H., McGlynn, S.E., and Jørgensen, B.B. (2019) Cryptic CH₄ cycling in the sulfate–methane transition of marine sediments apparently mediated by ANME-1 archaea. *ISME J* **13**: 250–262.
- Bhattacharai, S., Cassarini, C., and Lens, P.N.L. (2019) Physiology and distribution of archaeal methanotrophs that couple anaerobic oxidation of methane with sulfate reduction. *Microbiol Mol Biol Rev* **83**: e00074-18.
- Boetius, A., and Wenzhöfer, F. (2013) Seafloor oxygen consumption fueled by methane from cold seeps. *Nat Geosci* **6**: 725–734.
- Borrel, G., Adam, P.S., McKay, L.J., Chen, L.X., Sierra-García, I.N., Sieber, C.M., *et al.* (2019) Wide diversity of methane and short-chain alkane metabolisms in uncultured archaea. *Nat Microbiol* **4**: 603–613.
- Boyd, J.A., Jungbluth, S.P., Leu, A.O., Evans, P.N., Woodcroft, B.J., & Chadwick, G.L. (2019). Divergent methyl-coenzyme M reductase genes in a deep-subseafloor Archaeoglobi. *The ISME Journal*, **1**,
- Cai, C., Leu, A.O., Xie, G.J., Guo, J., Feng, Y., Zhao, J.X., *et al.* (2018) A methanotrophic archaeon couples anaerobic oxidation of methane to Fe (III) reduction. *ISME J* **12**: 1929–1939.
- Cassarini, C., Zhang, Y., and Lens, P.N. (2019) Pressure selects dominant anaerobic methanotrophic phylotype and sulfate reducing bacteria in coastal marine Lake Grevelingen sediment. *Front Environ Sci* **6**: 162.
- Chen, S.C., Musat, N., Lechtenfeld, O.J., Paschke, H., Schmidt, M., Said, N., *et al.* (2019) Anaerobic oxidation of ethane by archaea from a marine hydrocarbon seep. *Nature* **568**: 108–111.
- Coates, J.D., Woodward, J., Allen, J., Philp, P., and Lovley, D.R. (1997) Anaerobic degradation of polycyclic aromatic hydrocarbons and alkanes in petroleum-contaminated marine harbor sediments. *Appl Environ Microbiol* **63**: 3589–3593.
- Colman, D.R., Lindsay, M.R., and Boyd, E.S. (2019) Mixing of meteoric and geothermal fluids supports hyperdiverse chemosynthetic hydrothermal communities. *Nat Commun* **10**: 681.
- Conrad, R. (2009) The global methane cycle: recent advances in understanding the microbial processes involved. *Environ Microbiol Rep* **1**: 285–292.

- Cui, M., Ma, A., Qi, H., Zhuang, X., and Zhuang, G. (2015) Anaerobic oxidation of methane: an "active" microbial process. *Microbiologyopen* **4**: 1–11.
- Dedysch, S.N., and Knief, C. (2018) Diversity and phylogeny of described aerobic methanotrophs. In *Methane Biocatalysis: Paving the Way to Sustainability*. Cham, Switzerland: Springer, pp. 17–42.
- Dibrova, D.V., Galperin, M.Y., and Mulkidjanian, A.Y. (2014) Phylogenomic reconstruction of archaeal fatty acid metabolism. *Environ Microbiol* **16**: 907–918.
- Ermiler, U., Grabarse, W., Shima, S., Goubeaud, M., and Thauer, R.K. (1997) Crystal structure of methyl-coenzyme M reductase: the key enzyme of biological methane formation. *Science* **278**: 1457–1462.
- Evans, P.N., Parks, D.H., Chadwick, G.L., Robbins, S.J., Orphan, V.J., Golding, S.D., and Tyson, G.W. (2015) Methane metabolism in the archaeal phylum Bathyarchaeota revealed by genome-centric metagenomics. *Science* **350**: 434–438.
- Evans, P.N., Boyd, J.A., Leu, A.O., Woodcroft, B.J., Parks, D.H., Hugenholz, P., and Tyson, G.W. (2019) An evolving view of methane metabolism in the archaea. *Nat Rev Microbiol* **17**: 219–232.
- Felden, J., Ruff, S.E., Ertefai, T., Inagaki, F., Hinrichs, K.U., and Wenzhöfer, F. (2014) Anaerobic methanotrophic community of a 5346-m-deep vesicomyid clam colony in the Japan trench. *Geobiology* **12**: 183–199.
- Friedrich, M.W. (2005) Methyl-coenzyme M reductase genes: unique functional markers for methanogenic and anaerobic methane-oxidizing Archaea. *Methods Enzymol* **397**: 428–442.
- Hahn, C.J., Laso-Pérez, R., Vulcano, F., Vaziourakis, K.M., Stokke, R., Steen, I.H., et al. (2020) "Candidatus Ethanoperedens," a thermophilic genus of archaea mediating the anaerobic oxidation of ethane. *mBio* **28**: 11.
- Hallam, S.J., Putnam, N., Preston, C.M., Detter, J.C., Rokhsar, D., & Richardson, P.M. (2004). Reverse methanogenesis: testing the hypothesis with environmental genomics. *Science*, **305**(5689), 1457–1462.
- Haroon, M.F., Hu, S., Shi, Y., Imelfort, M., Keller, J., Hugenholz, P., et al. (2013) Anaerobic oxidation of methane coupled to nitrate reduction in a novel archaeal lineage. *Nature* **500**: 567–570.
- Hinrichs, K.U., Hayes, J.M., Sylva, S.P., Brewer, P.G., and DeLong, E.F. (1999) Methane-consuming archaeobacteria in marine sediments. *Nature* **398**: 802–805.
- Holler, T., Widdel, F., Knittel, K., Amann, R., Kellermann, M. Y., Hinrichs, K.U., et al. (2011) Thermophilic anaerobic oxidation of methane by marine microbial consortia. *ISME J* **5**: 1946–1956.
- Hua, Z.S., Wang, Y.L., Evans, P.N., Qu, Y.N., Goh, K.M., Rao, Y.Z., et al. (2019) Insights into the ecological roles and evolution of methyl-coenzyme M reductase-containing hot spring archaea. *Nat Commun* **10**: 1–11.
- Jaekel, U., Musat, N., Adam, B., Kuypers, M., Grundmann, O., and Musat, F. (2013) Anaerobic degradation of propane and butane by sulfate-reducing bacteria enriched from marine hydrocarbon cold seeps. *ISME J* **7**: 885–895.
- Jagersma, C.G., Meulepas, R.J., Timmers, P.H., Szperl, A., Lens, P.N., and Stams, A.J. (2012) Enrichment of ANME-1 from Eckernförde Bay sediment on thiosulfate, methane and short-chain fatty acids. *J Biotechnol* **157**: 482–489.
- Kallistova, A.Y., Merkel, A.Y., Tamovetskii, I.Y., and Pimenov, N.V. (2017) Methane formation and oxidation by prokaryotes. *Microbiology* **86**: 671–691.
- Kellermann, M.Y., Wegener, G., Elvert, M., Yoshinaga, M.Y., Lin, Y.S., Holler, T., et al. (2012) Autotrophy as a predominant mode of carbon fixation in anaerobic methane-oxidizing microbial communities. *Proc Natl Acad Sci USA* **109**: 19321–19326.
- Kleindienst, S., Ramette, A., Amann, R., and Knittel, K. (2012) Distribution and in situ abundance of sulfate-reducing bacteria in diverse marine hydrocarbon seep sediments. *Environ Microbiol* **14**: 2689–2710.
- Kniemeyer, O., Musat, F., Sievert, S.M., Knittel, K., Wilkes, H., Blumenberg, M., et al. (2007) Anaerobic oxidation of short-chain hydrocarbons by marine sulphate-reducing bacteria. *Nature* **449**: 898–901.
- Knittel, K., and Boetius, A. (2009) Anaerobic oxidation of methane: progress with an unknown process. *Annu Rev Microbiol* **63**: 311–334.
- Knittel, K., Lösekann, T., Boetius, A., Kort, R., and Amann, R. (2005) Diversity and distribution of methanotrophic archaea at cold seeps. *Appl Environ Microbiol* **71**: 467–479.
- Krüger, M., Meyerdierks, A., Glöckner, F.O., Amann, R., Widdel, F., Kube, M., et al. (2003) A conspicuous nickel protein in microbial mats that oxidize methane anaerobically. *Nature* **426**: 878–881.
- Krukenberg, V., Harding, K., Richter, M., Glöckner, F.O., Berg, J., & Knittel, K. (2016). Candidatus Desulfoterrivudis auxilii, a hydrogenotrophic sulfate-reducing bacterium of the HotSeep-1 cluster involved in the thermophilic anaerobic oxidation of methane. *Environ Microbiol*, **18**(9), 3073–3091.
- Krukenberg, V., Riedel, D., Gruber-Vodicka, H.R., Buttigieg, P.L., Tegetmeyer, H.E., Boetius, A., and Wegener, G. (2018) Gene expression and ultrastructure of meso- and thermophilic methanotrophic consortia. *Environ Microbiol* **20**: 1651–1666.
- Laso-Pérez, R., Wegener, G., Knittel, K., Widdel, F., Harding, K.J., Krukenberg, V., et al. (2016) Thermophilic archaea activate butane via alkyl-coenzyme M formation. *Nature* **539**: 396–401.
- Laso-Pérez, R., Hahn, C., van Vliet, D.M., Tegetmeyer, H.E., Schubotz, F., Smit, N.T., et al. (2019) Anaerobic degradation of non-methane alkanes by "Candidatus Methanoliparia" in hydrocarbon seeps of the Gulf of Mexico. *mBio* **10**: e01814-19.
- Lee, D.H., Lee, Y.M., Kim, J.H., Jin, Y.K., Paull, C., Niemann, H., et al. (2019) Discriminative biogeochemical signatures of methanotrophs in different chemosynthetic habitats at an active mud volcano in the Canadian Beaufort Sea. *Sci Rep* **9**: 1–13.
- Leu, A.O., Cai, C., Mcllroy, S.J., Southam, G., Orphan, V.J., Yuan, Z., et al. (2020) Anaerobic methane oxidation coupled to manganese reduction by members of the Methanoperedenaceae. *ISME J* **14**: 1030–1041.
- Liang, L., Wang, Y., Sivan, O., and Wang, F. (2019) Metal-dependent anaerobic methane oxidation in marine

- sediment: insights from marine settings and other systems. *Sci China Life Sci* **62**: 1287–1295.
- Lloyd, K.G., Alperin, M.J., and Teske, A. (2011) Environmental evidence for net methane production and oxidation in putative ANaerobic MEthanotrophic (ANME) archaea. *Environ Microbiol* **13**: 2548–2564.
- Mayr, S., Latkoczy, C., Kruger, M., Gunther, D., Shima, S., Thauer, R.K., *et al.* (2008) Structure of an F430 variant from archaea associated with anaerobic oxidation of methane. *J Am Chem Soc* **130**: 10758–10767.
- McGlynn, S.E. (2017) Energy metabolism during anaerobic methane oxidation in ANME archaea. *Microbes Environ* **32**: 5–13.
- McGlynn, S.E., Chadwick, G.L., Kempes, C.P., and Orphan, V.J. (2015) Single cell activity reveals direct electron transfer in methanotrophic consortia. *Nature* **526**: 531–535.
- McKay, L.J., Đlakić, M., Fields, M.W., Delmont, T.O., Eren, A.M., Jay, Z.J., *et al.* (2019) Co-occurring genomic capacity for anaerobic methane and dissimilatory sulfur metabolisms discovered in the Korarchaeota. *Nat Microbiol* **4**: 614–622.
- Meyerdierks, A., Kube, M., Kostadinov, I., Teeling, H., Glöckner, F.O., Reinhardt, R., and Amann, R. (2010) Metagenome and mRNA expression analyses of anaerobic methanotrophic archaea of the ANME-1 group. *Environ Microbiol* **12**: 422–439.
- Michaelis, W., Seifert, R., Nauhaus, K., Treude, T., Thiel, V., Blumenberg, M., *et al.* (2002) Microbial reefs in the Black Sea fueled by anaerobic oxidation of methane. *Science* **297**: 1013–1015.
- Mills, H.J., Hodges, C., Wilson, K., MacDonald, I.R., and Sobocky, P.A. (2003) Microbial diversity in sediments associated with surface-breaching gas hydrate mounds in the Gulf of Mexico. *FEMS Microbiol Ecol* **46**: 39–52.
- Nauhaus, K., Albrecht, M., Elvert, M., Boetius, A., and Widdel, F. (2007) In vitro cell growth of marine archaeal-bacterial consortia during anaerobic oxidation of methane with sulfate. *Environ Microbiol* **9**: 187–196.
- Niemann, H., Lösekann, T., de Beer, D., Elvert, M., Nadalig, T., Knittel, K., *et al.* (2006) Novel microbial communities of the Haakon Mosby mud volcano and their role as a methane sink. *Nature* **443**: 854–858.
- Niu, M., Fan, X., Zhuang, G., Liang, Q., and Wang, F. (2017) Methane-metabolizing microbial communities in sediments of the Haima cold seep area, northwest slope of the South China Sea. *FEMS Microbiol Ecol* **93**: fix101.
- Omeregge, E.O., Mastalerz, V., de Lange, G., Straub, K.L., Kappler, A., Røy, H., *et al.* (2008) Biogeochemistry and community composition of iron and sulfur-precipitating microbial mats at the Chefred mud volcano (Nile Deep Sea Fan, eastern Mediterranean). *Appl Environ Microbiol* **74**: 3198–3215.
- Orphan, V.J., House, C.H., Hinrichs, K.U., McKeegan, K.D., and DeLong, E.F. (2002) Multiple archaeal groups mediate methane oxidation in anoxic cold seep sediments. *Proc Natl Acad Sci USA* **99**: 7663–7668.
- Reeburgh, W.S. (2007) Oceanic methane biogeochemistry. *Chem Rev* **107**: 486–513.
- Ruff, S.E., Biddle, J.F., Teske, A.P., Knittel, K., Boetius, A., and Ramette, A. (2015) Global dispersion and local diversification of the methane seep microbiome. *Proc Natl Acad Sci USA* **112**: 4015–4020.
- Ruff, S.E., Kuhfuss, H., Wegener, G., Lott, C., Ramette, A., Wiedling, J., *et al.* (2016) Methane seep in shallow-water permeable sediment harbors high diversity of anaerobic methanotrophic communities, Elba, Italy. *Front Microbiol* **7**: 374.
- Ruff, S.E., Felden, J., Gruber-Vodicka, H.R., Marcon, Y., Knittel, K., Ramette, A., and Boetius, A. (2019) In situ development of a methanotrophic microbiome in deep-sea sediments. *ISME J* **13**: 197–213.
- Schreiber, L., Holler, T., Knittel, K., Meyerdierks, A., and Amann, R. (2010) Identification of the dominant sulfate-reducing bacterial partner of anaerobic methanotrophs of the ANME-2 clade. *Environ Microbiol* **12**: 2327–2340.
- Seitz, K.W., Dombrowski, N., Eme, L., Spang, A., Lombard, J., Sieber, J.R., *et al.* (2019) Asgard archaea capable of anaerobic hydrocarbon cycling. *Nat Commun* **10**: 1822.
- Shima, S., Krueger, M., Weinert, T., Demmer, U., Kahnt, J., Thauer, R.K., and Ermler, U. (2012) Structure of a methyl-coenzyme M reductase from Black Sea mats that oxidize methane anaerobically. *Nature* **481**: 98–101.
- Stokke, R., Roalkvam, I., Lanzen, A., Hafliðason, H., and Steen, I.H. (2012) Integrated metagenomic and metaproteomic analyses of an ANME-1-dominated community in marine cold seep sediments. *Environ Microbiol* **14**: 1333–1346.
- Stolper, D.A., Lawson, M., Davis, C.L., Ferreira, A.A., Neto, E.S., Ellis, G.S., *et al.* (2014) Formation temperatures of thermogenic and biogenic methane. *Science* **344**: 1500–1503.
- Teske, A., Hinrichs, K.U., Edgcomb, V., de Vera Gomez, A., Kyselá, D., Sylva, S.P., *et al.* (2002) Microbial diversity of hydrothermal sediments in the Guaymas Basin: evidence for anaerobic methanotrophic communities. *Appl Environ Microbiol* **68**: 1994–2007.
- Thauer, R.K. (2011) Anaerobic oxidation of methane with sulfate: on the reversibility of the reactions that are catalyzed by enzymes also involved in methanogenesis from CO₂. *Curr Opin Microbiol* **14**: 292–299.
- Thauer, R.K. (2019) Methyl (alkyl)-coenzyme M reductases: Nickel F-430-containing enzymes involved in anaerobic methane formation and in anaerobic oxidation of methane or of short chain alkanes. *Biochemistry* **58**: 5198–5220.
- Timmers, P.H., Welte, C.U., Koehorst, J.J., Plugge, C.M., Jetten, M.S., and Stams, A.J. (2017) Reverse methanogenesis and respiration in methanotrophic archaea. *Archaea* **2017**: 1–22.
- Vanwonterghem, I., Evans, P.N., Parks, D.H., Jensen, P.D., Woodcroft, B.J., Hugenholtz, P., and Tyson, G.W. (2016) Methylothrophic methanogenesis discovered in the archaeal phylum Verstraetearchaeota. *Nat Microbiol* **1**: 16170.
- Villanueva, L., Schouten, S., and Damsté, J.S.S. (2017) Phylogenomic analysis of lipid biosynthetic genes of archaea shed light on the ‘lipid divide’. *Environ Microbiol* **19**: 54–69.
- Wagner, T., Koch, J., Ermler, U., and Shima, S. (2017) Methanogenic heterodisulfide reductase (HdrABC-

- MvhAGD) uses two noncubane [4Fe-4S] clusters for reduction. *Science* **357**: 699–703.
- Wahlen, M. (1993) The global methane cycle. *Annu Rev Earth Planet Sci* **21**: 407–426.
- Walker, D.J., Adhikari, R.Y., Holmes, D.E., Ward, J.E., Woodard, T.L., Nevin, K.P., and Lovley, D.R. (2018) Electrically conductive pili from pilin genes of phylogenetically diverse microorganisms. *ISME J* **12**: 48–58.
- Wang, F.P., Zhang, Y., Chen, Y., He, Y., Qi, J., Hinrichs, K. U., *et al.* (2014) Methanotrophic archaea possessing diverging methane-oxidizing and electron-transporting pathways. *ISME J* **8**: 1069–1078.
- Wang, V.C.C., Maji, S., Chen, P.P.Y., Lee, H.K., Yu, S.S.F., and Chan, S.I. (2017) Alkane oxidation: methane monooxygenases, related enzymes, and their biomimetics. *Chem Rev* **117**: 8574–8621.
- Wang, Y., Feng, X., Natarajan, V.P., Xiao, X., and Wang, F. (2019a) Diverse anaerobic methane- and multi-carbon alkane-metabolizing archaea coexist and show activity in Guaymas Basin hydrothermal sediment. *Environ Microbiol* **21**: 1344–1355.
- Wang, Y., Wegener, G., Hou, J., Wang, F., and Xiao, X. (2019b) Expanding anaerobic alkane metabolism in the domain of archaea. *Nat Microbiol* **4**: 595–602.
- Wegener, G., Niemann, H., Elvert, M., Hinrichs, K.U., and Boetius, A. (2008) Assimilation of methane and inorganic carbon by microbial communities mediating the anaerobic oxidation of methane. *Environ Microbiol* **10**: 2287–2298.
- Wegener, G., Krukenberg, V., Riedel, D., Tegetmeyer, H.E., and Boetius, A. (2015) Intercellular wiring enables electron transfer between methanotrophic archaea and bacteria. *Nature* **526**: 587–590.
- Wegener, G., Krukenberg, V., Ruff, S.E., Kellermann, M.Y., and Knittel, K. (2016) Metabolic capabilities of microorganisms involved in and associated with the anaerobic oxidation of methane. *Front Microbiol* **7**: 46.
- Welte, C.U., Rasigraf, O., Vaksmaa, A., Versantvoort, W., Arshad, A., Op den Camp, H.J., *et al.* (2016) Nitrate- and nitrite-dependent anaerobic oxidation of methane. *Environ Microbiol Rep* **8**: 941–955.
- Wongnate, T., Sliwa, D., Ginovska, B., Smith, D., Wolf, M. W., Lehnert, N., *et al.* (2016) The radical mechanism of biological methane synthesis by methyl-coenzyme M reductase. *Science* **352**: 953–958.
- Yvon-Durocher, G., Allen, A.P., Bastviken, D., Conrad, R., Gudasz, C., St-Pierre, A., *et al.* (2014) Methane fluxes show consistent temperature dependence across microbial to ecosystem scales. *Nature* **507**: 488–491.
- Zhang, Y., Maignien, L., Zhao, X., Wang, F., and Boon, N. (2011) Enrichment of a microbial community performing anaerobic oxidation of methane in a continuous high-pressure bioreactor. *BMC Microbiol* **11**: 137.