

Evaluation of paralytic shellfish poisoning toxin profile of mussels from Bulgarian North Black Sea coast by HPLC-FID with post and pre-column derivatization

Z. V. Peteva^{1*}, G. N. Kalinova², B. Krock³, M. D. Stancheva⁴, St. K. Georgieva⁵

¹Medical University - Varna, Department of Chemistry, Tsar Osvoboditel, 84, 9000 Varna, Bulgaria,

²National Diagnostic and Research Veterinary Medicine Institute, National reference Laboratory for Marine Biotoxins, Bulgaria, Pencho Slaveikov Blvd., 15, 1606 Sofia,

³Alfred Wegener Institut, Helmholtz Zentrum für Polar- und Meeresforschung, Chemische Ökologie, am Handelshafen 12, 27570 Bremerhaven, Germany

⁴Medical University - Varna, Department of Chemistry, Tsar Osvoboditel, 84, 9000 Varna, Bulgaria,

⁵Medical University - Varna, Department of Chemistry, Tsar Osvoboditel, 84, 9000 Varna, Bulgaria,

Received October 27, 2016; Revised December 2, 2016

Marine toxins are produced by certain toxic phytoplankton species. Harmful toxins may accumulate in the shellfish tissue, potentially impacting human health. Paralytic shellfish poisoning (PSP) is a syndrome caused by ingestion of shellfish contaminated with paralytic shellfish toxins (PST) that comprise saxitoxin and its variants (neosaxitoxin, gonyautoxins and their decarbamoyl and *N*-sulfofocarbamoyl analogs).

The aim of this study was to evaluate the presence of paralytic shellfish toxins (PSTs) in plankton samples and in mussels intended for human consumption.

Mussels collected in the main areas of production and recreational harvesting off the north coast of Bulgaria have been investigated for PSP toxins. Individual toxins were determined using two methods both involving fluorescence detection: ion pair-liquid chromatography with post-column derivatization (method 1) and high-performance liquid chromatographic procedure employing pre-column oxidation of the toxins (method 2). The results according method 1 demonstrated the presence of gonyautoxin 2 in 53% of the mussel samples and no toxins were detected in the plankton samples. The toxicity level - 1.6 µg STX.2HCl .kg⁻¹ was far beneath the EU legislative limit of 800 µg STX.2HCl .kg⁻¹ concluding in negligible risk for human health.

Due to higher limits of detection no toxins were detected via method 2. Even though, considering method 2 is recognized by European Commission as official for regulatory purposes and the relative high value of the legislative threshold, thus obtained toxin levels are enough representative to conclude if mussels are safe for consumption or not. On the other hand, the more sensitive method 1 provides important data on extremely low toxin levels which would be useful for chronic exposure estimation and for completing the knowledge about occurrence of PSTs in certain locations.

Keywords: paralytic shellfish toxins, gonyautoxin 2, Bulgaria, mussels, plankton, LC-FLD

INTRODUCTION

Seafood products such as shellfish are a rich source of proteins and mineral components [1, 2]. Mussels and rapana are an important source of income for the fishing community in Bulgaria [3]. However, shellfish such as mussels may contain compounds harmful to human health such as marine biotoxins.

Among them are paralytic shellfish poisoning (PSP) toxins (PSTs) that comprise saxitoxin (STX) and its variants. A dose of approximately 1 mg of saxitoxin from a single serving of contaminated shellfish is fatal to humans. PSTs are neurotoxins that specifically bind to voltage gated sodium channels and thereby block the excitation current in nerve and muscle cells, finally resulting in paralysis and other disorders in consumers of contaminated

shellfish [4]. Intoxication with PSTs produces several symptoms that include tinkling, sensation or numbness around the lips, numbness of extremities, gastrointestinal problems, difficulty in breathing, and, in fatal cases, death [5]. The human PSP fatality cases vary worldwide and depend essentially on local health - care practitioner awareness of the intoxication and its treatment and the local medical system's capacity on assisting respiratory support needing victims. For example, in a series of intoxications in Europe and North America, no deaths occurred among more than 200 cases. In similar outbreaks in Southeast Asia and Latin America, where clinicians may be unexperienced with PSP, the case fatality proportion ranged between 2-14% [4]. Worldwide, PSP affects some 2000 people annually, with a 15% mortality rate [6].

* To whom all correspondence should be sent.
E-mail: zlatina.peteva@mu-varna.bg

Wild mussels (7 samples) were harvested manually by recreational harvesters and by our co-workers. Farmed mussels (10 samples) were provided from mussel farmers directly from cultivation ropes.

Sample preparation:

The digestive glands of at least 1.5 kg specimen (~ 37.3 g) were dissected and homogenized with dispersing instrument (POLYMIX®PT 1200E, KINEMATIKA AG, Germany) for 5-10 min at maximum speed.

Extraction of Plankton:

The sample concentrate from the plankton net was collected on 20 µm mesh size sieve and washed in 50 mL centrifuge tube. The suspension was centrifuged (4000 × g, 10 min at 10 °C), the supernatant was removed and 1 mL 0.03 M acetic acid was added to the plankton pellet. For the extraction of PSTs from the plankton cells, the resulted suspension was sonicated (40 Hz, 15 min) and centrifuged again. The supernatant was filtered through syringe filters (0.45 µm pore size, ø 25 mm, Minisart, Sartorius, Germany). Filtrates were transferred into chromatographic vials and kept at -20 °C until further analysis.

Extraction of digestive glands:

- for PCOX analysis:

An average 2.14 g of hepatopancreas homogenate was extracted in duplicate with 4 mL 0.2 M acetic acid with dispersing instrument for 2 min at maximum speed and centrifuged for 10-15 min at 4000 × g. The resulting supernatants were transferred to 50 mL centrifuge tubes and centrifuged again. An aliquot (1 mL) was filtered through syringe filters. Filtrates were kept in chromatographic vials at -20 °C until further analysis.

- for Lawrence method [16]:

Hepatopancreas homogenate (~ 5g) was extracted in duplicate with 3 mL 1% acetic acid. First extraction was performed at boiling water for 5 min. Subsequently a centrifugation at 4000 × g for 10 min was performed. Supernatants were combined and made up to volume of 10 mL.

Clean-up of 1 mL extract was performed with SPE C18 cartridge (SUPELCO, 500mg/3 ml) and the extract was adjusted to pH 6.5 with 1M NaOH.

Pre-column derivatization

The sample extracts were oxidized with periodate oxidant - 100 µL of sample extract were added to 100 µL of matrix modifier (PSP-free oysters were extracted and cleaned-up by the same procedure as samples) in a 1.5 mL vial. 500 µL of periodate

oxidant (prepared daily by mixing 5 mL each of 0.03 M periodic acid, 0.3 M ammonium formate, and 0.3 M Na₂HPO₄, and adjusted to pH to 8.2 with 0.2 M NaOH using pH meter) were added to the vial and mixed. The solution reacted at room temperature for 1 min. Glacial acetic acid (5 µL) was added to interrupt the reaction.

HPLC- FD analysis

PCOX method (Method 1)

All mussel and plankton extracts were analyzed for PSTs by reverse-phase ion-pair liquid chromatography on a LC1100 series liquid chromatograph (Agilent Technologies, Waldbronn, Germany) coupled to a PCX 2500 post-column derivatization system (Pickering Laboratories, Mountain View, CA, USA) and dual monochromator fluorescence detector (G1321A) following minor modifications of previously published methods [20]. Chromatographic conditions were as follows: Eluent A - 6 mM octanesulfonic acid, 6 mM heptanesulfonic acid and 40 mM ammonium phosphate; Eluent B - 13 mM octanesulfonic acid and 50 mM phosphoric acid by isocratic elution program. The flow rate was 1 mL/min and injection volume - 20 µL. The separation of analytes was performed on a 250 mm × 4.6 mm i.d., 5 µm, Luna C18 reversed-phase column (Phenomenex, Aschaffenburg, Germany) equipped with a Phenomenex SecuriGuard pre-column.

Post column derivatization

To the column eluate 10 mM periodic acid in 555 mM ammonium hydroxide was added continuously at a flow of 0.4 mL/min before the reaction coil (50 °C). After the reaction coil acidification of the eluate with 0.75 M nitric acid was performed. The toxins were detected by a dual monochromator fluorescence detector (λ_{ex} 333 nm; λ_{em} 395 nm).

Lawrence method (Method 2)

The analyses were performed on a liquid chromatograph UltiMate 3000 Thermo Fisher Scientific Dionex, equipped with quaternary pump, vacuum degasser, autosampler, injector, fluorescent detector Dionex RF 2000, chromatographic column Kinetex: C18, 150 mm × 4.6 mm, 5µm; flow rate of the mobile phase 1mL/min (Eluent A: 0.1 M ammonium formate, pH 6 with 1 M acetic acid; Eluent B: 0.1 M ammonium formate in 5% acetonitrile, pH 6 with 1 M acetic acid). Elution was achieved by a gradient program (0-5% mobile phase B/5 min, 5-70% B/4 min and 100% A for the last 6 min), injection volume was 50 µL.

The identification of PSTs for both methods was accomplished by comparison of retention time and fluorescence response at specified wavelengths to certified reference materials (CRMs) (C1/2, GTX1/4, dcGTX2/3, GTX2/3, GTX5, NEO, dc-STX, STX).

To prevent false positives caused by interference of auto-fluorescent compounds, using both methods, PSP extracts were analyzed with and without derivatization.

The limits of detection (LODs) of the nine toxins for both methods were calculated (Table 1).

Table 1. LODs of investigated PSTs via HPLC-FD with post- and pre-column derivatization (NH- net haul)

PSP toxins	HPLC FD with post column derivatization		HPLC FD with pre-column derivatization
	ng/NH/m plankton	LOD ($\mu\text{g STX.2HCl eq/kg}$) mussels	mussels
C1/C2	1.4	0.2	3
GTX 1	17.5	9.4	
GTX 4			60
	13.5	7.2	
dc GTX 2	0.6	0.9	
dc GTX 3	0.6	0.9	13
GTX 2	0.7	0.4	
GTX3	0.8	0.7	15
B1	3.7	0.5	3
NEO	6.9	12.1	36
dcSTX	1.0	2.2	11
STX	0.6	1.1	15

Calculations

Based on the relative potency of each STX-analogue, by means of their effect in sodium channels and in the mouse bioassay (MBA), the European Food Safety Authority (EFSA) published the toxicity equivalency factors (TEF) [15].


As the Lawrence pre-column HPLC-FD method [16] cannot distinguish between PST stereoisomers (GTX1 and GTX4; C1 and C2, dc-GTX2 and dc-GTX3, GTX2 and GTX3) by choosing the highest toxicity factor of the two co-eluted compounds. Total PSP toxicity was calculated by summing the toxicity contribution of each quantified toxin expressed in terms of $\mu\text{g STX .2HCl eq. kg}^{-1}$.

In the present study, mussel and plankton samples were subjected to HPLC-FD analysis in order to detect the presence of the following PSP toxin types: dc-GTX2/3, C1/2, dc-STX, GTX2/3; B1, STX, NEO, GTX1/4 and dc-NEO.

Initially we applied the official AOAC (Lawrence) method for PSTs determination in shellfish samples. No PSTs were detected in any of the investigated samples with the precolumn derivatization method (Figure 2). The calculated LODs (Table 1) were much lower than the legislative threshold ($800 \mu\text{g STX.2HCl eq.kg}^{-1}$) which concluded that obviously there was no risk for PSP if investigated samples were consumed.

However, investigations on plankton species of the Institute of Oceanology of Bulgaria documented the presence of *Alexandrium spp.* in the studied area [21]. This lead us to conclude that there was a principal possibility for PSTs to be detected in plankton and mussels. Therefore, we performed PSTs analysis firstly of plankton samples but no PSTs were detected in any of the analyzed samples. This result came in agreement with our previous study on plankton samples from the same region where no PSTs were detected as well [22]. However, since plankton toxin data represents the situation at the time of sampling and is not valid for longer periods, we considered to reanalyze PSTs in mussels with a more sensitive PCOX method, as mussels accumulate and retain toxins for a longer period of time. The data obtained would be more informative and allow statements for longer periods.


Surprisingly, low levels of PSTs were detected by this method. Figure 3 shows the chromatogram of the sample with the highest toxicity (MU48P), analyzed by HPLC-FD with pre-column derivatization method.



Operator:User Timebase:hplc Sequence:PSP_Varna_08062018

Page 1-1
11/5/2018 11:24 AM

MU 48 P			
Sample Name:	samp15-MU48 P ox	Injection Volume:	50.0
Vial Number:	RB5	Channel:	Emission
Sample Type:	unknown	Wavelength:	n.a.
Control Program:	PSP_Kinetex_flow_1	Bandwidth:	n.a.
Quantif. Method:	MIX-6_PERIODAT_guard Kinetex_flow 1	Dilution Factor:	1.0000
Recording Time:	6/8/2018 12:39	Sample Weight:	
Run Time (min):	15.00	Sample Amount:	



No.	Ret.Time min	Peak Name	Height mV	Area mV*min	Rel.Area %	Amount	Type
Total:			0.000	0.000	0.00	0.000	

Fig. 2. Chromatograms of a PSP CRMs after periodate oxidation (top) and extract of cultured mussel sample MU48P (bottom) obtained via HPLC-FD with pre-column derivatization

As samples were collected regularly throughout the investigated period a seasonal change in PSTs contamination throughout the year 2017 (Figure 4) became evident.




Figure 4. Seasonal changes in PSTs levels

With respect of the overall low contamination a peak in the PST level (1.6 µg STX.2HCl /kg) was observed in early summer. This increased concentration is more than two times higher than the mean PST level, which was around 0.78 µg STX.2HCl /kg. Obvious is the presence of PSTs overall harvesting seasons with exception of early spring and late autumn. However, it is still unclear whether 2017 was a typical situation or changes in environmental conditions could lead to higher *Alexandrium* densities and PSTs levels in mussels.

In similar study conducted in the same location and during the same period of time in 2015-2016 among 55 samples analyzed both STX and GTX2,3 were detected in only one sample with total toxicity of 55.1 µg STX.2HCl /kg [23], much higher than here determined.

The two methods used have their advantages and in certain cases result of both of them should be considered. HPLC-FD with pre-column derivatization is the official method recognized by the EU legislation. Thereby detected levels are low enough to consider if mussels are suitable for consumption or not. On the other hand, the method HPLC-FD with post-column derivatization is quick and lower levels of PSTs could be detected. This is important e.g. when exposure to chronic low levels of PSTs is investigated. Even more, it is still unclear whether *Alexandrium* is really responsible for PSTs in the mussels, and whether it can come in the future to higher PST loads. Therefore, research is needed to identify sources of PSTs on the Bulgarian coast, as well as whether *Alexandrium* (as a potential source of PSP) could form harmful blooms in the Black Sea.

CONCLUSIONS

This study highlights the low contamination with PSTs in mussels off the north coast of Bulgaria in 2017. Via HPLC-FD with post column derivatization were detected GTX2 levels that were lower than the LOD of the other method used – HPLC-FD with precolumn derivatization. Thus, combination of both methods will contribute in increasing awareness about PSP and by estimation of PSTs exposure.

Acknowledgement: This work was supported by Science Fund of Medical University Varna, Project Incoming number 16012/2016 and partially funded by the Helmholtz-Gemeinschaft Deutscher Forschungszentren through the research program “Polar regions And Coasts in the changing Earth System” (PACES II) of the Alfred Wegener Institut-Helmholtz Zentrum für Polar- und Meeresforschung.

REFERENCES

1. A. Fuentes, I. Fernández-Segovia, I. Escriche, J. Serra, *Food Chem.*, **112**(2) (2009).
2. O. Ozden, S. Ulusoy, N. Erkan, *J für Verbraucherschutz und Lebensmittelsicherheit*, **5**, (3-4),407 (2010).
3. Ministry of Agriculture and Food, *Annual Report on the situation and development of agriculture*, Sofia (2016).
4. C.Y. Kao, *Algal Toxins in Seafood and Drinking Water*, Academic, San Diego, 1993.
5. N. Lagos and D. Andrinolo, in: *Seafood and freshwater toxins*, Marcel Dekker, New York, 2000, p. 203.
6. G. Hallegraeff, in: *Manual of Harmful Marine Microalgae*, G. Hallegraeff, D. Anderson and A. Cembella, (Eds) International Oceanographic

- Z.V. Peteva et al.: Evaluation of paralytic shellfish poisoning toxin profile of mussels from Bulgarian North Black Sea coast ... Commission (IOC) Manual and Guides UNESCO, Paris, 1995, p. 1.
7. A. Chapelle, M. Le Gac and C. Labry, *Harmful Algae News*, **51**, 2015.
 8. P. Costa, A. Robertson and M. Quilliam, *Marine Drugs*, **13** (4), 2046 (2015).
 9. M. Rañada, R. Tabbada, A. Mendoza, *Regional studies in Marine Sciences*, **3**, 176 (2016).
 10. A. Cembella, in: Physiological Ecology of Harmful Algal Blooms, A. Cembella and G. Hallegraeff, (Eds.), Springer, Berlin, 1998, p. 381.
 11. R. Orr, A. Stüken, T. Rundberget, W. Eikrem and K. Jakobsen, *Harmful Algae*, **10**, 676 (2011).
 12. L. Gainey, J. Shumway, S. Shumway, *J. Shellfish Res*, **7**, 623 (1988).
 13. C. Dell'Aversano, J. Walter, I. Burton, D. Stirling, E. Fattorusso and M. Quilliam, *J. Nat. Prod*, **71**, 518 (2008).
 14. L.M. Durán-Riveroll, B. Krock, A. Cembella, J. Peralta-Cruz, J.J. Bustillos-Guzman, C.H. Band-Schmidt. *Natural Products Chemistry & Research*, **5**, 275 (2017).
 15. J. Alexander, D. Benford, A. Cockburn, J.-P. Cradevi, E. Dogliotti A. Domenico et al., *EFSA J.*, **1019**, 1 (2009).
 16. AOAC. Method 2005.06: Paralytic Shellfish Poisoning Toxins in Shellfish. Prechromatographic Oxidation and Liquid Chromatography with Fluorescence Detection (First Action 2005," *AOAC INTERNATIONAL*, 83 (2006).
 17. A. Turner, M. Dhanji-Rapkova, C. Baker and M. Algoet, *Journal of AOAC International*, **97** (2), 492 (2014).
 18. R. Berlinck, M. Kossuga, *Nat. Prod. Rep.*, **22**, 516 (2005)
 19. M. S. Harwood and M. Boundy, in: Comprehensive Analytical Chemistry Recent Advances in the Analysis of Marine Toxins, J. Diogène, M. Campàs, Elsevier, Amsterdam, 2017.
 20. B. Krock, C. G. Seguel and A. D. Cembella, *Harmful Algae*, **6**(5), 734 (2007).
 21. BAS-IO, Protocols phytoplankton, 2017. http://www.bgodc.io-bas.bg/monitoring_program.html
 22. Z. Peteva, S. Georgieva, B. Krock and M. Stancheva, in: CBU International Conference Proceeding 2018: Innovations in Science and Education, [In press] (2018).
 23. G. Krumova-Valcheva, G. Kalinova, *Acta Microbiologica Bulgarica*, **33**(1), 30 (2017)