




LETTER

Seasonal patterns of microbial diversity across the world oceansEric J. Raes ^{1,2,3*}, Shannon Myles,¹ Liam MacNeil ⁴, Matthias Wietz ^{5,6}, Christina Bienhold,^{5,6} Karen Tait,⁷ Paul J. Somerfield,⁷ Andrew Bissett,⁸ Jodie van de Kamp ⁸, Josep M. Gasol,⁹ Ramon Massana ⁹, Yi-Chun Yeh,¹⁰ Jed A. Fuhrman,¹¹ Julie LaRoche ^{1*}

¹The UWA Oceans Institute, The University of Western Australia, Crawley, Australia; ²Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada; ³Minderoo Foundation, Broadway, Nedlands, Australia; ⁴GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany; ⁵Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany; ⁶Max Planck Institute for Marine Microbiology, Bremen, Germany; ⁷Plymouth Marine Laboratory, Plymouth, UK; ⁸Commonwealth Scientific and Industrial Research Organisation, Hobart, Australia; ⁹Institut de Ciències del Mar, CSIC, Barcelona, Catalonia, Spain; ¹⁰Carnegie Institution of Science, Stanford, California, USA; ¹¹Department of Biological Sciences, University of Southern California, Los Angeles, California, USA

Scientific Significance Statement

Marine microbes are essential for sustaining life in our oceans, making it crucial to monitor changes in their diversity to better understand and predict microbially driven ecosystem functions. Global oceanographic expeditions and basin-wide transects reveal positive correlations between microbial diversity and variables like temperature and productivity, but these studies often lack seasonal data and include few observations from high-latitude regions during winter. Our research shows that, despite differences in collection methods, DNA extraction protocols, targeted *16S rRNA* hypervariable regions, sequencing technologies, and bioinformatics pipelines, seasonal trends in microbial community richness and evenness remain consistent across time-series sites in both the northern and southern hemispheres.

Abstract

Understanding the patterns of marine microbial diversity (Bacteria + Archaea) is essential, as variations in their alpha- and beta-diversities can affect ecological processes. Investigations of microbial diversity from global oceanographic expeditions and basin-wide transects show positive correlations between microbial diversity and

*Correspondence: ejraes@gmail.com, Julie.LaRoche@dal.ca

Associate editor: Zachary S Feiner

Author Contribution Statement: EJR and JLR developed the initial research idea. EJR, SM, LM, and JLR wrote the first draft of the manuscript and conducted statistical analyses. SM processed Bedford Basin and SPOT data. LM conducted HGAM analyses. MW processed FRAM data. Ecological insights and interpretation were contributed by EJR, JLR, SM, and LM (Bedford Basin); MW and CB (FRAM); KT and PJS (L4); EJR, JVDK and AB (Australian sites); YY and JF (SPOT); and JMG and RM (BBMO). All authors provided editorial comments on the manuscript.

Data availability Statement: All data presented in this manuscript including detailed descriptions of methodologies and bioinformatic workflows are publicly available. The sequence data and associated metadata are deposited at institutional and international data repositories as outlined in Supplementary Materials and Methods. The code to reproduce and plot the figures presented in the manuscript is available at <https://github.com/EricRaes/Time-series-Analyses>.

A dedication to Paul Somerfield (1963–2023).

Additional Supporting Information may be found in the online version of this article.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

either temperature or productivity, but these studies rarely captured seasonality, especially in polar regions. Here, using multiannual alpha-diversity data from eight time series in the northern and southern hemispheres, we show that marine microbial community richness and evenness generally correlate more strongly with daylength than with temperature or chlorophyll *a* (a proxy for photosynthetic biomass). This pattern is observable across time series found in the northern and southern hemispheres regardless of collection method, DNA extraction protocols, targeted *16S* rRNA hypervariable region, sequencing technology, or bioinformatics pipeline.

In terrestrial ecosystems, decreased species diversity can reduce central ecosystem functions (Tilman et al. 2014), such as primary productivity (Balvanera et al. 2006; Oehri et al. 2017), and disrupt ecosystem resilience and stability through time (Cardinale et al. 2011; Gross et al. 2014; Wagg et al. 2019). The relationship between microbial diversity and ecosystem functions, however, is complex, and remains poorly understood for oceanic environments (Sunagawa et al. 2020). At the base of the marine food web, Bacteria and Archaea (herein referred to as prokaryotes) are essential for ecosystem functioning (Falkowski 1997); alterations in their diversity and distribution could affect ecological dynamics (Horner-Devine et al. 2003; Hutchins and Feixue 2017; Cavicchioli et al. 2019). Untangling regional and global patterns in prokaryotic diversity across seasonal cycles is essential to understanding, modeling, and predicting microbially driven ecosystem functions (Hatosy et al. 2013; Vallina et al. 2014; Louca et al. 2016).

Global ocean research expeditions (e.g., Tara Oceans) or ocean basin-wide transects (e.g., GO-SHIP and BIOGOTRACES) have shown a positive correlation between bacterial diversity and either temperature (Ibarbalz et al. 2019) or primary production (Raes et al. 2018a, 2018b). Although basin-wide transects cover wide biogeochemical provinces, they do not systematically address seasonality. In addition, these programs collected few or no samples during winter in polar regions. Marine time series using an Eulerian sampling design (i.e., a fixed site) provide fundamental seasonal insights into the diversity patterns of marine microbes (Wiltshire et al. 2010; Fuhrman et al. 2015; Bryant et al. 2016; Marquardt et al. 2016; Brown et al. 2018; Buttigieg et al. 2018; Lambert et al. 2019; Auladell et al. 2022) and are complementary to basin-wide observations. In particular, highly resolved (fortnightly to monthly) seasonal observations from time series suggest that marine prokaryotic community diversity is strongly correlated with daylength (e.g., Gilbert et al. 2012; Bryant et al. 2016; Marquardt et al. 2016; Giner et al. 2019; Lambert et al. 2019; Raes et al. 2022).

Trends in archaeal and bacterial alpha-diversity have mostly been linked to (i) the kinetic energy hypothesis, that is, warmer temperatures increase metabolic reaction rates which in turn, affect genetic and evolutionary traits, ultimately resulting in higher alpha-diversities (Brown 2014); or (ii) the resource hypothesis, that is, a higher energy production can support more species through niche diversification (Mittelbach

et al. 2001). We utilized the established ecological temperature framework (Brown 2014) and the resources mechanism (Mittelbach et al. 2001) to predict the drivers behind shifts in prokaryotic diversity over time, with our primary focus on temporal fluctuations within a specific location, rather than spatial distinctions between different sites. Our rationale is based on the idea that the evolutionary timeline of prokaryotes within a given location or among various sites is likely to be similar. It is important to emphasize that prokaryotic evolution unfolds at a distinct pace compared to macrofauna, with prokaryotes demonstrating significant responsiveness to alterations in their environmental conditions, primarily due to their higher cell division rates (Hillebrand et al. 2022).

Here, we use seasonal data from eight time series in the northern and southern hemispheres (from 79°N to 42°S) to test the hypothesis that, on a multiannual basis, marine prokaryotic alpha-diversity (richness and evenness) correlates more strongly with daylength than with temperature or productivity. Regardless of the various factors that can introduce variation in prokaryotic diversity studies, we demonstrate the generality of a recurrent yearly cycle in community structure.

Materials and methods

Selection of time-series sites for perspective on prokaryotic community diversity

16S ribosomal RNA gene (rRNA) metabarcoding data were retrieved from eight time series (Fig. 1; Table 1), alongside physical and biochemical metadata. These include (1) the Arctic FRAM observatory at the long-term ecological research (LTER) site HAUSGARTEN at F4 and HG-IV moorings; (2) the English Channel coastal L4 (ECL4) site; (3) the Compass Buoy Station HLO in the Bedford Basin (BBNS), Canada; (4) the Blanes Bay Microbial Observatory LTER (BBMO), Spain; (5) the San Pedro Ocean Time Series (SPOT), California, U.S.A.; (6) Integrated Marine Observing System (IMOS) National Reference Stations (NRS) Yongala (YON), Australia; (7) NRS Rottneest Island (ROT), Australia; and (8) NRS Maria Island (MAI), Australia. Data obtained were generated from samples in the euphotic zone (shallower than 50 m depth) from polar, temperate, and subtropical climate zones. FRAM samples were collected autonomously using moored Remote Access Samplers (RAS; McLane), whereas all other samples were collected manually with Niskin bottles or a bucket (BBMO). The Supplementary Materials and Methods provide details on the selection criteria, discrete sampling depths at




Fig. 1. Legend on next page.

each site and protocols for physical and biogeochemical metadata.

DNA sampling, sequencing, and bioinformatics

Detailed information on sample collection dates, targeted 16S rRNA gene hypervariable regions, primers, DNA extraction, sequencing technology, and bioinformatics are shown in Table 1 and correspond to the individual specifications of the original publications. No reprocessing of the sequencing data was conducted and the original ASV/OTU tables were used for statistical analyses except for SPOT and Bedford Basin, whose raw sequencing data were reprocessed identically as originally published (Raes et al. 2022; Yeh and Fuhrman 2022a). Details on bioinformatic workflows are shown in the Supplementary Materials and Methods. The V4–V5 and V6 16S rRNA regions target Archaea, Bacteria, and Eukarya (Lane et al. 1985; Weisburg et al. 1991; Lee and Gutell 2012; Parada et al. 2016), but eukaryotic information was not analyzed here.

Statistical analyses

Data analyses, visualization, and maps were generated with R (v4.2.2; R Core Team 2014) in RStudio (v.1.3.1093; see our GitHub). Statistical tests and correlations were conducted between sample points matching in time (i.e., alpha diversity, temperature, chlorophyll *a* (Chl *a*) concentrations and daylength at time x_1). To visualize changes in beta-diversity over time, pairwise Aitchison's distances were calculated between each possible pair of samples (Fig. 2, y -axis) and recorded over the time difference between those samples, in days (Fig. 2, x -axis); mean values and associated 95% confidence intervals are plotted for every 10-d interval (Fig. 2) using ggplot2 (v.1.3.2; Wickham et al. 2019). Centered log-ratio (CLR) transformations of non-rarefied ASV/OTU tables with an added pseudo-count of 1 for zeros and distances were calculated using the Vegan package (v.2.6.4; Oksanen et al. 2007). To evaluate the response of diversity to different predictors under the three hypotheses described here (daylength, temperature, Chl *a*), we tested four versions (model S1, model G, model S2, and model GS following Pedersen et al. 2019) of hierarchical generalized additive models (HGAMs) in R package mgcv (Wood 2017). HGAMs were tested with richness and Shannon's diversity as response variables. See Supplementary Section 3.7 for model details.

Results and discussion

Physical and biochemical seasonality

Seasonality of physical and bio-geochemical parameters is presented in Supplementary Fig. S8A–H. Daylength, as

determined from latitude and time of year, exhibited the smallest and largest seasonal differences in the tropics (2.3 h) and in the Arctic (24 h), respectively. Overall, seasonality was apparent in the physical and biogeochemical parameters at all stations, except for YON, with oligotrophic conditions and low phytoplankton biomass throughout multiple years. Temperatures across the eight sites in the northern and southern hemisphere ranged from -1.7°C to 30°C . Dissolved inorganic nitrate and nitrite (NO_x) concentrations across all sites ranged from below detection limits ($< 0.02 \mu\text{mol L}^{-1}$) to $15 \mu\text{mol L}^{-1}$ at the ECL4. Highest NO_x concentrations occurred in the winter months, except at ROT (highest in summer) and YON (oligotrophic throughout the year). Chl *a* concentrations, a proxy for photosynthetic biomass, peaked in spring and autumn in BBNS (up to $40 \mu\text{g L}^{-1}$), in early spring in SPOT (up to $19.8 \mu\text{g L}^{-1}$), in spring and autumn in the ECL4 (up to $4.76 \mu\text{g L}^{-1}$), in winter and spring in BBMO (up to $2.9 \mu\text{g L}^{-1}$), and in summer in the FRAM (up to $2.9 \mu\text{g L}^{-1}$). In the southern hemisphere, productivity was higher in spring at MAI (up to $1.6 \mu\text{g L}^{-1}$) and in autumn at ROT (up to $0.9 \mu\text{g L}^{-1}$). Low Chl *a* concentrations ($< 0.5 \mu\text{g L}^{-1}$) were recorded at YON throughout the year, though with a trend of relatively higher concentrations during the summer months.

Consistent patterns in seasonality regardless of sampling and metabarcoding methodology

Despite different amplification targets and different resolution (OTUs vs. ASVs), all sites showed significant seasonality in prokaryotic diversity (community richness and evenness; Fig. 1). Highest species richness and evenness were always recorded in the winter ($2\text{--}3\times$ greater than in summer), even at the tropical YON ($p < 0.05$ for all Wilcoxon tests between summer and winter). These trends remained regardless of rarefaction depths and were also observed when considering the rare microbiome (OTUs/ASVs composing $< 1\%$ community proportions; Supplementary Figs. S9–S16).

The choice of target 16S rRNA variable region is tailored to specific research question and particular taxa (Choi et al. 2017), sequencing technology (Cruaud et al. 2014), and intercomparisons of results to other environments or older studies (Gilbert et al. 2009; Thompson et al. 2017; Brown et al. 2018). Considering the different PCR biases of primer sets, there is currently no perfect universal primer set for targeting variable 16S rRNA regions across microorganisms (McNichol et al. 2021). Several studies have investigated how different primers capture true prokaryotic diversity (e.g., Bukin et al. 2019; Willis et al. 2019; Soriano-Lerma et al. 2020). Yet, here we show a consistent signal in seasonal

Fig. 1. Global seasonal trends in prokaryotic alpha-diversity. Top: World map illustrating the location of eight time series Fram Strait (FRAM), English Channel (ECL4), Bedford Basin (BBNS), Blanes Bay (BBMO), San Pedro (SPOT), NRS Yongala (YON), NRS Rottneest Island (ROT), and NRS Maria Island (MAI). For each site, individual panels show alpha-diversity trends across week number including richness (Chao1), Shannon, and Pielou metrics. Data points are colored by sampling month, with Loess regression lines (in blue with 95% confidence band in gray) fitted to the alpha-diversity metrics. The x -axis shows time in number of weeks, with one corresponding to the first week in January and 52 corresponding to the last week in December.

Table 1. Time-series locations, climate zones, sampling schemes, and sequencing metadata.

Properties	Fram Strait, mooring									
	F4	F4	HG-IV	English Channel	Bedford Basin	Blanes Bay	San Pedro	Yongala	Rottneest Island	Maria Island
Acronym	FRAM	FRAM	FRAM	ECL4	BBNS	BBMO	SPOT	YON	ROT	MAI
Latitude (N)	79°0.708'	79°1.380'	79°1.380'	50°15'	44°41'37"	41°40'	33°30'	-19°18'	-32°0'	-42°36'
Longitude (E)	6°57.888"	4°15.708'	4°15.708'	-4°13'	-63°38'25"	2°48'	-118°30'	147°37'	115°25'	148°14'
Climate zone	Polar	Polar	Polar	Temperate	Temperate	Temperate	Temperate	Tropical	Subtropical	Temperate
Biogeochemical Province	Boreal polar/ Atlantic Arctic	Boreal polar/ Atlantic Arctic	Boreal polar/ Atlantic Arctic	Northeast Atlantic shelfe	Northwest Atlantic shelfe	Mediterranean Sea	Coastal Californian current	East Australian coast	West Australian coast	Tasman Sea
Depths sampled (m)	24-42	24-42	26-48	Surface	1, 5, 10	1	5	0, 10, 20, 26	0, 10, 20, 30, 40, 46	0, 10, 20, 50
Sampling frequency	Biweekly	Biweekly	Biweekly	Monthly	Weekly	Monthly	Monthly	Monthly	Monthly	Monthly
Sampling period	2016-2018	2016-2018	2016-2018	2003-2008	2014-2017	2004-2013	2005-2018	2015-2020	2015-2020	2012-2020
Total samples	46	47	47	72	191	120	147	54	58	89
16S rRNA region	V4-V5	V4-V5	V4-V5	V6	V4-V5	V4	V4-V5	V1-V3	V1-V3	V1-V3
Region length (bp)	~410	~410	~410	~68	~410	~435	~410	~490	~490	~490
Primers	515F + 926R*	515F + 926R*	515F + 926R*	967F + 1046R†	515F + 926R*	341F + 806RB‡	515Y + 926R*	27F + 519R§	27F + 519R§	27F + 519R§
Sequencing technology	Illumina MiSeq	Illumina MiSeq	Illumina MiSeq	454-pyrosequencing	Illumina MiSeq	Illumina MiSeq	Illumina HiSeq	Illumina MiSeq	Illumina MiSeq	Illumina MiSeq
Sequence denoising/ clustering (ASV/OTU)	ASV (DADA2)	ASV (DADA2)	ASV (DADA2)	OTU97 (DOTUR)	ASV (DADA2)	ASV (DADA2)	ASV (DADA2)	ASV (UNOISE 3)	ASV (UNOISE 3)	ASV (UNOISE 3)

Latitude and longitude in degrees minutes.

*Parada et al. (2016).

†Sogin et al. (2006).

‡Herlemann et al. (2011) and Apprill et al. (2015).

§Lane et al. (1985) and Weisburg et al. (1991).




Fig. 2. Global seasonal trends in prokaryotic community beta-diversity in the euphotic zone. Site-specific distributions of pairwise Aitchison distances between community compositions of each possible time difference (x -axis) between sample collections (95% confidence intervals around estimates of the mean every 10 d). Data from Fram Strait (FRAM), English Channel (ECL4), Bedford Basin (BNNS), Blanes Bay (BBMO), San Pedro Timeseries (SPOT), NRS Yongala (YON), NRS Rottneest Island (ROT), and NRS Maria Island (MAI) (panels from top to bottom, left to right, respectively). Dotted lines are shown every 365 d. The marine heatwave (the “BLOB”) is denoted by an arrow on SPOT between 2014 and 2015. Aitchison distance is the Euclidian distance calculated for the prokaryotic community composition; it is based on read counts which were center logratio transformed.

alpha-diversity across sites, regardless of location or targeted *16S rRNA* gene hypervariable region (V1–V3, V4, V4–V5, V6), DNA extraction methodology, or even sample collection method (FRAM samples were collected autonomously and

preserved with mercury chloride). Furthermore, our analyses display a recurring cyclical pattern in prokaryotic beta-diversity, with the strongest signal at the temperate and polar sites (Fig. 2; Supplementary Fig. S17). The only exceptions

occurred at YON, where a “drifting” signal characterized the community as diversity became more dissimilar over time (Fig. 2). A possible explanation could be that biological interactions have a greater impact on that community as daylength, and oligotrophic conditions remain relatively constant year-round as is typical for tropical oceans. The recurring compositional cycle at SPOT was disrupted around 2014–2015 (Fig. 2; Yeh and Fuhrman 2022b), due to an anomalous major heating event across the northeast Pacific Ocean related to El Niño (“the Blob”), which impacted the whole marine food web (Cavole et al. 2016; Traving et al. 2021).

Kinetic energy, resource, and daylength hypotheses

Various environmental and ecological factors correlate with marine prokaryotic diversity (reviewed in Fuhrman et al. 2015; Ibarbalz et al. 2019). The kinetic energy and resource hypotheses provide mechanisms for changes in marine prokaryotic community diversity. The kinetic energy hypothesis postulates that elevated temperature leads to higher local diversity, through increased metabolic rates, accelerated physiological processes, shorter generation times, and ultimately producing a more diverse community. The resource hypothesis postulates that increasing productivity scales with expanding diversity due to the greater resource availability, which can support a greater number of species (Brown 2014; Ibarbalz et al. 2019). We used log-scaled Chl *a* concentrations, although it is notably not the best estimator of resource limitation for prokaryotes, particularly in very oligotrophic sites and during summer where a diverse bacterial component support pigment diversification (Auladell et al. 2022). Although the kinetic energy and resource hypotheses provide a framework to explain global marine biodiversity patterns, individual time-series observations have shown that prokaryotic richness and evenness are negatively correlated with daylength (Gilbert et al. 2012; Wietz et al. 2021; Raes et al. 2022; Doane et al. 2023).

In addition to linear correlations and multiple linear regression models (Supplementary Figs. S18–S23) we explored a form of hierarchical generalized additive models (HGAMs), which provide flexible predictions of nonlinear relationships (Hastie and Tibshirani, 1986). We show, using HGAMs, that the negative correlation between prokaryotic alpha-diversity and daylength persists on a multiannual basis across sites in both northern and southern hemispheres. The primary finding of the HGAMs indicates a prominent seasonality effect: knowing the month, site, and year of an observation accounts for >80% of deviance explained (Fig. 3; Supplementary Table S1). Including daylength, temperature, and chlorophyll as fixed effects into HGAMs with global (model “G”) and site-specific (model “S2”) smoothers broadly reproduced similar patterns shown in linear regressions. Visualizing the partial dependence curves shows how model “G” captures a strong negative relationship between diversity and daylength, a positive relationship to temperature up to $\sim 20^{\circ}\text{C}$ —resembling a

thermal performance curve—and a right-skewed relationship to chlorophyll with a peak at $\sim 1 \mu\text{g L}^{-1}$. Alternatively, model S2 allows variation using site-specific smoothers, yet only 1% additional deviance was explained compared to model G. Here, the relationship between diversity and temperature or chlorophyll are mixed across sites, although still significant, and a consistently negative relationship to daylength persists. Daylength emerges as a key statistical variable explaining patterns in prokaryotic diversity across multiple time series in the northern and southern hemispheres factor across the entire dataset, albeit with some variation at site level.

A clear example of site level variation is noticed for the tropical site YON, where annual variability in daylength is minimal (Fig. 3). Similarly, for the Arctic site FRAM, the absolute Pearson correlation coefficients between $\log(\text{Chl } a)$ and Shannon diversity was higher compared to the correlation coefficient between Shannon diversity and day length (Supplementary Fig. S19). In the Arctic, extreme seasonal shifts between summer and winter dictate changes in communities. Light triggers primary productivity, leading to lower prokaryotic diversity in summer and higher diversity in winter. In winter, prokaryotic niche diversity increases and a diverse group of prokaryotes contributes to nutrient replenishment in winter (Wietz et al. 2021). The only positive relationship with $\log(\text{Chl } a)$ concentrations was found for ROT (<7% explained), with daylength still explaining between 15% and 24% of the trend in prokaryotic diversity (for Chao1 and Shannon, respectively; Supplementary Figs. S18, S19). A positive relationship between productivity and prokaryotic community richness has previously been noted by Raes et al. (2018a, 2018b) from a single transect along the Leeuwin Current flowing past Rottneest Island, explained by niche partitioning between nitrate-driven autotrophic and mixotrophic micro-eukaryotes. Multiple linear regression models (with temperature + $\log(\text{Chl } a)$ + daylength fitted last) corroborated our results (Supplementary Fig. S23). In a previous study we used Partial Mantel tests to separate the effects of day length and temperature on prokaryotic beta diversity. Despite the high temporal resolution of the Bedford Basin data (weekly over 4 years), it remained difficult to distinguish between light and temperature effects (Raes et al. 2022).

Studies at large spatial scales showed that temperature is the main (and positively correlated) explanatory variable for microbial community diversity in marine ecosystems (Fuhrman et al. 2008; Ibarbalz et al. 2019). Our analyses show that the relationship with temperature varied between sites; encompassing negative, positive, or no correlation with prokaryotic richness or evenness (Supplementary Figs. S6, S18, S19).

Our findings do not support the kinetic energy nor the resource hypothesis to explain the factors influencing prokaryotic diversity (Mittelbach et al. 2001; Ibarbalz et al. 2019). Instead, our analyses closely align with Ladau et al. (2013),




Fig. 3. Hierarchical generalized additive model plots (HGAM) showing the partial effects of explanatory variables: **(A)** daylength, **(B)** temperature, and **(C)** log (Chl *a*) on richness. The *y*-axis represents the contribution of a specific predictor to the response variable, after accounting for the effects of other predictors in the model (Supplementary Table S1). Positive values indicate an increase in the response variable, while negative values indicate a decrease. HGAM plots are shown for model G which uses global smoothers and random intercepts for sites (random effect). Gaussian quantiles for model G are shown in Supplementary Fig. S5. **(D)** Model S2 partial dependence curves for day length with site-specific smoothers and random intercepts for sites (random effect; see also Supplementary Fig. S6; Table S1). Global smoothers use default thin plate regression splines (TPRS) for each fixed effect covariate. Ticks are shown on each *x*-axis for the distribution of observed data points.

highlighting the primary impact of seasonality on regional, and potentially global, prokaryotic diversity. The true drivers for prokaryotic community diversity are difficult to identify with certainty as many parameters co-vary with daylength, and many potentially important parameters were not assessed here. Hence, it is not an “either/or verdict” for the hypotheses that were considered here, but rather a complex interplay of processes that dominate and differ when sampling at different spatial and temporal scales. We demonstrate that prokaryotic diversity patterns investigated locally, with seasonal resolution, are affected by different environmental drivers compared to large spatial studies, which usually do not account for temporal or seasonal variations. Our findings indicate that in high-latitude regions, diversity must be interpreted in the context of the particularly strong seasonal cycle, with highest diversity during the polar night. Hence, seasonality and local variation must be considered to understand marine prokaryotic community diversity, especially when comparing globally distributed sites.

The relevance of daylength has been described as a variable that integrates seasonal variability in temperature and stratification (Wietz et al. 2021). Shorter day length co-occurs with seasonal vertical mixing, thus the passive merging of surface (photic)

and deep (aphotic) prokaryotic communities resulting in higher diversity (García et al. 2015). In addition, increased daylength and associated higher solar irradiance might have negative (bactericidal) effects on photosensitive microbes that are poorly adapted to high solar irradiance (Ruiz-González et al. 2013). Alonso-Sáez et al. (2006), for example, carried out experimental work that suggested that some groups of non-photosynthetic, heterotrophic bacteria may be negatively impacted by light, as they lack bacteriorhodopsin or other light harvesting photosystems as well as protective mechanisms against UV radiation.

A multiannual recurring state

Time-series data allow the description of long-term seasonal patterns and their associated variability, affording explicit tests for how changes in biological diversity impact ecosystem functions and ecological stability (Fuhrman et al. 2015; Benway et al. 2019). This advantage of time series is unique in comparison to larger-scale spatial studies (O'Brien et al. 2017) such as Tara Oceans, which cannot capture seasonality. Our analyses confirm clear boundaries for prokaryotic community variability over multiannual scales, from subtropical to polar oceans. While the kinetic energy or resource hypotheses are not mutually exclusive—temperature

and Chl *a* covary—our findings and those from Fuhrman et al. (2015) show that daylength explains the largest variation in prokaryotic alpha- and beta-diversity. Daylength, as a variable, integrates various seasonal processes, including net heat flux, wind speed, total daily irradiance, stratification, depletion, and regeneration of nutrients, as well as the release of POC and DOC. These processes create a recurring pattern of changes in diversity that are not solely driven by temperature or Chl *a*.

Conclusions

Our results showed that daylength is statistically a strong variable accounting for prokaryotic diversity patterns across six of the eight independent prokaryotic time series. However, the inverse relationship between daylength and diversity is counterintuitive, and does not readily link to an underlying biological mechanism. Although physical mixing and seasonality in light-sensitive taxa are potential ecological mechanisms, other causal processes are possible, and a unifying mechanistic explanation still eludes formal description. Our study shows the necessity for considering the temporal dimension in both regional and global prokaryotic diversity analyses, as diversity trends are connected to our planet's seasonal rhythm. Diversity was up to threefold higher in winter across the majority of the eight time series analyzed. The highest variation was observed between the summer and winter months, with clear annually recurring seasonal beta-diversity patterns in polar and temperate waters. Notably, these patterns were independent of collection methods, DNA extraction chemistry, targeted *16S* rRNA hypervariable region, sequencing technology, resolution of taxonomic units, or bioinformatics pipeline. Our findings underscore the potential for improved global and networked observational initiatives by harmonizing methods so we can address comparisons of absolute diversity. This emphasizes the value in establishing standardized best practices for genomic data acquisition and reporting, encompassing the creation of a Minimum Information for an Omics Protocol, as recommended by Samuel et al. (2021). Such harmonization will promote data interoperability, facilitate collaborative research efforts, and promote comparisons of absolute diversity values between sites, which our work cannot address. Where climate changes do generate more frequent extreme events, this work highlights a potential, hitherto undescribed process whereby prokaryotic community fluctuations in response to an interference with the regular daylength-correlated cycles can be identified (see, e.g., the “Blob” 2-yr heatwave event Fig. 2e). We posit that based on the prokaryotic time series presented here, among the richest datasets yet available, the empirical evidence supports a primary role for daylength in explaining the seasonal diversity and ecological rhythm of prokaryotic microbes, the basal component of marine food webs, and warrants deeper investigation for causal processes.

References

- Alonso-Sáez, L., J. M. Gasol, T. Lefort, J. Hofer, and R. Sommaruga. 2006. Effect of natural sunlight on bacterial activity and differential sensitivity of natural bacterioplankton groups in northwestern Mediterranean coastal waters. *Appl. Environ. Microbiol.* **72**: 5806–5813. doi:[10.1128/AEM.00597-06](https://doi.org/10.1128/AEM.00597-06)
- Apprill, A., S. McNally, R. Parsons, and L. Weber. 2015. Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquat. Microb. Ecol.* **75**: 129–137. doi:[10.3354/ame01753](https://doi.org/10.3354/ame01753)
- Auladell, A., A. Barberán, R. Logares, E. Garcés, J. M. Gasol, and I. Ferrera. 2022. Seasonal niche differentiation among closely related marine bacteria. *ISME J.* **16**: 178–189. doi:[10.1038/s41396-021-01053-2](https://doi.org/10.1038/s41396-021-01053-2)
- Balvanera, P., A. B. Pfisterer, N. Buchmann, J.-S. He, T. Nakashizuka, D. Raffaelli, and B. Schmid. 2006. Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecol. Lett.* **9**: 1146–1156. doi:[10.1111/j.1461-0248.2006.00963.x](https://doi.org/10.1111/j.1461-0248.2006.00963.x)
- Benway, H. M., and others. 2019. Ocean time series observations of changing marine ecosystems: An era of integration, synthesis, and societal applications. *Front. Mar. Sci.* **6**: 393. doi:[10.3389/fmars.2019.00393](https://doi.org/10.3389/fmars.2019.00393)
- Brown, J. H. 2014. Why are there so many species in the tropics? *J. Biogeogr.* **41**: 8–22. doi:[10.1111/jbi.12228](https://doi.org/10.1111/jbi.12228)
- Brown, M. V., and others. 2018. Systematic, continental scale temporal monitoring of marine pelagic microbiota by the Australian marine microbial biodiversity initiative. *Sci. Data* **5**: 180130. doi:[10.1038/sdata.2018.130](https://doi.org/10.1038/sdata.2018.130)
- Bryant, J. A., F. O. Aylward, J. M. Eppley, D. M. Karl, M. J. Church, and E. F. DeLong. 2016. Wind and sunlight shape microbial diversity in surface waters of the North Pacific subtropical gyre. *ISME J.* **10**: 1308–1322. doi:[10.1038/ismej.2015.221](https://doi.org/10.1038/ismej.2015.221)
- Bukin, Y. S., Y. P. Galachyants, I. V. Morozov, S. V. Bukin, A. S. Zakharenko, and T. I. Zemskaia. 2019. The effect of 16S rRNA region choice on bacterial community Metabarcoding results. *Sci. Data* **6**: 190007. doi:[10.1038/sdata.2019.7](https://doi.org/10.1038/sdata.2019.7)
- Buttigieg, P. L., E. Fadeev, C. Bienhold, L. Hehemann, P. Offre, and A. Boetius. 2018. Marine microbes in 4D—Using time series observation to assess the dynamics of the ocean microbiome and its links to ocean health. *Curr. Opin. Microbiol.* **43**: 169–185. doi:[10.1016/j.mib.2018.01.015](https://doi.org/10.1016/j.mib.2018.01.015)
- Cardinale, B. J., K. L. Matulich, D. U. Hooper, J. E. Byrnes, E. Duffy, L. Gamfeldt, P. Balvanera, M. I. O'Connor, and A. Gonzalez. 2011. The functional role of producer diversity in ecosystems. *Am. J. Bot.* **98**: 572–592. doi:[10.3732/ajb.1000364](https://doi.org/10.3732/ajb.1000364)
- Cavicchioli, R., and others. 2019. Scientists' warning to humanity: Microorganisms and climate change. *Nat. Rev. Microbiol.* **17**: 569–586. doi:[10.1038/s41579-019-0222-5](https://doi.org/10.1038/s41579-019-0222-5)
- Cavole, L., and others. 2016. Biological impacts of the 2013–2015 warm-water anomaly in the Northeast Pacific:

- Winners, losers, and the future. *Oceanography* **29**: 273–285. doi:[10.5670/oceanog.2016.32](https://doi.org/10.5670/oceanog.2016.32)
- Choi, C. J., C. Bachy, G. S. Jaeger, C. Poirier, V. V. S. S. Lisa Sudek, A. M. Sarma, S. J. Giovannoni, and A. Z. Worden. 2017. Newly discovered deep-branching marine plastid lineages are numerically rare but globally distributed. *Curr. Biol.* **27**: R15–R16. doi:[10.1016/j.cub.2016.11.032](https://doi.org/10.1016/j.cub.2016.11.032)
- Cruaud, P., A. Vigneron, C. Lucchetti-Miganeh, P. E. Ciron, A. Godfroy, and M.-A. Cambon-Bonavita. 2014. Influence of DNA extraction method, 16S rRNA targeted hypervariable regions, and sample origin on microbial diversity detected by 454 pyrosequencing in marine chemosynthetic ecosystems. *Appl. Environ. Microbiol.* **80**: 4626–4639. doi:[10.1128/AEM.00592-14](https://doi.org/10.1128/AEM.00592-14)
- Doane, M. P., and others. 2023. Defining marine bacterioplankton community assembly rules by contrasting the importance of environmental determinants and biotic interactions. *Environ. Microbiol.* **1462–2920**: 16341. doi:[10.1111/1462-2920.16341](https://doi.org/10.1111/1462-2920.16341)
- Falkowski, P. G. 1997. Evolution of the nitrogen cycle and its influence on the biological sequestration of CO₂ in the ocean. *Nature* **387**: 272–275. doi:[10.1038/387272a0](https://doi.org/10.1038/387272a0)
- Fuhrman, J. A., J. A. Cram, and D. M. Needham. 2015. Marine microbial community dynamics and their ecological interpretation. *Nat. Rev. Microbiol.* **13**: 133–146. doi:[10.1038/nrmicro3417](https://doi.org/10.1038/nrmicro3417)
- Fuhrman, J. A., J. A. Steele, I. Hewson, M. S. Schwabach, M. V. Brown, J. L. Green, and J. H. Brown. 2008. A latitudinal diversity gradient in planktonic marine bacteria. *Proc. Natl. Acad. Sci. U.S.A.* **105**: 7774–7778. doi:[10.1073/pnas.0803070105](https://doi.org/10.1073/pnas.0803070105)
- García, F. C., L. Alonso-Sáez, X. A. Morán, and Á. López-Urrutia. 2015. Seasonality in molecular and cytometric diversity of marine bacterioplankton: The re-shuffling of bacterial taxa by vertical mixing. *Environ. Microbiol.* **17**: 4133–4142. doi:[10.1111/1462-2920.12984](https://doi.org/10.1111/1462-2920.12984)
- Gilbert, J. A., D. Field, P. Swift, L. Newbold, A. Oliver, T. Smyth, P. J. Somerfield, S. Huse, and I. Joint. 2009. The seasonal structure of microbial communities in the Western English Channel. *Environ. Microbiol.* **11**: 3132–3139. doi:[10.1111/j.1462-2920.2009.02017.x](https://doi.org/10.1111/j.1462-2920.2009.02017.x)
- Gilbert, J. A., and others. 2012. Defining seasonal marine microbial community dynamics. *ISME J.* **6**: 298–308. doi:[10.1038/ismej.2011.107](https://doi.org/10.1038/ismej.2011.107)
- Giner, C. R., V. Balagué, A. K. Krabberød, I. Ferrera, A. Reñé, E. Garcés, J. M. Gasol, R. Logares, and R. Massana. 2019. Quantifying long-term recurrence in planktonic microbial eukaryotes. *Mol. Ecol.* **28**: 923–935. doi:[10.1111/mec.14929](https://doi.org/10.1111/mec.14929)
- Gross, K., B. J. Cardinale, J. W. Fox, A. Gonzalez, H. Michel Loreau, W. Polley, P. B. Reich, and J. van Ruijven. 2014. Species richness and the temporal stability of biomass production: A new analysis of recent biodiversity experiments. *Am. Nat.* **183**: 1–12. doi:[10.1086/673915](https://doi.org/10.1086/673915)
- Hastie, T., and R. Tibshirani. 1986. Generalized additive models. *Stat. Sci.* **1**: 297–318. doi:[10.1201/9780203738535](https://doi.org/10.1201/9780203738535)
- Hatosy, S. M., J. B. H. Martiny, R. Sachdeva, J. Steele, J. A. Fuhrman, and A. C. Martiny. 2013. Beta diversity of marine bacteria depends on temporal scale. *Ecology* **94**: 1898–1904. doi:[10.1890/12-2125.1](https://doi.org/10.1890/12-2125.1)
- Herlemann, D. P. R., M. Labrenz, K. Jürgens, S. Bertilsson, J. J. Waniek, and A. F. Andersson. 2011. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J.* **5**: 1571–1579. doi:[10.1038/ismej.2011.41](https://doi.org/10.1038/ismej.2011.41)
- Hillebrand, H., E. Acevedo-Trejos, S. D. Moorthi, A. Ryabov, M. Striebel, P. K. Thomas, and M. L. Schneider. 2022. Cell size as driver and sentinel of phytoplankton community structure and functioning. *Funct. Ecol.* **36**: 276–293. doi:[10.1111/1365-2435.13986](https://doi.org/10.1111/1365-2435.13986)
- Horner-Devine, C. M., M. A. Leibold, V. H. Smith, and B. J. M. Bohannan. 2003. Bacterial diversity patterns along a gradient of primary productivity. *Ecol. Lett.* **6**: 613–622. doi:[10.1046/j.1461-0248.2003.00472.x](https://doi.org/10.1046/j.1461-0248.2003.00472.x)
- Hutchins, D. A., and F. Feixue. 2017. Microorganisms and ocean global change. *Nat. Microbiol.* **2**: 1–11. doi:[10.1038/nmicrobiol.2017.58](https://doi.org/10.1038/nmicrobiol.2017.58)
- Ibarbalz, F. M., and others. 2019. Global trends in marine plankton diversity across kingdoms of life. *Cell* **179**: 1084–1097.e21. doi:[10.1016/j.cell.2019.10.008](https://doi.org/10.1016/j.cell.2019.10.008)
- Ladau, J., T. J. Sharpton, M. M. Finucane, G. Jospin, S. W. Kembel, J. O'Dwyer, A. F. Koeppel, J. L. Green, and K. S. Pollard. 2013. Global marine bacterial diversity peaks at high latitudes in winter. *ISME J.* **7**: 1669–1677. doi:[10.1038/ismej.2013.37](https://doi.org/10.1038/ismej.2013.37)
- Lambert, S., M. Tragin, J.-C. Lozano, J.-F. Ghiglione, D. Vaultot, F.-Y. Bouget, and P. E. Galand. 2019. Rhythmicity of coastal marine Picoeukaryotes, bacteria and archaea despite irregular environmental perturbations. *ISME J.* **13**: 388–401. doi:[10.1038/s41396-018-0281-z](https://doi.org/10.1038/s41396-018-0281-z)
- Lane, D. J., B. Pace, G. J. Olsen, D. A. Stahl, M. L. Sogin, and N. R. Pace. 1985. Rapid determination of 16S ribosomal RNA sequences for phylogenetic analyses. *Proc. Natl. Acad. Sci. U.S.A.* **82**: 6955–6959. doi:[10.1073/pnas.82.20.6955](https://doi.org/10.1073/pnas.82.20.6955)
- Lee, J. C., and R. R. Gutell. 2012. A comparison of the crystal structures of eukaryotic and bacterial SSU ribosomal RNAs reveals common structural features in the hypervariable regions. *PLoS One* **7**: e38203. doi:[10.1371/journal.pone.0038203](https://doi.org/10.1371/journal.pone.0038203)
- Louca, S., and others. 2016. Integrating biogeochemistry with multiomic sequence information in a model oxygen minimum zone. *Proc. Natl. Acad. Sci. USA* **113**: E5925–E5933. doi:[10.1073/pnas.1602897113](https://doi.org/10.1073/pnas.1602897113)
- Marquardt, M., A. Väder, E. I. Stübner, M. Reigstad, and T. M. Gabrielsen. 2016. Strong seasonality of marine microbial eukaryotes in a high-Arctic fjord (Isfjorden, in West Spitsbergen, Norway). *Appl. Environ. Microbiol.* **82**: 1868–1880. doi:[10.1128/AEM.03208-15](https://doi.org/10.1128/AEM.03208-15)

- McNichol, J., P. M. Berube, S. J. Biller, and J. A. Fuhrman. 2021. Evaluating and improving small subunit rRNA PCR primer coverage for bacteria, archaea, and eukaryotes using metagenomes from Global Ocean surveys. *MSystems* **6**: e0056521. doi:10.1128/mSystems.00565-21
- Mittelbach, G. G., C. F. Steiner, S. M. Scheiner, K. L. Gross, H. L. Reynolds, R. B. Waide, M. R. Willig, S. I. Dodson, and L. Gough. 2001. What is the observed relationship between species richness and productivity? *Ecology* **82**: 2381–2396. doi:10.1890/0012-9658(2001)082[2381:WITORB]2.0.CO;2
- O'Brien, T. D., L. Lorenzoni, K. Isensee, and L. Valdes. 2017. What are marine ecological time series telling us about the ocean? A status report. IOC Technical Series No. 129. IOC-UNESCO. Available from https://www.uncclearn.org/wp-content/uploads/library/ioc_unesco.pdf
- Oehri, J., B. Schmid, G. Schaepman-Strub, and P. A. Niklaus. 2017. Biodiversity promotes primary productivity and growing season lengthening at the landscape scale. *Proc. Natl. Acad. Sci. USA* **114**: 10160–10165. doi:10.1073/pnas.1703928114
- Oksanen, J., and others. 2007. Vegan: Community ecology package. Available from <https://CRAN.R-project.org/package=vegan>
- Parada, A. E., D. M. Needham, and J. A. Fuhrman. 2016. Every base matters: Assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environ. Microbiol.* **18**: 1403–1414. doi:10.1111/1462-2920.13023
- Pedersen, E. J., D. L. Miller, G. L. Simpson, and N. Ross. 2019. Hierarchical generalized additive models in ecology: An introduction with mgcv. *PeerJ*. **7**: e6876. doi:10.7717/peerj.6876
- R Core Team. 2014. R: A language and environment for statistical computing. MSOR Connections 1. Available from <https://www.semanticscholar.org/paper/R%3A-A-language-and-environment-for-statistical-Team/659408b243cec55de8d0a3bc51b81173007aa89b>
- Raes, E. J., L. Bodrossy, J. van de Kamp, A. Bissett, M. Ostrowski, M. V. Brown, S. L. S. Sow, B. Sloyan, and A. M. Waite. 2018a. Oceanographic boundaries constrain microbial diversity gradients in the South Pacific Ocean. *Proc. Natl. Acad. Sci. USA* **115**: E8266–E8275. doi:10.1073/pnas.1719335115
- Raes, E. J., L. Bodrossy, J. van de Kamp, A. Bissett, and A. M. Waite. 2018b. Marine bacterial richness increases towards higher latitudes in the Eastern Indian Ocean. *Limnol. Oceanogr. Lett.* **3**: 10–19. doi:10.1002/loi2.10058
- Raes, E. J., J. Tolman, D. Desai, J.-M. Ratten, J. Zorz, B. M. Robicheau, D. Haider, and J. LaRoche. 2022. Seasonal bacterial niche structures and Chemolithoautotrophic ecotypes in a North Atlantic Fjord. *Sci. Rep.* **12**: 15335. doi:10.1038/s41598-022-19165-w
- Ruiz-González, C., R. Simó, R. Sommaruga, and J. M. Gasol. 2013. Away from darkness: A review on the effects of solar radiation on heterotrophic bacterioplankton activity. *Front. Microbiol.* **4**: 131. doi:10.3389/fmicb.2013.00131
- Samuel, R. M., and others. 2021. Toward a global public repository of community protocols to encourage best practices in biomolecular ocean observing and research. *Front. Mar. Sci.* **8**: 758694. doi:10.3389/fmars.2021.758694
- Sogin, M. L., H. G. Morrison, J. A. Huber, D. M. Welch, S. M. Huse, P. R. Neal, J. M. Arrieta, and G. J. Herndl. 2006. Microbial diversity in the deep sea and the underexplored 'rare biosphere. *Proc. Natl. Acad. Sci. USA* **103**: 12115–12120. doi:10.1073/pnas.0605127103
- Soriano-Lerma, A., V. Pérez-Carrasco, M. Sánchez-Marañón, M. Ortiz-González, V. Sánchez-Martín, J. Gijón, J. M. Navarro-Mari, J. A. García-Salcedo, and M. Soriano. 2020. Influence of 16S rRNA target region on the outcome of microbiome studies in soil and saliva samples. *Sci. Rep.* **10**: 13637. doi:10.1038/s41598-020-70141-8
- Sunagawa, S., and others. 2020. Tara oceans: Towards global ocean ecosystems biology. *Nat. Rev. Microbiol.* **18**: 428–445. doi:10.1038/s41579-020-0364-5
- Thompson, L. R., and others. 2017. A communal catalogue reveals Earth's multiscale microbial diversity. *Nature* **551**: 457–463. doi:10.1038/nature24621
- Tilman, D., F. Isbell, and J. M. Cowles. 2014. Biodiversity and ecosystem functioning. *Annu. Rev. Ecol. Evol. Syst.* **45**: 471–493. doi:10.1146/annurev-ecolsys-120213-091917
- Traving, S. J., and others. 2021. Prokaryotic responses to a warm temperature anomaly in northeast subarctic Pacific waters. *Commun. Biol.* **4**: 1–12. doi:10.1038/s42003-021-02731-9
- Vallina, S. M., M. J. Follows, S. Dutkiewicz, J. M. Montoya, P. Cermenou, and M. Loreau. 2014. Global relationship between phytoplankton diversity and productivity in the ocean. *Nat. Commun.* **5**: 4299. doi:10.1038/ncomms5299
- Wagg, C., K. Schläeppli, S. Banerjee, E. E. Kuramae, and M. G. A. van der Heijden. 2019. Fungal-bacterial diversity and microbiome complexity predict ecosystem functioning. *Nat. Commun.* **10**: 4841. doi:10.1038/s41467-019-12798-y
- Weisburg, W. G., S. M. Barns, D. A. Pelletier, and D. J. Lane. 1991. 16S ribosomal DNA amplification for phylogenetic study. *J. Bacteriol.* **173**: 697–703. doi:10.1128/jb.173.2.697-703.1991
- Wickham, H., and others. 2019. Welcome to the tidyverse. *J. Open Sour. Softw.* **4**: 1686. doi:10.21105/joss.01686
- Wietz, M., C. Bienhold, K. Metfies, S. Torres-Valdés, W.-J. von Appen, I. Salter, and A. Boetius. 2021. The polar shift: Seasonal dynamics and drivers of Arctic Ocean microbiomes revealed by autonomous sampling. *ISME Commun.* **1**: 1–12. doi:10.1038/s43705-021-00074-4
- Willis, C., D. Desai, and J. LaRoche. 2019. Influence of 16S rRNA variable region on perceived diversity of marine microbial communities of the northern North Atlantic. *FEMS Microbiol. Lett.* **366**: fnz152. doi:10.1093/femsle/fnz152

- Wiltshire, K. H., and others. 2010. Helgoland roads, North Sea: 45 years of change. *Estuar. Coasts* **33**: 295–310. doi:[10.1007/s12237-009-9228-y](https://doi.org/10.1007/s12237-009-9228-y)
- Wood, S. N. 2017. Generalized additive models: An introduction with R. Chapman and Hall/CRC. doi:[10.1201/9781315370279](https://doi.org/10.1201/9781315370279)
- Yeh, Y.-C., and J. A. Fuhrman. 2022a. Contrasting diversity patterns of prokaryotes and Protists over time and depth at the San-Pedro Ocean time series. *ISME Commun.* **2**: 1–12. doi:[10.1038/s43705-022-00121-8](https://doi.org/10.1038/s43705-022-00121-8)
- Yeh, Y.-C., and J. A. Fuhrman. 2022b. Effects of phytoplankton, viral communities, and warming on free-living and particle-associated marine prokaryotic community structure. *Nat. Commun.* **13**: 7905. doi:[10.1038/s41467-022-35551-4](https://doi.org/10.1038/s41467-022-35551-4)

Acknowledgments

We acknowledge the Mi'kmaq First Nation people and the indigenous name Kwipek (Head of the Tide) of the Bedford area. We pay respect to the traditional and original owners of Australia, and specifically the Noongar people from Rottnest Island, the Puthikwilayti people from Maria Island and the Biri language group from Northern Queensland (Yongala). This work was funded by NSERC Discovery and OFI grants. Additionally, EJR was supported by an International Postdoctoral Scholarship from the Ocean Frontier Institute. We thank Bill Li, Emmanuel Devred, Lindsay Bailey, the Bedford Institute of Oceanography, and the Department for Fisheries and Ocean for initiating and maintaining the time series and the CERC-OCEAN team for providing the nutrient analyses. Samples from Fram Strait were obtained in the framework of the Helmholtz Infrastructure Program FRAM, funded by the Alfred Wegener Institute Helmholtz Centre

for Polar and Marine Research, with major contributions to conceptualization and coordination from Antje Boetius, Ian Salter, and Katja Metfies. KT and PJS acknowledge funding support from the UK Natural Environment Research Council (NERC), originally through Oceans2025 Theme 4, and more recently through its National Capability Long-term Single Centre Science Programme, Climate Linked Atlantic Sector Science (Grant no. NE/R015953/1). We acknowledge the contribution of the Australian Microbiome consortium in the generation of data used in this publication. The Australian Microbiome initiative is supported by funding from Bioplatforms Australia and the Integrated Marine Observing System (IMOS) through the Australian Government's National Collaborative Research Infrastructure Strategy (NCRIS), Parks Australia through the Bush Blitz program funded by the Australian Government and BHP, and the CSIRO. We thank the people running the BBMO site and the PIs of the projects that have funded research at the site during the years, particularly I. Ferrera and E. Garcés for the sequencing here analyzed. The ICM authors acknowledge the "Severo Ochoa Centre of Excellence" accreditation (CEX2019-000928-S). We also thank the anonymous reviewers for their time and constructive feedback and the valuable editorial suggestions and comments during proofreading from Dr. R. M. Franco-Santos. We thank Dr. Jahangir Vajedsamiei for his kind advice addressing autocorrelation issues.

Conflict of Interest

None declared.

Submitted 26 May 2023

Revised 22 May 2024

Accepted 14 June 2024