

EGU24-12738, updated on 15 Apr 2025

<https://doi.org/10.5194/egusphere-egu24-12738>

EGU General Assembly 2024

© Author(s) 2025. This work is distributed under the Creative Commons Attribution 4.0 License.



Evaluating the PolarRES regional models using tailor-made climate indices for Arctic reindeer herding communities

Heidrun Matthes¹, Jussi T Eronen⁴, Xavier Fettweis⁵, Bruce C Forbes³, Ella Gilbert⁶, Joachim Otto Habeck², Tim Horstkotte⁹, Kirill Istomin², Teresa Komu³, Oskar Landgren¹², Jan Landwehrs¹, Roza Laptander², Priscilla Mooney⁸, Ruth Mottram¹⁰, Christiaan van Dalum¹¹, Willem Jan van den Berg¹¹, Sirpa Rasmus³, Annette Rinke¹, and Hans Tømmervik⁷

¹Alfred Wegener Institut Helmholtz-Zentrum für Polar-und Meeresforschung, Potsdam, Germany

²Universität Hamburg, Hamburg, Germany

³Lapin Yliopisto, Rovaniemi, Finland

⁴Helsingin Yliopisto, Helsinki, Finland

⁵Université de Liège, Liège, Belgium

⁶British Antarctic Survey, Cambridge, UK

⁷Norsk institutt for naturforskning, Tromsø, Norway

⁸Norwegian Research Centre, Bergen, Norway

⁹Umeå universitet, Umeå, Sweden

¹⁰Danmarks Meteorologiske Institut, København, Denmark

¹¹Universiteit Utrecht, Utrecht, Netherlands

¹²Meteorologisk institutt, Oslo, Norway

From a combination of the operational system of reindeer herding and meteorological seasonality, we developed a range of climate indices reflecting critical events in the reindeer herding year that influence the success of this livelihood. These critical events can be described as combinations of specific meteorological conditions, and therefor rendered as equations we can compute from climate model output, creating a capability for analysing different projections of the future and delivering relevant information on climate change to reindeer herding communities.

For this purpose, we can use a wealth of different global and regional climate projections, with distinct advantages and disadvantages (e.g. model resolution, different greenhouse gas futures, high number of models in the ensemble for uncertainty estimates, process representation, availability of variables). For example, the CMIP6 ensemble enables the analysis of a broad range of greenhouse gas futures from a wide variety of models, allowing us to assess scenario uncertainty, but it is limited by its coarse spatial resolution. On the other hand, the PolarRES ensemble has a higher spatial resolution but is only available for one RCP/greenhouse gas future. The PolarRES ensemble consists of regional climate simulations generated by multiple regional climate models that dynamically downscale CMIP6 global climate simulations selected using a novel storyline approach. Both ensembles provide hindcast simulations that allow us to evaluate the ensemble performance with regard to the climate indices we defined.

This study uses these simulations to evaluate and compare model performance to understand the potential and limitations of future projections of specific climate indices relevant for reindeer herding. We use in-situ based observations from the data set Global Summary of the Day to evaluate onset and end of the continuous freezing period, hot summer days, thawing days in autumn and freeze-thaw cycles in both spring and autumn.