



ICES
CIEM

International Council for
the Exploration of the Sea

Conseil International pour
l'Exploration de la Mer

Pathways for converting zooplankton traits to ecological insights are paved with findable, accessible, interoperable, and reusable (FAIR) data practices

Jessica Titocci ^{1,*}, Patrick R. Pata ^{2,3,*}, Tiziana Durazzano ⁴, Sakina-Dorothee Ayata ^{5,6}, Corentin Clerc ^{5,7}, Astrid Cornils ⁸, Patrick Duffy ⁹, Adam T. Greer ⁹, Claudia Halsband ¹⁰, Ryan F. Heneghan ¹¹, Anaïs Lacoursière-Roussel ¹², Fabien Lombard ^{6,13}, Sanna Majaneva ¹⁴, Evgeny A. Pakhomov ^{15,2,3}, Carolina Reis ^{15,16}, Sinja Rist ¹⁷, Alix C.M. Rommel ¹⁸, Teresa Silva ¹⁹, Lars Stemmann ^{15,20}, Kevin Ugwu ²¹, Alberto Basset ²², Ilaria Rosati ¹, Kieran J. Murphy ²³, Brian P.V. Hunt ^{2,3}

¹National Research Council (CNR), Research Institute for Terrestrial Ecosystems (IRET), Lecce, 73100, Italy

²Institute for the Oceans and Fisheries, University of British Columbia, Vancouver, BC, V6T 1Z4, Canada

³Department of Earth, Ocean and Atmospheric Sciences, University of British Columbia, Vancouver, BC, V6T 1Z4, Canada

⁴Department of Arctic and Marine Biology, UiT The Arctic University of Norway, Tromsø, 9037, Norway

⁵Sorbonne Université, CNRS, IRD, MNHN, Laboratoire d'Océanographie et du Climat: Expérimentations et Approches Numériques, LOCEAN-IPSL, Paris, 75005, France

⁶Institut universitaire de France (IUF), Paris, 75005, France

⁷Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich, Zürich, 8092, Switzerland

⁸Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, 27570, Germany

⁹Skidaway Institute of Oceanography, University of Georgia, Savannah, GA, 31411, United States

¹⁰Akvaplan-niva, Fram Centre, Tromsø, 9007, Norway

¹¹Australian Rivers Institute, School of Environment and Science, Griffith University, Nathan, QLD 4111, Australia

¹²St. Andrews Biological Station, Fisheries and Oceans Canada, St. Andrews, New Brunswick, E5B 0E4, Canada

¹³Sorbonne Université, Laboratoire d'Océanographie de Villefranche, UMR 7093, CNRS, Institut de la Mer de Villefranche, Villefranche sur mer, 06230, France

¹⁴Akvaplan-niva, Trondheim, 7010, Norway

¹⁵Marine Biotechnology Post-Graduation Program (PPGBM), IEAPM/UFF, Niterói, Rio de Janeiro, 24020-141, Brazil

¹⁶Flanders Marine Institute (VLIZ), Jacobsenstraat 1, 8400 Ostend, Belgium

¹⁷National Institute of Aquatic Resources (DTU Aqua), Technical University of Denmark, Henrik Dams Allé, 2800 Kgs. Lyngby, Denmark

¹⁸Scottish Oceans Institute, University of St Andrews, Gatty Marine Laboratory, St Andrews KY16 8LB, United Kingdom,

¹⁹Marine and Freshwater Research Institute (MFRI), Fornubúðir 5, 220 Hafnafjörður, Iceland

²⁰Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche (LOV), Villefranche-sur-Mer, 06230, France

²¹Man–Technology–Environment Research Centre (MTM), Örebro University, 70182, Örebro, Sweden

²²Department of Biological and Environmental Sciences and Technologies (DiSTeBA), University of Salento, Lecce, 73100, Italy

²³The Australian Centre for Excellence in Antarctic Science, University of Tasmania, Hobart, 7001, Australia

*Corresponding authors. Jessica Titocci, National Research Council (CNR), Research Institute for Terrestrial Ecosystems (IRET), Lecce, 73100, Italy. E-mail: jessica.titocci@cnr.it; Patrick Pata, Institute for the Oceans and Fisheries, University of British Columbia, Vancouver, BC, V6T 1Z4, Canada. E-mail: p.pata@oceans.ubc.ca.

[†]These two authors contributed equally to this work.

Abstract

The use of trait-based approaches and trait data in zooplankton ecology is rapidly growing to better understand and predict the patterns of zooplankton distributions and their role in aquatic ecosystems and biogeochemical cycles. Although the number of zooplankton trait-based studies and available trait datasets is increasing, several challenges remain for the findability, accessibility, interoperability, and reusability (FAIR) in trait-based approaches that, if unaddressed, may stifle progress in this research area. Here, we review recent applications of trait-based approaches in zooplankton research and summarize the currently available trait data resources. To realize the potential of trait-based approaches to resolve ecological roles of zooplankton, datasets and approaches must adhere to FAIR principles. We provide recommendations and pathways forward to ensure FAIRness while highlighting the importance of collaborative efforts. These practical and easily implementable strategies will enhance the FAIRness of trait data, ultimately advancing zooplankton ecological research and connecting these findings to aquatic ecosystem functioning.

Keywords: trait; zooplankton; FAIR; trait-based approaches




Figure 1. Characteristics of trait-based approaches in zooplankton ecology listed as advantages, challenges, and methodological or ecological applications. Zooplankton vectors adapted from Pata, P. (2023). Marine organism line art. <https://doi.org/10.5281/zenodo.13685461>.

Table 1. Examples of applications of trait-based approaches in zooplankton ecology categorized thematically based on the study topic.

Category	References
Individual-level analysis, trade-offs between traits	Kjørboe (2008), Gorokhova et al. (2013), Bianco et al. (2014), Kjørboe and Hirst (2014), Kjørboe et al. (2015)
Population-level analysis	García et al. (2007), Frances et al. (2021), Ilić et al. (2021), Shaw et al. (2021)
Community-level analysis	Litchman et al. (2013), Pomerleau et al. (2015), Romagnan et al. (2015), Benedetti et al. (2016, 2023), Hébert et al. (2016b), Teuber et al. (2019), Becker et al. (2021), Vilgrain et al. (2021), Cornils et al. (2022), Feuilloley et al. (2022), Li et al. (2022), Beck et al. (2023), Perhirin et al. (2024)
Biodiversity and ecosystem functioning	Barnett et al. (2007), Hébert et al. (2016a), St.-Gelais et al. (2023)
Modeling biogeography	Barton et al. (2013), Brun et al. (2016), Prowe et al. (2019), Drago et al. (2022)
Modelling food webs, including size-based ecosystem models	Heneghan et al. (2016), Prowe et al. (2019, 2020, 2023), Everett et al. (2022), Negrete-García et al. (2022), Clerc et al. (2023)
Modeling biogeochemistry	Renaud et al. (2018), Archibald et al. (2019), Luo et al. (2020), Wright et al. (2021), Serra-Pompei et al. (2022), Pinti et al. (2023), Clerc et al. (2024)

This table is an open access source that can be updated regularly, and we invite all zooplankton researchers to contribute to its future version at this link <https://osf.io/gqu53>.

such as GenBank, Barcode of Life Data Systems (BOLD, doi.org/10.17616/R3PP7J), and MetaZooGene Atlas and Database (Bucklin et al. 2021, O'Brien et al. 2024) provide an additional source of taxonomic data that can be integrated with trait data and facilitate interspecies comparisons and the linkage of genetic information with phenotypic traits. Another emerging trait data resource is Ecotaxa (<https://ecotaxa.obs-vlfr.fr/>), the largest database of sorted plankton images containing ~450 million images, which is dedicated to the taxonomic identification of images of plankton and includes measurements of morphological traits (Picheral et al. 2017).

Collectively, these resources support the advancement of trait-based research in zooplankton ecology by providing readily accessible information, courses, and documentation of FAIR data practices to researchers worldwide. The repositories also have implemented quality assurance (QA) and quality control (QC) measures to maintain high standards of data submission. Users collaborate through defined responsibilities shared among data providers, data curators, and reviewers, and provide direct support and guidance to facilitate data preparation and submission. This ensures rigorous QC through evaluation processes that may include pre-entry checks, formal criteria assessments, and content review. However, trait data resources remain decentralized and widely dispersed across the Web, which makes it challenging to develop robust information systems to aggregate and disseminate data according to FAIR principles. To date, there has been no specific online repository dedicated to zooplankton traits. Nevertheless, the Open Traits Network (OTN) represents one of the major efforts to provide a centralized hub for trait-related data (<https://opentraits.org>). It includes a collection of trait datasets, often specific to a particular taxonomic domain or biogeographic region, with some datasets including zooplankton taxa. Similarly, a catalogue of trait databases for a plethora of aquatic organisms, including zooplankton, has been provided in Martini et al. (2021).

As more trait data become available, collaborative projects and initiatives are establishing standardized methods for producing FAIR trait data (Gallagher et al. 2015, Schneider et al. 2019, Keller et al. 2023). In recent years, particular attention has been paid to addressing issues of heterogeneity in terminologies of trait data in units or categorical variables by applying standardized definitions. One of the most important examples of improving the annotation, standardization, and interoperability of trait data and metadata is the Ecological Trait-data Standard (ETS) vocabulary (Schneider et al. 2019). The ETS vocabulary is a single resource terminology that provides a starting point for the development of a common language and terminology around traits and trait-based research across disciplines. Specifically, for zooplankton, the Zooplankton Trait Thesaurus (<https://ecoportal.lifewatch.eu/ontologies/ZOOPLANKTRAITS>) was initiated to make zooplankton-related trait terminologies machine-readable and actionable, to ensure data consistency, and to promote zooplankton data harmonization and integration across studies. This resource provides standards mainly for morphological traits and is currently being improved and extended to include standards for physiological, behavioural, and life-history traits. Furthermore, this thesaurus has been merged with the Phytoplankton Trait Thesaurus (Rosati et al. 2017), the Macroalgae Trait Thesaurus, and the Fish Trait Thesaurus in a unique resource: the Traits Thesaurus (<https://www.doi.org/10.48373/sa6p-ta25>).

The ultimate goal of this Traits Thesaurus is to interact with the ETS and to be linked and aligned with other existing biodiversity and trait data terminology initiatives within The Biodiversity Information Standards (<https://www.tdwg.org/>). Establishing clear trait definitions and semantically annotating trait terminologies with rich metadata using controlled vocabularies is essential for standardizing traits and associated information, such as units of measurement, the location or environment where the trait was measured, the level of measurement (e.g. individual or species), and the protocol or instrument used. This limits confusion during data aggregation and allows for accommodating multiple trait records for a single species, which facilitates documenting intraspecific variation.

Recently, the identification of zooplankton and the collection of trait-based measurements have seen significant advances through the integration of image processing pipelines, machine learning applications, and various analytical tools, Web services, and open-source code and software (Supplementary Table). Among these, Ecotaxa is a valuable example of an interactive database that allows collaborative image processing and classification according to a universal taxonomy and machine learning techniques to automate the identification process, thereby reducing manual effort and improving data consistency. Ecotaxa uses the UniEuk taxonomic framework, which is based on curated molecular phylogenies and has established new standards relevant to marine biodiversity image networks (Irisson et al. 2022, Martin-Cabrera et al. 2022). Other openly available analytical services for zooplankton trait data acquisition, processing, analysis, and modelling include The LifeWatch Data Explorer, Plankton Toolbox, Plankton Identifier, Plankton Lifeform Extraction Tool, and Plankton Inversion Model (Supplementary Table). The Zoo and Phytoplankton EOY Product from the Blue-Cloud VRE and the Plankton Genomics VLab are examples of Virtual Lab (VLab) and VREs that promote e-Science and collaborative research (Supplementary Table). They do so through platforms that provide integrated access to data, computational resources, and analytical workflows for interpolating sparse *in situ* measurements and modelling phytoplankton-zooplankton interactions, and for in-depth assessment of plankton distributions by mining biomolecular, imaging, and environmental data. All of these data resources and Web services have the potential to greatly motivate researchers to organize, share, and use trait data, fostering the adoption of common protocols and analytical processes in accordance with the FAIR principles.

Existing challenges in zooplankton trait-based approaches

Despite the growing interest in trait-based approaches in zooplankton ecology, their application in fundamental and generalizable ecological research is limited by (i) data availability, (ii) data FAIRness, and (iii) some inherent shortcomings and uncertainties in recent implementations of trait-based approaches.

Trait data availability

There are significant gaps in the taxonomic coverage of zooplankton trait information where most available data are extracted from morphological information (e.g. size, shape, body mass) or categorical behavioural traits, with less avail-

able information on physiological, biochemical, and life-history traits (Pata and Hunt 2023). The existing databases mainly focus on crustaceans, with other taxa, such as pelagic cnidarians, ctenophores, and tunicates, largely absent. Important zooplankton groups that need to be incorporated into or linked with global zooplankton trait databases include meroplankton, microzooplankton, macrozooplankton, and freshwater species. Additionally, several important traits, such as growth rates, clearance rates, reproduction frequency, vertical migration, motility, and size at maturation, are largely missing from the available databases even for relatively well-studied species.

A strategy for resolving the gaps in trait data availability is to use available data as proxies for the traits of interest or estimating trait values through broad taxonomic generalizations, allometric scaling equations, or imputation (Litchman *et al.* 2021, Thorson *et al.* 2023). Standardization of these estimation procedures is necessary, with recommendations provided by de Bello *et al.* (2021). The accuracy of these various methods in estimating zooplankton traits was found to be strongly dependent on the number of existing trait records (Pata and Hunt 2023). Thus, acquiring new trait data is essential.

It is not clear whether the limitations on zooplankton trait data's findability and accessibility stem from actual gaps in data availability or from the scarcity of FAIR zooplankton trait data. At the same time, zooplankton trait data that are not available in public open data repositories but are stored in log books or on computer hard drives and discs in research laboratories need to be identified, digitized, and uploaded to appropriate data repositories. Furthermore, trait data can also be 'hidden' in publications that collect these data for other purposes and are therefore not incorporated in trait databases. However, mining for all the hidden data would require extensive manual effort to harmonize the datasets and integrate them into trait data repositories.

Trait data FAIRness

Beyond the challenges in data availability, the majority of available zooplankton trait data are decentralized, and, currently, there is no single access point or repository specifically tailored for accessing zooplankton trait data and metadata. As a result, many available data exist primarily as trait datasets attached to publications or uploaded to general-purpose data repositories such as Figshare, Zenodo, ResearchGate, and Data Dryad. Although these general-purpose repositories are open source and provide DOIs, they typically allow data to be archived with minimal standards for metadata documentation and data interoperability, resulting in variable tabular structures and labelling conventions for trait variables. Therefore, they often do not fully meet the requirements of FAIR principles. Beyond general-purpose data repositories, zooplankton trait data and metadata are also often scattered across multiple domain-specific repositories for marine and freshwater systems (Supplementary Table) or available through institutional websites and e-Science research infrastructure (e.g. LifeWatch, Long Term Ecological Research Network, European Marine Biological Resource Centre, European Multidisciplinary Seafloor and water column Observatory). In addition, FAIR data practices and data quality assurance measurements are still not consistent across research data hosting centres and vary considerably depending on the repository

used (Kindling and Strecker 2022). Thus, the scattered data are complicated to extract and subsequently reuse. This issue is exacerbated by the occurrence of multiple parallel and concurrent efforts that may share the same objectives but employ their own approaches to data FAIRness and data and metadata quality assurance, resulting in differences in the metadata schemas, data formats, standards used, and individual approaches to QA/QC assessment. Currently, the lack of common agreement, linkage, and interoperability among these data resources underscores the urgent need for alignment and harmonization. Moreover, similar to the peer review of research papers, research on the assessment and certification of data repositories is needed to improve the value of data, policies, procedures, and services for data management and sharing.

Another source of misunderstanding that limits data FAIRness is, fundamentally, in the definition of what a trait is, which previous studies have attempted to constrain (Violle *et al.* 2007). Trait-based approaches include a heterogeneity in the sources of trait observations and in the level of biological organization in which traits were measured. Thus, practically, ecologists use varying, although closely related, definitions of what a trait is (Dawson *et al.* 2021). Beyond resolving the term 'trait', the terminologies and units of the traits themselves are context-dependent and remain challenging to standardize and decode.

In recent years, the DarwinCore (DwC) standard (Wieczorek *et al.* 2012) and DwC-Archive have gained popularity for biodiversity data. However, the recommended data architecture was created with the intention of harmonizing occurrence data and does not provide standards for trait data. The 'Ecological Trait-data Standard' (ETS; Schneider *et al.* 2019) addresses this gap, providing a vocabulary for trait datasets with essential terms such as 'trait name', 'trait value', and 'trait units'. The ETS, however, does not offer standardized descriptions for trait names and units, which can be domain specific. There are currently a number of glossaries aimed at standardizing terminology and facilitating communication between trait-based research communities (e.g. Marine Species Traits, BIOTIC, ICES vocabulary, Ecotaxonomy). However, a common problem is that many of these glossaries often lack the ability to create linked data, as they do not provide Uniform Resource Identifiers (URIs) for terms that are required to connect between different datasets and resources (Parr *et al.* 2016, Schneider *et al.* 2019). This limitation in interoperability hinders their usefulness in automated systems and data integration efforts and is further exacerbated by the lack of alignment between the semantic resources mentioned previously.

Another key challenge when integrating and ensuring the FAIRness of zooplankton trait data lies in the quality of the metadata associated with the collected trait data, which can be widely heterogeneous (e.g. various levels of taxonomic specificity, non-systematic information about the location and time of sampling). Due to the rapid changes in marine environments caused by climate change, harmonized metadata are needed to evaluate and establish trait baselines, as well as to track the changes over time and to test their robustness. In particular, the recurrent absence of spatial metadata for zooplankton traits poses difficulties for understanding ecological and biogeographical patterns and limits the ability to relate trait variations with environmental conditions.

Trait-based approaches

Trait selection is limited by the amount of available information from literature and the cost of making new trait measurements. Reliance on analysing only the functional traits that are deemed to have sufficiently available information introduces a risk of oversimplification that may fail to capture the functional diversity or ecological process of interest. One outcome of this is focusing mainly on some easily measurable traits such as size, shape, and body mass (or biovolume), and putting less emphasis on behavioural or physiological traits, which may require more detailed analyses and more specific equipment to obtain accurate and reliable trait data (Petchey and Gaston 2006). To address this, we recommend identifying the relevant traits *a priori* and explicitly linking these to the associated organismal or ecosystem processes of interest. This approach is consistent with the work already undertaken within the Essential Ocean Variables (EOVs) and the Essential Biodiversity Variables (EBVs), which provide international co-ordination on best practices for observing and producing data (Muller-Karger et al. 2018). The EOV and EBV frameworks demonstrate the value of recommended and commonly agreed standards and protocols for data collection and management at the international level to ensure interoperability and integrity of data from local to global scales. Furthermore, since trait information could be derived from various levels of biological organization, focusing only on taxonomic level trait values (e.g. using the same trait value for all individuals of the same species) will effectively miss the intrataxa variability. This presents a particular challenge when trait plasticity is ecologically relevant, whether from developmental changes in life history, regional variability in environmental conditions, or from the long-term responses to environmental changes. The appropriateness of trait records obtained from data compilations needs to be evaluated based on the ecological context for when the trait was measured, which are ideally provided by the metadata. This would be consistent with FAIR data practices necessary for utilizing trait data to address ecological questions.

The selection of measured or applied traits can affect estimates and applications of trait data (de Bello et al. 2021). For practical reasons, researchers tend to focus on understanding the ecosystem effects of a single or a few traits. The use of a ‘representative trait’ approach, often involving a single trait (i.e. size), may oversimplify the interpretation of these ecological aspects. For instance, recent work documenting morphological traits of mesozooplankton revealed that carbon export may be more influenced by body transparency (i.e. more transparent individuals are often gelatinous) than by their size (Perhirin et al. 2024). Instead, measuring more traits may better represent the functional diversity and variation of the community (Maire et al. 2015), although measuring a large number of traits at different levels of ecological complexity quickly becomes impractical, requiring increased sampling effort and costs for analysis. Traits are, however, known to intersect in the organismal processes they capture and so identifying the commonalities between traits in the statistical trait space (Vilgrain et al. 2021) would be useful for trait selection.

Currently, the relationships between traits and associated ecological functions are poorly investigated and understood (Degen et al. 2018). Furthermore, the trade-offs between resource acquisition and defence govern the diversity of communities, as they allow the coexistence of many trait configura-

tions with similar fitness (Hébert et al. 2017). The execution of any one of these functions, however, may conflict with the others, as they cannot all be maximized simultaneously (Bremner et al. 2006). Thus, quantifying the risks and trade-offs associated with key traits is necessary to predict the morphology, behaviour, and physiology that optimizes the fitness of an organism in any environment (Violle et al. 2007), which requires cross-disciplinary investigations.

Finally, an important challenge to address in zooplankton ecology is understanding how zooplankton traits and processes contribute to and interact within the overall food web, from virioplankton to large marine mammals. Other than body size, there are traits that are common across trophic groups (Litchman et al. 2021) and the use of these cross-trophic level traits will stimulate trait-based research on entire ecosystems (Martini et al. 2021). Moreover, it will be worthwhile to learn from other ecological domains, such as terrestrial plants, that may have led to the development of trait-based approaches in terms of trait data infrastructure, forming expert networks, and applying traits for predictive studies (Green et al. 2022).

Future directions

Although trait-based approaches have been used in ecological research for some time, coordinated efforts in trait data collection, harmonization, and analysis are needed to realize the full potential of zooplankton trait-based approaches. Broadly, future zooplankton trait-based research involves the need to incorporate new sources of trait data and facilitate how trait-based studies utilize common standards, practices, and technologies for FAIR data to achieve a more cohesive and comprehensive understanding of zooplankton trait-based knowledge. We believe that several steps are necessary to coordinate and promote the collection and standardization of zooplankton trait data and the increase of FAIR zooplankton trait data (Fig. 2).

Enabling zooplankton trait data findability and accessibility

Continuing to collect zooplankton trait data from field and laboratory studies, as well as finding and synthesizing trait records from literature into existing databases, is an ongoing effort that will advance trait-based approaches. This process involves digitizing existing data and metadata, extracting information from papers and grey literature, and encouraging potential data holders to standardize and share their data in domain-specific repositories through persistent identifiers (PIDs). It is essential to establish structured connections between zooplankton data and related information through high-quality and rich metadata. This includes detailed descriptions of the methodology, environmental conditions of the trait measurements, data provenance, and other relevant attributes. For image data, assigning a unique identifier to high-resolution images will allow for potential revisions or additions to the trait measurements. When publishing, it is crucial to share data and metadata openly through domain-specific repositories rather than in supplementary materials, ensuring that the data and metadata have a PID. Additionally, the published data need to be associated with a clear usage licence. The high volume and diversity of trait data generated by different systems and research groups underscore the critical




Figure 2. Summary of next steps to achieve FAIRness in zooplankton trait-based research. Recommendations are listed thematically for next steps related to FAIR data practices, collaboration within the scientific community, methods in trait data analysis, and expanding trait data collection. Sources: Illustration of the UMCES research vessel, *RV Rachel Carson*, adapted from Tracey Saxby, Integration and Application Network (ian.umces.edu/media-library), CC BY-SA 4.0. Zooplankton vectors adapted from Pata, P. (2023). Marine organism line art. <https://doi.org/10.5281/zenodo.13685461>.

importance of reproducible and platform-independent data management strategies.

To support these efforts, a dedicated data rescue working group should be established, potentially within the ICES framework, to facilitate connections and common practices and to provide technical support in adjusting existing data to fit the structure of databases. The contributions of data providers to making data FAIR and harmonized must be acknowledged, credited, and incentivized, thereby increasing the value of data reuse and promoting even more the adoption of FAIR data management practices. Additionally, centralized

databases such as the Plant Trait Database (TRY, <https://www.try-db.org/de/TabDetails.php>) and FishBase (<https://fishbase.org>) have demonstrated that providing accessible resources in a single access point facilitates innovation in terrestrial plant and fish functional ecology. Since there is currently no specific online repository dedicated to zooplankton traits, the zooplankton research community must decide on a suitable path forward. Options include improving data FAIRness across existing repositories by aligning and standardizing FAIR data management practices along the different repositories or developing a centralized trait database or data portal specific

for zooplankton traits. A centralized platform has the potential to benefit zooplankton ecologists by facilitating the findability, accessibility, and integration of zooplankton trait data into their studies. However, managing a centralized repository is challenging and requires consistent funding to remain operational and ongoing investment in technical digital infrastructure, staff, and user support to maintain their functionality. Collaborations with digital research infrastructure, academic institutions, funding bodies, and data-sharing consortia can help ensure the long-term sustainability of the repository.

Enabling zooplankton trait data interoperability

Even if trait data are available in open data repositories, data interoperability seems to be the most challenging aspect, especially among data generated by different devices and research groups (Bi et al. 2024). A critical first step in adhering to FAIR principles is the need for semantic and syntactic data standardization in zooplankton ecology. This includes standardizing data and metadata from disparate trait data collection systems into interoperable and reusable formats and adopting widely used protocols and manuals for zooplankton trait measurements (e.g. Harris et al. 2000). For this reason, zooplankton ecologists must achieve a general agreement on the definitions of specific traits and how these are measured. This would require developing agreed-upon trait terminologies and trait measurement protocols and the need for more Linked Open Data (LOD) for sharing machine-readable and actionable interlinked data on the Web. This should be achieved by the improved use of existing standards and controlled vocabularies for biodiversity and trait data annotation (Supplementary Table). Although initial steps have been taken to include species traits as EBVs, and several datasheets, templates, and guidelines for trait data collection have been proposed (Muller-Karger et al. 2018), progress is slow for integrating and operationalizing species traits in global monitoring. This is due partly to the challenges highlighted in this paper, such as the lack of a clear consensus on what is considered a ‘trait’ and insufficient standardization of trait data and metadata from data providers. In this context, developing tools to assist in aligning datasets to data templates and in mapping the measured traits to trait-based semantic resources remains a useful and ongoing task. These tools should include software for Web-based access to trait data and semantic Web standards.

Pre-configured spreadsheet templates for capturing trait data may be an efficient way to identify and share a consistent structure and terminology extracted from controlled vocabularies (i.e. thesauri, ontologies) for trait data collection that can be used in workflow analyses through Web services and Virtual Research Environments (VREs). Similar initiatives already exist for occurrence data and phytoplankton trait data (i.e. PhytoplanktonData Template, <https://www.phytovre.lifewatchitaly.eu/phyto-data-template/>). For plankton imaging data management, best practices and recommendations have already been individuated among EMODnet Biology, OBIS, and EurOBIS networks (Martin-Cabrera et al. 2022) and some specific data templates and standard terminologies have been suggested (De Pooter et al. 2017). However, usually, data templates have limitations, as they can be restrictive, inflexible, and may not cover all possible scenarios exhaustively. An alternative approach is to use template generators, such as The Nansen Legacy Template Generator for

DwC and CF-NetCDF (Marsden and Schneider 2024), which enables the addition of terms such as the Climate and Forecasting convention standard names and DwC terms to spreadsheets, allowing scientists to create semantically aware templates without needing to understand the underlying technology. Similar initiatives should be shared in the zooplankton community to provide open source easily built templates with standardized structures for trait data collection and acquisition that is flexible enough to ideally contain the details related to the trait record (e.g. the prey type and concentrations used when measuring feeding rates).

Although the use of data templates may be helpful in trait data harmonization, they have limited applications if data interoperability is weak. Therefore, there is a strong need to identify mappings among already existing trait-based semantic resources and align these resources. In this context, the use of semantic technologies that promote automated ontology matching approaches is a promising solution to the semantic heterogeneity problem. Moreover, trait-based terminologies included in glossaries and vocabularies with no URIs should be assigned with URIs to facilitate the sharing and reuse of LOD on the Web. Thus, in the case of zooplankton data, providing the specific standardized definitions of traits and units through the Traits Thesaurus will enable an automated tool to extract relevant parameters from these diverse data sources. It is essential that each type of measurement is uniquely identifiable and accessible in a machine-readable format. The list of possible traits is extensive, yet many are seldom measured or recorded as continuous data, but rather as categorical descriptions. With regard to categorical traits, each value or level of a trait would require standardized definitions as well. As a result, there is an increasing demand for domain-specific thesauri and ontologies tailored to address these limitations and improve annotation and specificity, ensuring machine-readability and actionability. These actions, aligned with preexisting government-backed frameworks and product distribution initiatives, should result as synergistic efforts for enhancing the information available for existing EBVs. This would also facilitate inclusion of species traits as EBVs, particularly within the zooplankton domain.

Enabling zooplankton trait data reusability

Once zooplankton trait data are findable, accessible, and interoperable, it is essential to enhance their full reusability by documenting the analytical workflow. This will ensure that zooplankton trait data are not only accessible but also reusable in different analytical contexts, thus maximizing the value of the data collected. Since the handling and processing of large amounts of zooplankton trait data is difficult to perform manually, the development of efficient tools for data integration, analysis, and modelling, open-source codes, user-friendly Web services, workflows, and VREs specifically designed for zooplankton trait-based data analysis should be strongly encouraged by digital research infrastructure and IT developers. Reusability also includes the appropriate recognition of the data providers and preserving the history of the PIDs linked to the data. Digital research infrastructure plays an important role in propagating good practices in data use and citation through providing user interfaces and tools that facilitate reusability. In plankton imaging, significant advances in machine learning and computer vision algorithms

have led to an immense increase in recorded data quantity and quality (Orenstein *et al.* 2022). These data are now part of large-scale monitoring programmes, such as the Canadian Aquaculture Monitoring Program (Finnis *et al.* 2023). Efforts are underway to make these images and associated processing algorithms available for broad-scale plankton analyses of taxa and functional traits (Picheral *et al.* 2017, Drago *et al.* 2022, Dugenne *et al.* 2024), with initiatives such as iFDO providing standards and frameworks for FAIR marine images (<https://marine-imaging.com/fair/ifdos/iFDO-overview/>).

The promotion of open-source codes and software, workflows, and pipelines for data processing can support high-throughput data analyses, and various statistical and computational techniques and modelling, including machine learning, and enhance the scalability of research efforts. Dedicated virtual zooplankton labs and VRE should be promoted as collaborative spaces where researchers can work together on zooplankton trait data analysis, sharing tools, resources, and data, facilitating collaboration and knowledge sharing. In addition, incorporating more traits into models allows for multi-platform analyses that comprehensively address trait-based ecological questions and improve predictive power. These models can simulate different ecological scenarios and help predict the effects of environmental changes on zooplankton communities. By implementing these strategies, the zooplankton community can augment the reusability of zooplankton trait data, driving advances in ecological research and supporting sustainable ecosystem management.

Collaborating towards FAIR trait-based zooplankton research

All the steps previously stated depend heavily on support from the institutions and digital research infrastructure that are revolutionizing data management practices across a range of scientific disciplines. They are essential frameworks that facilitate the organization, storage, and dissemination of the vast amounts of research data produced worldwide. However, the effectiveness of this infrastructure depends on the commitment of individual researchers to be open and to improve the FAIRness of their own research, together with the support of data curators and data managers. The existing digital research infrastructure could play an important role in providing funding and professional support in maintaining data centres. In the future, the funding, development, and promotion of a network of aligned physical and digital research infrastructure will be a major asset to the trait-based research community.

Strengthening collaboration and networking in zooplankton research is the basis for integrating diverse expertise, methodologies, and data sources. Stronger networks and collaborative efforts will lead to improved data standardization, and more robust and accurate analytical processes and models. Collaboration will be facilitated through forming working groups to establish expert networks with the skills necessary to navigate the complexities of FAIR data principles. Future working groups would need to solicit financial support specifying the funds and labour necessary for data management.

Encouraging the development of specialized programmes that integrate training in computer science, zooplankton ecology, and data management is also essential. These actions may have the potential to cultivate new professionals who have expertise in both ecological science and data management, fostering dialogue and collaboration and ultimately advancing

research in both fields. Furthermore, implementing workshops and courses related to open science and zooplankton trait data management will facilitate the dissemination of FAIR zooplankton trait data. These initiatives should be promoted early in scientific careers, even at the undergraduate level, and within the framework of digital research infrastructure, which plays a key role in enabling data-driven research by providing digital resources, tools, and Web services. Moreover, connecting with and learning from experts in other domains of ecology, such as marine microbes, fishes, and terrestrial plants, who have more extensive experience in organizing trait data and applying trait-based approaches, would be helpful in acquiring best practices for zooplankton ecology.

Finally, incentives must be generated to foster a culture of data sharing in the scientific community. Improving the way we attribute datasets in peer-reviewed research and academic evaluations is critical. Recognizing and rewarding data sharing efforts through proper citation and academic credit can provide avenues for professional advancement for researchers contributing their data to public repositories. By treating datasets and data papers as research outputs on par with traditional publications, the scientific community can foster a more collaborative and transparent research environment. Future trait data rescue and data standardization activities should consider how to promote the achievements of the scientists generating the original trait data.

The pathways forward are paved with FAIR data practices, and strengthening collaborations among the zooplankton research community with the existing FAIR initiatives and frameworks is much needed. By adhering to FAIR principles and practices, trait-based approaches in zooplankton research hold significant promise for transforming trait data into actionable knowledge leading to a more holistic understanding of the critical role of zooplankton in aquatic ecosystems and biogeochemical cycles. Working towards establishing a consolidated network of trait data providers, stewards, and users will be key in promoting the benefits of FAIR data practices and in achieving the steps we outlined towards converting traits to ecological insights.

Acknowledgements

The authors would like to thank all the participants during the 7th ICES-PICES Zooplankton Production Symposium workshop on ‘Approaches towards findable, accessible, interoperable and reusable (FAIR) zooplankton trait data as stepping stones to improved functional ecology’ for contributing to the vibrant discussions that resulted in the ideas featured in this manuscript. We especially thank Inessa Corney for assisting in organizing the workshop. We also thank Leen Vandepitte and Stefanie Dekeyzer for their feedback on the initial version of the manuscript. The authors thank the three anonymous reviewers for their helpful feedback on the previous version of the manuscript.

Author contributions

J.T. and P.R.P.: conceptualization, writing—original draft, review and editing, project administration, and supervision. T.D.: writing—review and editing and visualization. S-D.A., C.C., A.C., P.D., A.T.G., C.H., R.F.H., A.L-R., F.L., S.M., E.A.P., C.R., S.R., A.C.M.R., T.S., L.S., and K.U.: writing—review and editing. A.B., I.R., K.J.M., B.P.V.H.: conceptualiza-

tion, writing—original draft, review and editing, project administration, and funding acquisition.

Supplementary data

Supplementary data is available at *ICES Journal of Marine Science* online.

Conflict of interest: None declared.

Funding

P.P. was supported by a Natural Sciences and Engineering Research Council of Canada Doctoral Fellowship, MEOPAR grant NCE 2-02-03-038.2, Canadian Space Agency grant FAST 50568, and Brian Hunt's NSERC Discovery Grant RGPIN-2017-04499. J.T. was supported by the project 'Italian Integrated Environmental Research Infrastructure System' (ITINERIS) in the framework of Next Generation EU PNRR—Mission 4 'Education and Research'—Component 2: 'From research to business'—Investment 3.1: 'Fund for the realisation of an integrated system of research and innovation infrastructures', Notice 3264/2021, IR0000032, CUP B53C22002150006. C.R. has received funding from European Union's Horizon 2020 Research and Innovation Project Mission Atlantic (grant # 862428) and by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brasil (CAPES) (Finance Code 001). J.T. and P.P. would also like to acknowledge the support received through the North Pacific Marine Science Organization (PICES) travel grant, which ensured their successful participation at the 7th International Zooplankton Production Symposium in Hobart, Australia.

Data availability

No new data were generated or analysed in support of this research. Information containing the contents of Table 1 and the supplementary table are archived in this Open Science Framework project link (<https://osf.io/gqu53/>).

References

- Archibald KM, Siegel DA, Doney SC. Modeling the impact of zooplankton diel vertical migration on the carbon export flux of the biological pump. *Global Biogeochem Cycles* 2019;33:181–99. <https://doi.org/10.1029/2018GB005983>
- Barnett AJ, Finlay K, Beisner BE. Functional diversity of crustacean zooplankton communities: towards a trait-based classification. *Freshw Biol* 2007;52:796–813. <https://doi.org/10.1111/j.1365-2427.2007.01733.x>
- Barton AD, Pershing AJ, Litchman E *et al.* The biogeography of marine plankton traits. *Ecol Lett* 2013;16:522–34. <https://doi.org/10.1111/ele.12063>
- Beck M, Cailleton C, Guidi L *et al.* Morphological diversity increases with decreasing resources along a zooplankton time series. *Proc R Soc B Biol Sci* 2023;290:20232109. <https://doi.org/10.1098/rspb.2023.2109>
- Becker EC, Mazzocchi MG, Pinto de Macedo-Soares LC *et al.* Latitudinal gradient of copepod functional diversity in the South Atlantic Ocean. *Prog Oceanogr* 2021;199:102710. <https://doi.org/10.1016/j.pcean.2021.102710>
- Benedetti F, Gasparini S, Ayata S-D. Identifying copepod functional groups from species functional traits. *J Plankton Res* 2016;38:159–66. <https://doi.org/10.1093/plankt/fbv096>
- Benedetti F, Wydler J, Vogt M. Copepod functional traits and groups show divergent biogeographies in the global ocean. *J Biogeogr* 2023;50:8–22. <https://doi.org/10.1111/jbi.14512>
- Bi H, Cheng Y, Cheng X *et al.* Taming the data deluge: a novel end-to-end deep learning system for classifying marine biological and environmental images. *Limnol Oceanogr Methods* 2024;22:47–64. <https://doi.org/10.1002/lom3.10591>
- Bianco G, Mariani P, Visser AW *et al.* Analysis of self-overlap reveals trade-offs in plankton swimming trajectories. *J R Soc Interface* 2014;11:20140164. <https://doi.org/10.1098/rsif.2014.0164>
- Biard T, Stemmann L, Picheral M *et al.* In situ imaging reveals the biomass of giant protists in the global ocean. *Nature* 2016; 532:504–7. <https://doi.org/10.1038/nature17652>
- Bremner J, Rogers SI, Frid CLJ. Methods for describing ecological functioning of marine benthic assemblages using biological traits analysis (BTA). *Ecol Indic* 2006;6:609–22. <https://doi.org/10.1016/j.ecolind.2005.08.026>
- Brun P, Payne MR, Kiørboe T. Trait biogeography of marine copepods—an analysis across scales. *Ecol Lett* 2016;19:1403–13. <https://doi.org/10.1111/ele.12688>
- Brun P, Payne MR, Kiørboe T. A trait database for marine copepods. *Earth Syst Sci Data*, 2017;9:99–113. <https://doi.org/10.5194/essd-9-99-2017>
- Bucklin A, Peijnenburg KT, Kosobokova KN *et al.* Toward a global reference database of COI barcodes for marine zooplankton. *Mar Biol* 2021;168:78. <https://doi.org/10.1007/s00227-021-03887-y>
- Butt N, Halpern BS, O'Hara CC *et al.* A trait-based framework for assessing the vulnerability of marine species to human impacts. *Ecosphere* 2022;13:e3919. <https://doi.org/10.1002/ecs2.3919>
- Clerc C, Bopp L, Benedetti F *et al.* Including filter-feeding gelatinous macrozooplankton in a global marine biogeochemical model: model-data comparison and impact on the ocean carbon cycle. *Biogeosciences* 2023;20:869–95. <https://doi.org/10.5194/bg-20-869-2023>
- Clerc C, Bopp L, Benedetti F *et al.* Effects of mesozooplankton growth and reproduction on plankton and organic carbon dynamics in a marine biogeochemical model. *Glob Biogeochem Cycles* 2024;38:e2024GB008153. <https://doi.org/10.1029/2024GB008153>
- Cornils A, Thomisch K, Hase J *et al.* Testing the usefulness of optical data for zooplankton long-term monitoring: taxonomic composition, abundance, biomass, and size spectra from ZooScan image analysis. *Limnol Oceanogr Methods* 2022;20:428–50. <https://doi.org/10.1002/lom3.10495>
- Dawson SK, Carmona CP, González-Suárez M *et al.* The traits of 'trait ecologists': an analysis of the use of trait and functional trait terminology. *Ecol Evol* 2021;11:16434–45. <https://doi.org/10.1002/ece3.8321>
- De Bello F, Carmona CP, Dias A *et al.* *Handbook of Trait-Based Ecology: From Theory to R Tools*. Cambridge: Cambridge University Press, 2021,1–295. <https://doi.org/10.1017/9781108628426>
- Degen R, Aune M, Bluhm B *et al.* Trait-based approaches in rapidly changing ecosystems: a roadmap to the future polar Oceans. *Ecol Indic* 2018; 91:722–36. <https://doi.org/10.1016/j.ecolind.2018.04.050>
- Degen R, Faulwetter S. The Arctic Traits Database—a repository of Arctic benthic invertebrate traits. *Earth Syst Sci Data* 2019;11:301–22. <https://doi.org/10.5194/essd-11-301-2019>
- Deiner K, Bik HM, Mächler E *et al.* Environmental DNA metabarcoding: transforming how we survey animal and plant communities. *Mol Ecol* 2017;26:5872–95. <https://doi.org/10.1111/mec.14350>
- Demetrius L, Ziehe M. Darwinian fitness. *Theor Popul Biol* 2007;72:323–45. <https://doi.org/10.1016/j.tpb.2007.05.004>
- De Pooter D, Appeltans W, Bailly N *et al.* Toward a new data standard for combined marine biological and environmental datasets—expanding OBIS beyond species occurrences. *Biodivers Data J* 2017;5:e10989. <https://doi.org/10.3897/BDJ.5.e10989>

- Drago L, Panaïotis T, Irisson J-O *et al.* Global distribution of zooplankton biomass estimated by in situ imaging and machine learning. *Front Mar Sci* 2022; 9:894372. <https://doi.org/10.3389/fmars.2022.894372>
- Dugenne M, Corrales-Ugalde M, Luo J *et al.* First release of the Pelagic Size Structure database: global datasets of marine size spectra obtained from plankton imaging devices. *Earth Syst Sci Data* 2024;16:2971–99. <https://doi.org/10.5194/essd-16-2971-2024>
- Everett J, Heneghan R, Blanchard J *et al.* Self-organisation of zooplankton communities produces similar food chain lengths throughout the ocean. *Res Sq*. <https://doi.org/10.21203/rs.3.rs-1186379/v1>, 25 April 2022, preprint: not peer reviewed.
- Faulwetter S, Markantonatou V, Pavloudi C *et al.* Polytraits: a database on biological traits of marine polychaetes. *Biodivers Data J* 2014;2:e1024. <https://doi.org/10.3897/BDJ.2.e1024>
- Feuilloley G, Fromentin J-M, Saraux C *et al.* Temporal fluctuations in zooplankton size, abundance, and taxonomic composition since 1995 in the North Western Mediterranean Sea. *ICES J Mar Sci* 2022;79:882–900. <https://doi.org/10.1093/icesjms/fsab190>
- Finnis S, Guyondet T, McKindsey CW *et al.* *Guidance on Sampling Effort to Monitor Mesozooplankton Communities at Canadian Bivalve Aquaculture Sites Using an Optical Imaging System*/by Stephen Finnis [and 14 Others]. St. Andrews, New Brunswick: St. Andrews Biological Station, Fisheries and Oceans Canada, 2023.
- Frances DN, Barber AJ, Tucker CM. Trait-density relationships explain performance in cladoceran zooplankton. *Ecology*, 2021;102:e03294. <https://doi.org/10.1002/ecy.3294>
- Gallagher J, Orcutt J, Simpson P *et al.* Facilitating open exchange of data and information. *Earth Sci Inf* 2015;8:721–39. <https://doi.org/10.1007/s12145-014-0202-2>
- García CE, De Jesús Chaparro-Herrera D, Nandini S *et al.* Life-history strategies of *Brachionus havanaensis* subject to kairomones of vertebrate and invertebrate predators. *Chem Ecol* 2007;23:303–13. <https://doi.org/10.1080/02757540701525988>
- Gleiber MR, Hardy NA, Roote Z *et al.* The Pelagic Species Trait Database, an open data resource to support trait-based ocean research. *Sci Data* 2024;11:2. <https://doi.org/10.1038/s41597-023-02689-9>
- Gorokhova E, Lehtiniemi M, Motwani NH. Trade-offs between predation risk and growth benefits in the Copepod *Eurytemora affinis* with contrasting pigmentation. *PLoS One*, 2013;8:e71385. <https://doi.org/10.1371/journal.pone.0071385>
- Green SJ, Brookson CB, Hardy NA *et al.* Trait-based approaches to global change ecology: moving from description to prediction. *Proc R Soc B Biol Sci* 2022;289:20220071. <https://doi.org/10.1098/rspb.2022.0071>
- Greer AT, Cowen RK, Guigand CM *et al.* Fine-scale planktonic habitat partitioning at a shelf-slope front revealed by a high-resolution imaging system. *J Mar Syst* 2015;142:111–25. <https://doi.org/10.1016/j.jmarsys.2014.10.008>
- Greer AT, Lehrter JC, Binder BM *et al.* High-resolution sampling of a broad marine life size spectrum reveals differing size- and composition-based associations with physical oceanographic structure. *Front Mar Sci* 2020;7:542701. <https://doi.org/10.3389/fmars.2020.542701>
- Harris R, Wiebe P, Lenz J *et al.* *ICES Zooplankton Methodology Manual*. San Diego, California, USA: Academic Press, 2000.
- Hébert M-P, Beisner BE, Maranger R. A meta-analysis of zooplankton functional traits influencing ecosystem function. *Ecology*, 2016a;97:1069–80. <https://doi.org/10.1890/15-1084.1>
- Hébert M-P, Beisner BE, Maranger R. A compilation of quantitative functional traits for marine and freshwater crustacean zooplankton. *Ecology*, 2016b;97:1081–. <https://doi.org/10.1890/15-1275.1>
- Hébert M-P, Beisner BE, Maranger R. Linking zooplankton communities to ecosystem functioning: toward an effect-trait framework. *J Plankton Res* 2017;39:3–12. <https://doi.org/10.1093/plankt/fbw068>
- Heneghan RF, Everett JD, Blanchard JL *et al.* Zooplankton are not fish: improving zooplankton realism in size-spectrum models mediates energy transfer in food webs. *Front Mar Sci* 2016;3:201. <https://doi.org/10.3389/fmars.2016.00201>
- Heneghan RF, Everett JD, Blanchard JL *et al.* Climate-driven zooplankton shifts cause large-scale declines in food quality for fish. *Nat Clim Change* 2023;13:470–7. <https://doi.org/10.1038/s41558-023-01630-7>
- Heneghan RF, Everett JD, Sykes P *et al.* A functional size-spectrum model of the global marine ecosystem that resolves zooplankton composition. *Ecol Modell* 2020;435:109265. <https://doi.org/10.1016/j.ecolmodel.2020.109265>
- Ilić M, Cordellier M, Fink P. Intrapopulation variability in a functional trait: susceptibility of *Daphnia* to limitation by dietary fatty acids. *Freshw Biol* 2021;66:130–41. <https://doi.org/10.1111/fwb.13623>
- Irisson J-O, Ayata S-D, Lindsay DJ *et al.* Machine learning for the study of plankton and marine snow from images. *Ann Rev Mar Sci* 2022;14:277–301. <https://doi.org/10.1146/annurev-marine-041921-013023>
- Keller A, Ankenbrand MJ, Bruelheide H *et al.* Ten (mostly) simple rules to future-proof trait data in ecological and evolutionary sciences. *Methods Ecol Evol* 2023;14:444–58. <https://doi.org/10.1111/2041-210X.14033>
- Kindling M, Strecker D.,. Data quality assurance at research data repositories. *Data Sci J*, 2022;21:18–. <https://doi.org/10.5334/dsj-2022-018>
- Kjørboe T. Optimal swimming strategies in mate-searching pelagic copepods. *Oecologia* 2008;155:179–92. <https://doi.org/10.1007/s00442-007-0893-x>
- Kjørboe T, Ceballos S, Thygesen UH. Interrelations between senescence, life-history traits, and behavior in planktonic copepods. *Ecology* 2015;96:2225–35. <https://doi.org/10.1890/14-2205.1>
- Kjørboe T, Hirst AG. Shifts in mass scaling of respiration, feeding, and growth rates across life-form transitions in marine pelagic organisms. *Am Nat* 2014;183:E118–30. <https://doi.org/10.1086/675241>
- Kjørboe T, Visser A, Andersen KHA. Trait-based approach to ocean ecology. *ICES J Mar Sci* 2018;75:1849–63. <https://doi.org/10.1093/icesjms/fsy090>
- Laakmann S, Auel H. Longitudinal and vertical trends in stable isotope signatures ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of omnivorous and carnivorous copepods across the South Atlantic Ocean. *Mar Biol* 2010;157:463–71. <https://doi.org/10.1007/s00227-009-1332-9>
- Lacoursière-Roussel A, Howland K, Hernandez C *et al.* eDNA metabarcoding as a new surveillance approach for coastal Arctic biodiversity. *Ecol Evol* 2018;8:7763–77. <https://doi.org/10.1002/ece3.4213>
- Li Y, Ge R, Chen H *et al.* Functional diversity and groups of crustacean zooplankton in the southern Yellow Sea. *Ecol Indic* 2022;136:108699. <https://doi.org/10.1016/j.ecolind.2022.108699>
- Litchman E, Edwards KF, Boyd PW. Toward trait-based food webs: universal traits and trait matching in planktonic predator–prey and host–parasite relationships. *Limnol Oceanogr* 2021;66:3857–72. <https://doi.org/10.1002/lno.11924>
- Litchman E, Ohman MD, Kjørboe T. Trait-based approaches to zooplankton communities. *J Plankton Res* 2013;35:473–84. <https://doi.org/10.1093/plankt/fbt019>
- Lombard F, Boss E, Waite AM *et al.* Globally consistent quantitative observations of planktonic ecosystems. *Front Mar Sci* 2019;6:196. <https://doi.org/10.3389/fmars.2019.00196>
- Luo JY, Condon RH, Stock CA *et al.* Gelatinous zooplankton-mediated carbon flows in the global oceans: a data-driven modeling study. *Global Biogeochem Cycles* 2020;34:e2020GB006704. <https://doi.org/10.1029/2020GB006704>
- McManus MA, Greer AT, Timmerman AH *et al.* Characterization of the biological, physical, and chemical properties of a toxic thin layer in a temperate marine system. *Mar Ecol Prog Ser* 2021;678:17–35. <https://doi.org/10.3354/meps13879>

- Madin JS, Anderson KD, Andreassen MH *et al.* The Coral Trait Database, a curated database of trait information for coral species from the global oceans. *Sci Data* 2016;3:160017. <https://doi.org/10.1038/sdata.2016.17>
- Maire E, Grenouillet G, Brosse S *et al.* How many dimensions are needed to accurately assess functional diversity? A pragmatic approach for assessing the quality of functional spaces. *Global Ecol Biogeogr* 2015;24:728–40. <https://doi.org/10.1111/geb.12299>
- Maps F, Storozhenko PP, Świeżewski J *et al.* Automatic estimation of lipid content from in situ images of Arctic copepods using machine learning. *J Plankton Res* 2024;46:41–7. <https://doi.org/10.1093/plankt/fbad048>
- Marsden L, Schneider O. The Nansen legacy template generator for Darwin core and CF-NetCDF. *Data Sci J*, 2024;23:38. <https://doi.org/10.5334/dsj-2024-038>
- Martin-Cabrera P, Perez Perez R, Irissou J-O *et al.* Best Practices and Recommendations for Plankton Imaging Data Management: Ensuring Effective Data Flow Towards European Data Infrastructures. Ostend: Flanders Marine Institute, 2022, 1–31. <https://doi.org/10.25607/OBP-1742>
- Martini S, Larras F, Boyé A *et al.* Functional trait-based approaches as a common framework for aquatic ecologists. *Limnol Oceanogr* 2021;66:965–94. <https://doi.org/10.1002/lno.11655>
- Meunier CL, Boersma M, El-Sabaawi R *et al.* From elements to function: toward unifying ecological stoichiometry and trait-based ecology. *Front Environ Sci* 2017;5:18. <https://doi.org/10.3389/fenvs.2017.00018>
- Morim T, Henriques S, Vasconcelos R *et al.* A roadmap to define and select aquatic biological traits at different scales of analysis. *Sci Rep* 2023;13:22947. <https://doi.org/10.1038/s41598-023-50146-9>
- Muller-Karger FE, Miloslavich P, Bax NJ *et al.* Advancing marine biological observations and data requirements of the complementary essential ocean variables (EOVs) and essential biodiversity variables (EBVs) frameworks. *Front Mar Sci* 2018;5:211. <https://doi.org/10.3389/fmars.2018.00211>
- Negrete-García G, Luo JY, Long MC *et al.* Plankton energy flows using a global size-structured and trait-based model. *Prog Oceanogr* 2022;209:102898. <https://doi.org/10.1016/j.pocean.2022.102898>
- O'Brien TD, Blanco-Bercial L, Questel JM *et al.* MetaZooGene Atlas and database: reference sequences for marine ecosystems. In: R. DeSalle (ed.) *DNA Barcoding. Methods in Molecular Biology*. Vol. 2744. New York, NY: Humana, 2024. https://doi.org/10.1007/978-1-0716-3581-0_28
- Orenstein EC, Ayata S-D, Maps F *et al.* Machine learning techniques to characterize functional traits of plankton from image data. *Limnol Oceanogr* 2022;67:1647–69. <https://doi.org/10.1002/lno.12101>
- Parr CS, Schulz KS, Hammock J *et al.* TraitBank: practical semantics for organism attribute data. *Semant Web J* 2016;7:577–88. <https://doi.org/10.3233/SW-150190>
- Pata PR, Hunt BPV. Harmonizing marine zooplankton trait data toward a mechanistic understanding of ecosystem functioning. *Limnol Oceanogr* 2023. <https://doi.org/10.1002/lno.12478>
- Perhirin M, Gossner H, Godfrey J *et al.* Morphological and taxonomic diversity of mesozooplankton is an important driver of carbon export fluxes in the ocean. *Mol Ecol Resour* 2024;24:e13907. <https://doi.org/10.1111/1755-0998.13907>
- Petchey OL, Gaston KJ. Functional diversity: back to basics and looking forward. *Ecol Lett* 2006;9:741–58. <https://doi.org/10.1111/j.1461-0248.2006.00924.x>
- Picheral M, Colin S, Irissou J-O. *EcoTaxa, a Tool for the Taxonomic Classification of Images*. 2017. <http://ecotaxa.obs-vlfr.fr>. Date accessed 30 July 2024.
- Pinti J, DeVries T, Norin T *et al.* Model estimates of meta-zoans' contributions to the biological carbon pump. *Biogeosciences* 2023;20:997–1009. <https://doi.org/10.5194/bg-20-997-2023>
- Pomerleau C, Sastri AR, Beisner BE. Evaluation of functional trait diversity for marine zooplankton communities in the Northeast subarctic Pacific Ocean. *J Plankton Res* 2015;37:712–26. <https://doi.org/10.1093/plankt/fbv045>
- Prowe AEF, Visser AW, Andersen KH *et al.* Biogeography of zooplankton feeding strategy. *Limnol Oceanogr* 2019;64:661–78. <https://doi.org/10.1002/lno.11067>
- Razouls C, de Bovée F, Kouwenberg J *et al.* Diversity and Geographic Distribution of Marine Planktonic Copepods. 2005–2024. <http://copepodes.obs-banyuls.fr/en>. Date accessed 30 July 2024.
- Renaud PE, Daase M, Banas NS *et al.* Pelagic food-webs in a changing Arctic: a trait-based perspective suggests a mode of resilience. *ICES J Mar Sci* 2018;75:1871–81. <https://doi.org/10.1093/icesjms/fsy063>
- Rohr T, Richardson AJ, Lenton A *et al.* Zooplankton grazing is the largest source of uncertainty for marine carbon cycling in CMIP6 models. *Commun Earth Environ* 2023;4:212. <https://doi.org/10.1038/s43247-023-00871-w>
- Romagnan JB, Legendre L, Guidi L *et al.* Comprehensive model of annual plankton succession based on the whole-plankton time series approach. *PLoS One* 2015;10:e0119219. <https://doi.org/10.1371/journal.pone.0119219>
- Rosati I, Bergami C, Stanca E *et al.* A thesaurus for phytoplankton trait-based approaches: development and applicability. *Ecol Inform* 2017;42:129–38. <https://doi.org/10.1016/j.ecoinf.2017.10.014>
- Sayers EW, Beck J, Brister JR *et al.* Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 2020;48:D9–D16. <https://doi.org/10.1093/nar/gkz899>
- Schneider FD, Fichtmueller D, Gossner MM *et al.* Towards an ecological trait-data standard. *Methods Ecol Evol* 2019;10:2006–19. <https://doi.org/10.1111/2041-210X.13288>
- Serra-Pompei C, Ward BA, Pinti J *et al.* Linking plankton size spectra and community composition to carbon export and its efficiency. *Global Biogeochem Cycles* 2022;36:e2021GB007275. <https://doi.org/10.1029/2021GB007275>
- Shaw CT, Bi H, Feinberg LR *et al.* Cohort analysis of *Euphausia pacifica* from the Northeast Pacific population using a gaussian mixture model. *Prog Oceanogr* 2021; 191:102495. <https://doi.org/10.1016/j.pocean.2020.102495>
- St.-Gélais NF, Giorgio del, P. A *et al.* Functional community structure modulates zooplankton production rates across boreal lakes. *Freshw Biol* 2023;68:837–46. <https://doi.org/10.1111/fwb.14067>
- Steinberg DK, Landry MR. Zooplankton and the ocean carbon cycle. *Ann Rev Mar Sci* 2017;9:413–44. <https://doi.org/10.1146/annurev-marine-010814-015924>
- Stemmann L, Boss E. Plankton and particle size and packaging: from determining optical properties to driving the biological pump. *Ann Rev Mar Sci* 2012;4:263–90. <https://doi.org/10.1146/annurev-marine-120710-100853>
- Teuber L, Hagen W, Bode M *et al.* Who is who in the tropical Atlantic? Functional traits, ecophysiological adaptations and life strategies in tropical calanoid copepods. *Prog Oceanogr* 2019;171:128–35. <https://doi.org/10.1016/j.pocean.2018.12.006>
- Thomsen PF, Willerslev E. Environmental DNA—an emerging tool in conservation for monitoring past and present biodiversity. *Biol Conserv* 2015;183:4–18. <https://doi.org/10.1016/j.biocon.2014.11.019>
- Thorson JT, Maureaud AA, Frelat R *et al.* Identifying direct and indirect associations among traits by merging phylogenetic comparative methods and structural equation models. *Methods Ecol Evol* 2023;14:1259–75. <https://doi.org/10.1111/2041-210X.14076>
- Vilgrain L, Maps F, Picheral M *et al.* Trait-based approach using in situ copepod images reveals contrasting ecological patterns across an Arctic ice melt zone. *Limnol Oceanogr* 2021;66:1155–67. <https://doi.org/10.1002/lno.11672>
- Violle C, Navas ML, Vile D *et al.* Let the concept of trait be functional! *Oikos* 2007;116:882–92. <https://doi.org/10.1111/j.0030-1299.2007.15559.x>
- Visconti A, Caroni R, Rawcliffe R *et al.* Defining seasonal functional traits of a freshwater zooplankton community using $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ stable isotope analysis. *Water* 2018;10:108. <https://doi.org/10.3390/w10020108>

- Wang Z, Gerstein M, Snyder M. RNA-seq: a revolutionary tool for transcriptomics. *Nat Rev Genet* 2009;10:57–63. <https://doi.org/10.1038/nrg2484>
- Wieczorek J, Bloom D, Guralnick R *et al.* Darwin Core: an evolving community-developed biodiversity data standard. *PLoS One* 2012;7:e29715. <https://doi.org/10.1371/journal.pone.0029715>
- Wilkinson MD, Dumontier M, Aalbersberg IJ *et al.* The FAIR Guiding Principles for scientific data management and stewardship. *Sci Data* 2016;3:160018. <https://doi.org/10.1038/sdata.2016.18>
- Wright RM, Le Quéré C, Buitenhuis E *et al.* Role of jellyfish in the plankton ecosystem revealed using a global ocean biogeochemical model. *Biogeosciences* 2021;18:1291–320. <https://doi.org/10.5194/bg-18-1291-2021>

Handling Editor: Howard Browman