


Regulated Ecosystem Model (REcoM) Recent advances and plans ahead

Judith Hauck, Christopher Danek, Özgür Gürses, Miriam Seifert, Christoph Völker,
Ying Ye


natESM Focus Workshop on Ocean Biogeochemistry




Ocean carbon



Ocean carbon



Ocean carbon under change



Marine ecosystem drivers



Warming, acidification,
nutrients, oxygen, plankton
productivity



**SUSTAINABLE
DEVELOPMENT** **GOALS**
17 GOALS TO TRANSFORM OUR WORLD

SUSTAINABLE DEVELOPMENT GOAL 14

Conserve and sustainably use the oceans, seas and marine resources for sustainable development




Ocean carbon & modelling

**AWI- Earth
System Model**

**Biogeochemical
Model REcoM**

**Process
understanding**




FESOM2 REcoM
Finite volumE Sea ice-Ocean Model Regulated Ecosystem Model

CMIP6 Interim
data available in
concentration-
and emission-
driven mode

Preparing for
CMIP7

REcoM Regulated Ecosystem Model



“Regulated Ecosystem”

- Photosynthesis, nitrate assimilation, chlorophyll synthesis are described based on environmental conditions and internal stoichiometry of the cells (Geider et al., 1998)
- Silicate assimilation added by Hohn (2009)
- Comprehensive representation of iron chemistry (e.g. Ye et al., 2020)
- Biogeochemical cycling of carbon, nitrogen, silicon, iron, oxygen

Geider model





Photosynthesis is a saturating function of irradiance

Initial slope increases with increasing Chl:C

Light-saturated rate increases with increasing N:C

Chlorophyll synthesis coupled to protein synthesis and thus to nitrate assimilation

Chl synthesis downregulated at high light (high light saturation E_0/E_k)




Nitrate assimilation is a saturating function of nitrate concentration

Maximum uptake rate is downregulated at high values of N:C

Major respiratory costs are associated with protein synthesis (reduction of nitrate to ammonium, incorporation of ammonium into amino acids and polymerization of amino acids into proteins)


Various levels of complexity



tracers	22	25	30	33
Wall clock time per simulated year (1152 CPUs)	35 min	39	46	50

REcoM features

- Explicit calcifiers, CO₂-sensitivity of phytoplankton growth, interactive effects of drivers



- Explicit calcifiers, CO₂-sensitivity of phytoplankton growth, interactive effects of drivers (Seifert et al.)
- Three zooplankton groups, and particle-sinking scheme that accounts for ballast minerals, seawater viscosity, and oxygen-dependent remineralization

Microzooplankton (e.g., ciliates)



Fast-growing herbivore group, main grazer of (small) phytoplankton

Mesozooplankton (e.g., copepods)



Grazes on larger phytoplankton and microzooplankton, fast sinking fecal pellets


(Polar) macrozooplankton (krill)



Slow-growing, cold-water species, fast-sinking fecal pellets with high C:N ratio

REcoM features

- Explicit calcifiers, CO₂-sensitivity of phytoplankton growth, interactive effects of drivers (Seifert et al.)
- Three zooplankton groups, and advanced particle-sinking (Karakus et al.)
- Carbon isotopes (Butzin et al., 2024)
- CFCs



REcoM features




PAL
MOD

GERMAN
CLIMATE
MODELING
INITIATIVE



- Explicit calcifiers, CO₂-sensitivity of phytoplankton growth, interactive effects of drivers (Seifert et al.)
- Three zooplankton groups, and advanced particle-sinking (Karakus et al.)
- Carbon isotopes (Butzin et al., 2024)
- CFCs
- Coupled to sediment model MEDUSA (Ye et al., 2025)



REcoM features




PAL
MOD

GERMAN
CLIMATE
MODELING
INITIATIVE




- Explicit calcifiers, CO₂-sensitivity of phytoplankton growth, interactive effects of drivers (Seifert et al.)
- Three zooplankton groups, and advanced particle-sinking (Karakus et al.)
- Carbon isotopes (Butzin et al., 2024)
- CFCs
- Coupled to sediment model MEDUSA (Ye et al., 2025)
- Tracer parallelization (Himstedt, 2025)




REcoM features

- Explicit calcifiers, CO₂-sensitivity of phytoplankton growth, interactive effects of drivers (Seifert et al.)
- Three zooplankton groups, and advanced particle-sinking (Karakus et al.)
- Carbon isotopes (Butzin et al., 2024), CFCs
- Coupled to sediment model MEDUSA (Ye et al., 2025)
- Tracer parallelization (Himstedt, 2025)
- Data-assimilation (ensemble Kalman filter, PDAF)




REcoM features

- Explicit calcifiers, CO₂-sensitivity of phytoplankton growth, interactive effects of drivers (Seifert et al.)
- Three zooplankton groups, and advanced particle-sinking (Karakus et al.)
- Carbon isotopes (Butzin et al., 2024), CFCs
- Coupled to sediment model MEDUSA (Ye et al., 2025)
- Tracer parallelization (Himstedt, 2025)
- Data-assimilation (ensemble Kalman filter, PDAF)
- Adding marine N₂O emissions, MSc thesis M. Vollmayr (split reactive nitrogen into redox states, model production of N₂O)



Interfaces/Integration


- Coupled to FESOM, AWI-CM/ESM (versions 1, 2, 3), MITgcm



Interfaces/Integration

- Coupled to FESOM, AWI-CM/ESM (versions 1, 2, 3), MITgcm

Atmospheric CO₂ in AWI-ESM3 (work in progress)



Interfaces/Integration




**PAL
MOD**


GERMAN
CLIMATE
MODELING
INITIATIVE



- Coupled to FESOM, AWI-CM/ESM (versions 1, 2, 3), MITgcm
- Coupled set-up for paleo reconstruction



Silicon redistribution after Heinrich Stadials stimulated low-latitude diatom growth




- AWI-ESM reproduces patterns of opal deposition observed in sediment cores from equatorial regions for the PI, LGM and deglacial periods.
- Higher opal deposition during deglaciation due to redistribution of Si from Southern Ocean surface waters



Interfaces/Integration

- Coupled to FESOM, AWI-CM/ESM (versions 1, 2, 3), MITgcm
- Coupled set-up for paleo reconstruction
- Sprint proposal in review: implement REcoM into FABM (Framework for Aquatic Biogeochem. Models)
- Exploiting FESOM flexible mesh to resolve physical processes in regions of interest (Southern Ocean, Arctic, coasts)

Arctic Ocean: 4.5 km resolution



Abruptly attenuated carbon sequestration



→ A variety of changes lower the density and volume of newly formed bottom waters and reduce the associated carbon transport to the abyss.

Tipping point cascade from physics to ecosystems initiated on Weddell Sea continental shelves, avoidable in a 2°C scenario: from “dense shelf” to “warm shelf”


Erosion of density differences shelf vs. open ocean

→ Flushing of shelf with warm deep water

→ Increase ice-shelf basal melt rates


→ Impacts on

- Global sea level
 - destabilization of ice sheets
- ocean circulation and climate
 - AABW: heat and carbon transfer to depth
- Ecosystems
 - deep ocean oxygenation
 - acidification and oxygen decrease on the shelves



Regime shift Weddell Sea shelves

Tipping point cascade from physics to ecosystems initiated on Weddell Sea continental shelves, avoidable in a 2°C scenario: from “dense shelf” to “warm shelf”



A vast icefish breeding colony discovered in the Antarctic




Purser et al., 2022


Arctic land-ocean carbon coupling

- Despite increased primary production, **the efficiency of the Arctic Ocean's biological carbon pump is projected to decrease by 40%**, largely due to enhanced remineralization.
- Terrigenous inputs lead to intense outgassing on the Arctic coasts.

Negative feedback loop




Positive feedback loop




Arctic land-ocean carbon coupling

The full picture is more complicated and involves stoichiometry changes.



Interfaces/Integration

- Coupled to FESOM, AWI-CM/ESM (versions 1, 2, 3), MITgcm
 - Coupled set-up for paleo reconstruction
 - Sprint proposal in review: implement REcoM into FABM
 - Exploiting FESOM flexible mesh to resolve physical processes in regions of interest (Southern Ocean, Arctic, coasts)
-
- Running on supercomputers at AWI, DKRZ (Levante), NHR
 - REcoM GitHub repository used as submodule in FESOM
 - Planned: continuous integration




Model application in emission-driven AWI-ESM

**AWI- Earth
System Model**



**Biogeochemical
Model REcoM**



**Process
understanding**




Ice

FESOM2 REcoM
Finite volumE Sea ice-Ocean Model Regulated Ecosystem Model

CO₂

Future carbon cycle in emissions-driven ESMs


- Analysing differences between emissions- and concentration-driven simulations in various scenarios
- Characterising pathway-dependent change in ocean and land sink



Ocean alkalinity enhancement and phytoplankton


Effects of OAE on net primary production

Without carbonate system
effects on phytoplankton




Ocean alkalinity enhancement and phytoplankton

Effects of OAE on net primary production




Indirect effects of OAE **decrease phytoplankton NPP by up to 15%** in the model version with carbonate system effects on phytoplankton

Effects of phytoplankton on OAE efficiency



Stronger CO_2 uptake by the ocean **with carbonate system effects on phytoplankton**

Role of ocean biology for atmospheric CO₂



Effect of biology, ca 220 ppm
Preindustrial control experiment

$500/280 = 1.8$ “nearly twofold higher”

The absolute change is related to the amount of biological carbon in the ocean.
The relative change is not transferable to the present-day with higher atmospheric CO₂.
Land feedbacks missing, how does that compare to state-of-the-art models?

No-biology experiments in emission-driven AWI-ESM

- Lower and slower than in Maier-Reimer et al.
- Less diffuse ocean models and land carbon feedbacks

On-going & outlook


Other on-going projects:

- Southern Ocean freshwater effects on carbon (SOFIA)
- Quantifying and addressing ventilation biases
- Mapping polar plankton habitats and their change
- Sprint proposal in review: implement REcoM into FABM
- Coastal set-up

Within REcoM:

- Additional calcifiers and improved representation of alkalinity
- New phytoplankton types: phaeocystis, second diatom
- Tracer for terrestrial POC
- Plankton response to warming and heatwaves
- Marine nitrogen cycling
- pH/temperature-controlled trace metal cycling

FESOM2 **REcoM**
Finite volume Sea ice-Ocean Model Regulated Ecosystem Model



Thank you!

