

Vegetation Differentiation across a topographic Yedoma – Alas Transect in the High Arctic Tundra of Oyogos Yar, East Siberia

Frank Kienast

Research Institute and Museum for Natural History Senckenberg, Research Station for Quaternary Palaeontology Weimar, Am Jakobskirchhof 4, D-99423 Weimar, Germany,
e-mail: fkienast@senckenberg.de

Lutz Schirrmeister

Alfred Wegener Institute for Polar and Marine Research, Telegrafenberg A43, D-14473 Potsdam, Germany

Sebastian Wetterich

Alfred Wegener Institute for Polar and Marine Research, Telegrafenberg A43, D-14473 Potsdam, Germany

Abstract

Excessive moisture is regarded the main cause for the fall of Pleistocene tundra-steppe and the rise of modern tundra. The arctic tundra of Oyogos Yar is low diverse in plant species (ca 100). The floral composition is mainly the result of differences in moisture or drainage, respectively. We describe a vegetational profile recorded in August 2007 along a 10 km transect from the top of a Yedoma ridge down to the adjacent Alas depression between 40 m to 10 m a.s.l.. Six main landscape units are described with respect to their floristic composition: Yedoma top with thermokarst mounds, mud boils, Yedoma slopes, small thermokarst ponds, thermo-erosional valleys, and the bottom of thermokarst depressions. Arctic thermokarst landscapes with Yedoma ridges and Alas depressions can be well classified according to their vegetation. The main constituents of the plant cover at well drained sites are grasses and polar willows, whereas excessively wet sites are occupied by sedges, cotton grass and peat moss.

Keywords: Alas depression; Bioindication; Moisture regime; Thermokarst; Tundra vegetation; Yedoma elevation

Introduction

The arctic vegetation cover reflects very well the small-scale periglacial landscape differentiation. Detailed surveys of plant associations are essential for the understanding of biotic responses to changes in permafrost landscapes. Around Beringia, the great influence of topography on arctic vegetation has been described from Alaska (Walker, 2000, Kade et al., 2005) and from the Taymyr Peninsula (Matveyeva, 1994).

Such studies of modern tundra vegetation are however little-known from Arctic Yakutia. Within the frame of the joint Russian-German expedition “Lena – New Siberian Islands 2007”, we studied relief-vegetation interactions at the coast of the Dimitrii Laptev Strait in August 2007.

Regional Setting

Oyogos Yar is the name of the mainland coast of the Dimitrii Laptev Strait (Fig. 1) between the mouth of the Kondrat’eva River in the east and Cape Svyatoy Nos in the west. This landscape is part of the Yana-Indigirka Lowland in Northeastern Siberia. Up to 500 m thick continuous permafrost and wide spread thermokarst characterize the coastal lowland. Oyogos Yar’s topography is dominated by extremely flat plains covered by mires and shallow lakes.

There are two main topographic elements: low elevations, so-called Yedoma, which represent the Pleistocene ground level and thermokarst depressions (alases), which formed as result of thermal degradation of the ice-rich permafrost that constitutes the Yedoma.

According to Aleksandrova (1980), Oyogos Yar belongs to the Sellyakh Inlet – Indigirka Delta district of the East Siberian province of the southern arctic tundra characterized by the dominance of *Alopecurus alpinus* and *Salix polaris*, the presence of *Carex ensifolia* ssp. *arctisibirica* and the absence of subarctic elements like *Betula nana* s.l. According to the Circumpolar Arctic vegetation map (CAVM-Team, 2003), the study area is covered with sedge/




Figure 1: Position of the study area (black square corresponds to Figure 2) at the mainland coast of the Dimitrii Laptev Strait

grass, moss wetland (W1) with *Carex aquatilis*, *Arctophila fulva*, *Dupontia*, and *Eriophorum* spp..

The study area is located about 8 km west of the Kondrat'eva River mouth (Fig. 2) opposite to Cape Shalaurova, the eastern edge of Bol'shoy Lyakhovsky Island. The climate is characterized by cold winters, cool summers and low precipitation. Climate data from the weather station Cape Shalaurova, about 80 km north of the study site, reflect a mean July air temperature of 2.8 °C, a mean January air temperature of -32.2 °C and an annual precipitation of 253 mm (Rivas-Martínez, 1996-2004).




Figure 2: Study transect west of the Kondrat'eva River mouth

Site description

The study transect extends across the bottom of a large alas depression of about 10 km in diameter (5 to 10 m a.s.l.) and the adjacent slope and top areas of a Yedoma hill of up to 40 m height (Figs. 2 & 3). The alas bottom dominantly consists of polygonal wetland tundra with a 0.5 to 1.0 m thick peat cover.



Figure 3: View from the alas bottom to the Yedoma hill

The thermokarst depression is cut by the coast of the Dimitrii Laptev Strait in the north and additionally intersected by several thermo-erosional valleys that drained to the coast.

Within the recorded transect, six main landscape units are following described with respect to their floristic composition: the Yedoma with thermokarst mounds, mud boils, Yedoma slopes, small thermokarst ponds, thermo-erosional valleys, and the bottom of thermokarst depressions.

Vegetation characteristics

Thermokarst mounds on the Yedoma

Thermokarst mounds are the best drained habitats in the study area (Fig. 4). Their plant cover is mainly composed of *Salix polaris*, *Dryas punctata*, and *Alopecurus alpinus*. Other grasses such as *Festuca brachyphylla* and *Deschampsia borealis* and dicots like *Potentilla hyperctica*, *Oxyria digyna*, *Papaver polare* and *Valeriana subcapitata* also occur.



Figure 4: Thermokarst mounds on the Yedoma at Oyogas Yar

Mud boils

Mud boils are the result of cryoturbation caused by frost pressing. In consequence, muddy soil flooded the ground. The substrate is silty and well drained. Mud boils occur at places most exposed and windswept on the Yedoma. Plants occur here only between such mud spots; the coverage is consequently very low with 20 to 40 % (Fig. 5). *Potentilla hyperctica*, *Salix polaris* and low growing grasses and rushes like *Festuca brachyphylla*, *Deschampsia borealis* and *Luzula confusa* are the main constituents of such habitats. In addition, herbs such as *Lloydia serotina*, *Cardamine bellidifolia*, *Androsace triflora*, and *Tephrosieris atropurpurea* occur in lower abundances.

This vegetation is similar in composition to cryptogam, herb barren (B1) or to the gramioid tundra (G1), described in the Circumpolar Arctic vegetation map (CAVM-Team 2003).



Figure 5: Mud boil at the top of the Yedoma visible in Fig.3

Yedoma slopes

At Yedoma slopes, the coverage is in general > 80%. In the upper parts of slopes in SW exposition, *Dryas punctata* is one of the main constituents. *Salix polaris* and several grass species (*Alopecurus alpinus*, *Deschampsia borealis* and *Festuca brachyphylla*) and *Luzula confusa* are characteristic of Yedoma slopes. In lower parts of the slopes, where it is less drained and consequently moister, *Arctagrostis latifolia*, *Petasites frigidus*, several saxifrages (*S. nelsoniana*, *S. cernua*, *S. hieracifolia*) and other herbs (*Gastrolychnis apetala*, *Tephroseris atropurpurea*, *Ranunculus spp.*) are typical (Fig. 6).



Figure 6: Lower part of a Yedoma slope with dominating *Arctagrostis latifolia*. In the background, a thermo-erosional valley with reddish spectral signature (here: dark) is visible

Small thermokarst ponds

In small depressions on the Yedoma, ponds with *Pleuropogon sabinei* and on the shore, *Arctophila fulva*, *Arctagrostis latifolia*, *Ranunculus hyperboreus*, *Dupontia fischeri*, *Eriophorum polystachion* and *E. scheuchzeri* occur (Fig. 7).

Interestingly, genuine aquatics were widely lacking. Only *Hippuris vulgaris* was solitarily found in a sterile form. The white-flowered *Ranunculus pallasii* and *Caltha palustris* grew emerged in some ponds within the Alas depressions.



Figure 7: Thermokarst pond with dominating *Pleuropogon sabinei* growing emerged in the water (inserted photo)

Thermo-erosional valleys

Thermo-erosional valleys are permanently supplied by running water. They are characteristically colored and recognizable from far distant (Fig. 6). Dark green and reddish signatures are mainly caused by different *Eriophorum* species: green – *E. scheuchzeri*, red – *E. polystachion* (Figs. 6 & 8). Other plants of thermo-erosional valleys are *Petasites frigidus* and several crowfoot and grass species (*Dupontia fischeri*, *Calmagrostis holmii*).



Figure 8: Thermo-erosional valley intersecting the Yedoma of Oyogos Yar. Main constituent is here *Eriophorum scheuchzeri* causing a green spectral signature.

Bottom of thermokarst depressions

The bottom of thermokarst depressions, Alases, is in contrast to thermo-erosional valleys characterized by stagnant water and covered mainly with sedges (*Carex ensifolia* ssp. *arctisibirica*) and cotton grass (*Eriophorum polystachion*).

The vascular plant diversity is here the lowest in the study area. *Sphagnum* moss is widely present, causing irregular pale green spots in polygonal wetlands where the surface of water is above the ground (Fig. 9).

At sites outside water bodies, rushes (*Luzula nivalis*, *L. confusa*) cover large areas together with several grasses



Figure 9: Bottom of a thermokarst depression with a water table above the ground. The polygonal surface patterns are visually strengthened by vegetational differentiation (compare Fig. 10).

(*Dupontia fischeri*, *Calamagrostis holmii*, *Poa alpigena* and *Arctophila fulva*). The wettest places are almost exclusively occupied by *Eriophorum polystachion*, which produces reddish patterns on the ground indicating the water trenches between polygons from afar (Figs. 9 & 10).



Figure 10: The wettest places in high center polygonal wetland tundra are the inter-polygonal trenches. Here, *Eriophorum polystachion* is the main constituent, causing a reddish (here dark) pattern.

Conclusions

The main landscape units in thermokarst affected landscapes can be well distinguished by their vascular plant cover.

Moisture or drainage respectively is the most important ecological factor in the study area resulting in the strongest vegetation differentiation.

Subordinate factors are exposure and declination. There were no really dry places in the study area.

Excessive wetness is well indicated by plants with characteristic spectral properties and therefore visible from far distances.

The plant species composition can alter quickly on short distances reflecting moisture changes resulting from the damming effect of the frozen ground.

The existence of such small scale variations in the plant cover gives important implications for the interpretation of palaeobotanical records.

Acknowledgments

We thank all participants of the joint Russian-German expedition "Lena - New Siberian Islands 2007", who helped us collecting, drying and preparing the plants for the herbarium. The studies were supported by the German Science Foundation (KI 849/1).

References

- Aleksandrova, V.D., 1980. The Arctic and Antarctic: their division into geobotanical areas. Cambridge University Press, Cambridge.
- CAVM-Team, 2003. Circumpolar Arctic Vegetation Map, Conservation of Arctic Flora and Fauna (CAFF) Map no. 1. U.S. Fish and Wildlife Service, Anchorage, Alaska.
- Kade, A., Walker, D. A., Raynolds, M. K., 2005. Plant communities and soils in cryoturbated tundra along a bioclimate gradient in the Low Arctic, Alaska. *Phytocoenologia* 35 (4), 761-820.
- Matveyeva, N.V., 1994. Floristic classification and ecology of tundra vegetation of the Taymyr Peninsula, northern Siberia. *Journal of Vegetation Science* 5, 813-828.
- Rivas-Martinez, S., 1996-2004. Climate diagrams, Worldwide; *Bioclimatic Classification System*. Phytosociological Research Center, Spain. Online database, <http://www.globalbioclimatics.org/plot/rumys-s.htm>
- Walker, D.A., 2000. Hierarchical subdivision of Arctic tundra based on vegetation response to climate, parent material and topography. *Global Change Biology* 6 (Suppl. 1), 19-34.