

A consistent dataset of Antarctic ice sheet topography, cavity geometry, and global bathymetry

R. Timmermann¹, A. Le Brocq², T. Deen³, E.Domack⁴, P. Dutrieux³,
B. Galton-Fenzi⁵, H. Hellmer¹, A. Humbert⁶, D. Jansen⁷, A. Jenkins³,
A. Lambrecht⁸, K. Makinson³, F. Niederjasper¹, F. Nitsche⁹,
O.A. Nøst¹⁰, L.H. Smedsrud¹¹, and W.H.F. Smith¹²

Ralph.Timmermann@awi.de

¹ Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany

² University of Exeter, Exeter, UK

³ British Antarctic Survey, Cambridge, UK

⁴ Hamilton College, Clinton, NY, USA

⁵ University of Tasmania, Hobart, Tasmania

⁶ KlimaCampus, University of Hamburg, Hamburg, Germany

⁷ Swansea University, Swansea, UK

⁸ University Innsbruck, Innsbruck, Austria

⁹ Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA

¹⁰ Norwegian Polar Institute, Tromsø, Norway

¹¹ Bjerknæs Centre for Climate Research, Bergen, Norway

¹² Laboratory for Satellite Altimetry, NOAA NESDIS, Silver Spring, MD, USA

Abstract

Sub-ice shelf circulation and freezing/melting rates in ocean general circulation models depend critically on an accurate and consistent representation of cavity geometry. Existing global or pan-Antarctic data sets have turned out to contain various inconsistencies and inaccuracies. The goal of this work is to compile independent regional fields into a global data set. We use the S-2004 global 1-minute bathymetry as the backbone and add an improved version of the BEDMAP topography (ALBMAP bedrock topography) for an area that roughly coincides with the Antarctic continental shelf. The position of the merging line is individually chosen in different sectors in order to get the best out of each data set. High-resolution gridded data for upper and lower ice surface topographies and cavity geometry of the Amery, Fimbul, Filchner-Ronne, Larsen C and George VI Ice Shelves, and for Pine Island Glacier are carefully merged into the ambient ice and ocean topographies. Multibeam survey data for bathymetry in the former Larsen B cavity and the southeastern Bellingshausen Sea have been obtained from the data centers of Alfred Wegener Institute (AWI), British Antarctic Survey (BAS) and Lamont-Doherty Earth Observatory (LDEO), gridded, and blended into the existing bathymetry map. The resulting global 1-minute topography data set (RTopo-1) contains maps for upper and lower ice surface heights, bedrock bathymetry, and consistent masks for open ocean, grounded ice, floating ice, and bare land surface. The data set is available in NetCDF format from the PANGAEA database at [doi:10.1594/pangaea.741917](https://doi.org/10.1594/pangaea.741917).

1 Introduction

Heat and salt fluxes at the base of any ice shelf, the properties of water masses within the cavity, and the exchange with the open ocean in numerical simulations strongly depend on an accurate and consistent representation of ice-shelf draft and sub-ice shelf bedrock topography. Early attempts to quantify the contribution of ice shelf water to the Southern Ocean's hydrography (e.g. Hellmer and Jacobs, 1995; Beckmann et al., 1999; Timmermann et al., 2001; Assmann et al., 2003) had to admit significant uncertainties due to partly crude assumptions about the geometry of ice shelf cavities. Local or regional simulations of individual cavities and the adjacent seas were able to use more detailed datasets but suffered from uncertainties arising from the choice of open or closed boundary conditions (e.g. Gerdes et al., 1999; Grosfeld et al., 2001; Williams et al., 2001; Thoma et al., 2006; Dinniman et al., 2007) and had no possibility to investigate larger-scale impacts and feedbacks.

An estimate of the rate of ice mass loss from the Antarctic ice sheet is an important component in the IPCC's Fifth Assessment Report. Given that most of the Antarctic ice sheet drains into floating glaciers or ice shelves, model estimates of sub-ice shelf melting rates are crucial to obtain a reliable estimate of the southern hemisphere's ice mass budget. In order to reduce error bars for high-resolution simulations of the coupled ocean–sea ice–ice shelf system in circum-polar or global ocean general circulation models, we compiled consistent maps for Antarctic ice surface heights and global ocean topography that combine available gridded data with independent high-resolution datasets and original multi-beam echosounder surveys. To preserve as much as possible information from the source datasets and to ensure an easy interpolation to any model grid, we chose a resolution of 1 minute in zonal and meridional direction, although we are aware that large parts of the available information in high latitudes is less detailed than this grid spacing might suggest.

In this paper, we present the datasets used, discuss their spatial coverage and the preprocessing applied, the strategies followed for merging datasets, and the resulting maps of ice and bedrock topography. In contrast to the ongoing International Bathymetric Chart of the Southern Ocean (IBCSO) activities (e.g., Schenke and Ott, 2009), our main goal is not a remapping of bathymetric data in the region south of 50°S, but a consistent representation of Antarctic ice sheet/shelf topography and global bathymetry in a dataset that contains enough detail for a wide range of regional and larger-scale studies. The main target group includes, but is not restricted to ocean modellers who aim at a realistic representation of Southern

Ocean ice shelf processes in ocean general circulation models.

2 Datasets and processing

2.1 Overview

The RTopo-1 dataset (see Figs. 1 and 2 for an overview) comprises global 1-minute fields for

- bedrock topography (ocean bathymetry; surface topography of continents; bedrock topography under ice sheets/shelves)
- surface elevation (upper ice surface height for the Antarctic ice sheet/shelves; bedrock elevation for ice-free continent; zero for ocean)
- bottom surface height of the Antarctic ice sheet/ice shelf system (ice draft for ice shelves; zero in the absence of ice)
- masks to identify open ocean, ice sheet, ice shelf, and bare land surface
- locations of coast and grounding lines (consistent with mask) for easy plotting

For ice-free land surface, the *bedrock topography* and *surface elevation* maps are identical. Naturally, *bedrock topography* and *ice bottom surface height* are equal for grounded ice. Ice not connected to the Antarctic ice sheet, including glaciers on subantarctic islands and the Greenland ice sheet, is not covered in our dataset; these areas are treated as bare land surface.

Note that in the following we use the term “ice shelf topography” for a consistent combination of ice top and bottom surface heights. The term “cavity geometry” refers to a compilation of ice-shelf draft and sub-ice (cavity) bathymetry.

2.2 Data sources

2.2.1 World Ocean bathymetry

The nucleus of RTopo-1 is the S-2004 ocean bathymetry (Marks and Smith, 2006), which merges the gravimetry-based Smith and Sandwell (1997) map with the GEBCO 1-minute bathymetry. For the continental shelf regions of the Southern Ocean, where satellite gravity data are not available, this dataset features several






Figure 1: RTopo-1 global bedrock topography. On continents outside Antarctica, this represents the topography of the solid surface (e.g. bottom of lakes, surface elevation of glaciers and continental ice caps) and mirrors the S-2004 dataset.

uncertainties. Ice shelves are not represented at all; instead they appear as areas with zero surface height. For the Antarctic continent, S-2004 contains the (ice) surface elevation, but not the bedrock topography. For most of the World Ocean, however, S-2004 represents ocean bathymetry and continental bedrock topography with an impressive amount of detail. We therefore use this dataset as the global backbone bathymetry outside the immediate vicinity of Antarctica (see Fig. 3 for the location of merging lines and the following section for a discussion of their location). We keep lakes and other features with a surface height below mean sea level and no connection to the world ocean, but mark them as continent in the surface type mask (see Section 2.3).



2.2.2 Antarctic ice and bedrock topographies

Upper and lower surface elevations for the Antarctic ice sheet and some of the ice shelves (see below for those not included), and ocean bathymetry for most of the Antarctic continental shelf is based on an improved version of the BEDMAP (Lythe et al., 2001) dataset that has been compiled by Le Brocq et al. (2010) and will be referred to as ALBMAP hereafter. While the BEDMAP effort was mostly directed towards mapping the continental ice sheet (and its bed), ALBMAP features a greatly improved representation of sub-ice shelf cavities, which has been



Bedrock topography



Surface elevation



Ice bottom



Surface type mask




Figure 2: Antarctic subset of the RTopo-1 dataset. Top left: bathymetry (surface height of bedrock). Top right: surface elevation. Bottom left: ice bottom surface height (including ice shelf draft). Bottom right: surface type mask, blue = ocean, yellow = grounded ice, orange = floating ice, grey = bare bedrock (appears only on islands not attached to the Antarctic Ice Sheet here). Grey line marks the grounding line, black line the ice shelf front or coast.

achieved mostly by using an improved interpolation scheme and carefully correcting (mostly increasing) bedrock depth towards the grounding line for many ice shelves. Maybe most important, ALBMAP ensures consistency of the different maps along the grounding lines and introduces a surface type mask for grounded ice, floating ice, and open water.

Like BEDMAP, ALBMAP uses a stereographic projection, which is very convenient for studies of the Antarctic ice sheet/shelf system, but much less so for circumpolar or global ocean models. We use spherical Delauney triangulation as a basis for linear interpolation to our regular 1-minute (lon,lat) grid.

To ensure a smooth continuation of bathymetry into the sub-ice shelf cavities, we merge the ALBMAP data into S-2004 not strictly along the ice-shelf edge, but at a line that has been carefully adjusted to keep the best out of both datasets (Fig. 3). In the East Antarctic and many other places, the transition line follows the ice shelf front or Antarctic coast. In order to avoid the spurious signature of General Belgrano Bank, which has been reported to be only a minor rise (with an elevation of only 20 m above the surrounding flat bottom) by Nicholls et al. (2003), we use ALBMAP for the entire continental shelf in the southwestern Weddell Sea south of 66.15° S (which is the latitude between the Larsen B and C Ice Shelf areas) with the transition occurring between the 1000 m and 2000 m isobaths. Similar arguments apply to the choice to use ALBMAP bathymetry in the Amundsen and Ross Seas with the transition to S-2004 placed between 2000 and 4000 m water depth. Tangens hyperbolicus functions are used to ensure a smooth transition between the datasets without too much spurious blurring of gradients. The width of the transition corridor varies and be seen in the left panel of Fig. 3.

In order to use topographic information from surveys that have been conducted after the BEDMAP dataset had been published, we replace ice and ocean topographies by newer, regional datasets in several places outlined in the following sections. Again we use spherical triangulation as a basis for interpolation and tangens hyperbolicus functions of the ratio between "new" and "background" data in a data window of typically 1° width for the transition between datasets.

2.2.3 Filchner-Ronne Ice Shelf

Upper and lower surface heights for Filchner-Ronne Ice Shelf and the position of the ice shelf front are derived from the ice thickness datasets of Lambrecht et al. (2007) using their Eq. (2) with the proposed densities and parameter values. Given their small-scale and transient nature, the inlets downstream from Hemmen Ice Rise are filled with interpolated values.




Figure 3: Data sources for ocean bathymetry (left) and ice sheet/shelf topography (right). See Table 1 for an explanation of numbers. White areas mark transition zones between the datasets.

Bathymetry in the Filchner-Ronne Ice Shelf cavity is derived from the dataset compiled by Makinson and Nicholls (1999). In order to maintain floating conditions in the embayment with the inflow of Support Force Glacier, we deviate from the ALBMAP grounding line here and apply the grounding line position of Rignot and Jacobs (2002) instead, which is consistent with grounding line locations suggested by Makinson and Nicholls (1999) and Lambrecht et al. (2007) here. The ice plain to the west and the ice rumples east of Bungenstockrücken (Heidrich et al., 1992; Scambos et al., 2004) are treated as grounded ice.

Inconsistencies between the two datasets (ice thickness and bathymetry) are addressed by applying a minimum water column thickness of 10 m in the area of floating ice and enforcing zero water column thickness in locations with grounded ice. Given the high accuracy of the ice shelf thickness estimates (error quantified to be less than 25 m by the authors), we retain the Lambrecht et al. (2007) draft field and apply the corrections to the bathymetry; they are typically between 100 and 200 m and occur localized in a narrow band along the grounding line.

| Region | Data obtained from | Original data source(s) |
|---|---|--|
| 1. World Ocean bathymetry | S-2004 bathymetry (Marks and Smith, 2006) | 1. GEBCO 1-minute bathymetry 2. Smith and Sandwell (1997) |
| 2. Antarctic ice sheet/shelf upper and lower surface heights and bedrock topography | ALBMAP-v1 (Le Brocq et al., 2010) | BEDMAP (Lythe et al., 2001) |
| 3. Filchner-Ronne Ice Shelf • ice shelf topography • cavity bathymetry | radar and seismic sounding Makinson and Nicholls (1999) | Vaughan et al. (1995), Johnson and Smith (1997), Mayer et al. (1995) |
| 4. Amundsen Sea: • Pine Island Bay bathymetry | Nitsche et al. (2007) | AGASEA, ship data |
| 5. Pine Island Glacier • ice draft and cavity bathymetry | Jenkins et al. (2010) | orig. AUTOSUB data |
| 6. Larsen C Ice Shelf • ice shelf topography • cavity bathymetry | Jansen et al. (2010) | BEDMAP + ICESat altimetry estimate using grounding line depth and bathymetry at ice shelf front |
| 7. former Larsen B Ice Shelf cavity bathymetry | Domack (pers. comm.), AWI database | <i>Nathaniel B. Palmer</i> multibeam data, <i>Polarstern</i> multibeam data |
| 8. Bellingshausen Sea • bathymetry • George VI sub-ice bathymetry • George VI ice shelf topography | Deen (2009), Rottmann et al. (1996) Potter and Paren (1985) Humbert (2007) | BAS archived multibeam data, <i>Polarstern</i> multibeam data radar, plumb-line, seismics BEDMAP + ICESat laser altimetry |
| 9. Amery Ice Shelf • ice draft and cavity bathymetry | Galton-Fenzi et al. (2008) | borehole and ship-based measurements, radar, seismics, tidal modelling |
| 10. Fimbul Ice Shelf • ice draft and cavity bathymetry | Smedsrud et al. (2006) | Nøst et al. (2004) |

Table 1: Data sources for individual regions of the Southern Ocean, as merged into RTopo-1. Numbers in the Region column correspond to the source flag in Fig. 3.

2.2.4 Amundsen Sea ice shelves and Pine Island Glacier

As can be seen in Fig. 3, ocean bathymetry in most of the Amundsen Sea is derived from ALBMAP, with the transition to S-2004 occurring between 2000 and 4000 m water depth. Open ocean bottom topography in Pine Island Bay, however, utilizes the dataset from Nitsche et al. (2007), which is a combination of ship data with the *Airborne Geophysical Survey of the Amundsen Embayment* (AGASEA) and BEDMAP datasets. This dataset has already been included in ALBMAP, but to preserve its fine resolution we use the original Nitsche et al. (2007) data here.

For upper and lower ice surface height and cavity bathymetry for most ice shelves in the Amundsen Sea we retain the data from ALBMAP, which in turn uses AGASEA data (Vaughan et al., 2006; Holt et al., 2006) for Thwaites Glacier

Tongue, and Crosson and Dotson Ice Shelves.

An important exception is Pine Island Glacier, where the geometry of the sub-ice cavern (ice-shelf draft and sub-ice bathymetry) is interpolated from the very recent AUTOSUB data of Jenkins et al. (2010). While BEDMAP and S-2004 do not contain any meaningful information for the sub-PIG cavity, ALBMAP suggests a trough with a depth of about 900 m at the glacier front and a sill of about 100 m height at about two thirds of the distance between the ice shelf front and the grounding line. The AUTOSUB survey revealed that the trough is more than 1000 m deep at the ice front; the sill is located halfway between glacier front and grounding line, and its height is about 300 m (see Fig. 6 below). Water column thickness in the cavity thus varies from about 700 m near the glacier front to only 275 m at the sill, but then again increases to a maximum of 360 m towards the grounding line.

2.2.5 Larsen C Ice Shelf

For Larsen C Ice Shelf topography (surface elevation and draft) and grounding line location, we use data from Jansen et al. (2010), who combined BEDMAP data with surface heights obtained from ICESat altimetry. The grounding line location in this dataset has been picked from interferometry from the ERS-1/ERS-2 tandem mission in the north, and from MODIS imagery in the south. It is a rather conservative estimate in a sense that for all areas denoted as ice shelf we can be sure that the ice is really floating. Differences from BEDMAP draft occur mainly along the ice shelf front, where the ice is thinner now, and towards the grounding line, where the depth of the ice shelf base increases to about 550 m below sea level (Fig. 4, left).

Bottom topography in the cavity below Larsen C Ice Shelf is based on ALBMAP, but has been modified in order to ensure a minimum water column thickness of 10 m near the grounding line and a gradual rise to the bottom depth found near the ice shelf front. Bathymetry now features a depth between 500 and 600 m under most of the ice shelf with deep troughs towards the grounding line (Fig. 5). The existence of such deep troughs in immediate vicinity to the mountains of the Antarctic Peninsula might seem doubtful at first glance; multibeam bathymetry surveys in Antarctic Sound and the former Larsen B Ice Shelf cavity (Gavahan and Domack, 2006), however, show that indeed troughs of very similar depth and horizontal scale have been carved out where ice streams interfered with the ocean bottom. In any case it should be kept in mind that bathymetry in the Larsen C Ice Shelf cavity in our dataset is hardly more than an educated guess.




Figure 4: Surface elevation (left) and ice shelf draft (right) for the Antarctic Peninsula, including Larsen C ice shelf and the remnants of Larsen B ice shelf. The ice shelf front is marked with a black line, the grounding line location with a grey line.




Figure 5: Different representations of bathymetry (bedrock topography) in the area of the Larsen Ice Shelves. Top row: BEDMAP (left), S-2004 (middle), and ALBMAP (right). Bottom row: Multibeam track data (left), and the RTopo-1 product (right). Ice shelf front and coastline are marked with a black line, ground-line location with a grey line.

2.2.6 Larsen A and B Ice Shelves

The disintegration of Larsen A and B Ice Shelves in January 1995 and February 2002 left the former cavities as open water embayments. For the bathymetry in this area, we combine original data from *Polarstern* cruise ANT-XXIII/8 (Pugacheva and Lott, 2008) with digitized maps of along-track multibeam data from the *Nathaniel B. Palmer* NBP0107 and NBP0603 cruise reports (E. Domack, pers. comm.). Again spherical triangulation is used as a basis to fill the gaps between cruise tracks (Fig. 5). The coastline of the Larsen B embayment has been carefully corrected wherever the existence of ship tracks suggests the presence of open water instead of ice or continent.

2.2.7 Bellingshausen Sea and George VI Ice Shelf

For the eastern Bellingshausen Sea shelf area (i.e. the area shallower than 1000 m in the sector east of 90°W and south of 67°S), bathymetry in the open ocean is constructed from multibeam swath data from the BAS bathymetry database (Deen, 2009) augmented with original data from R/V *Polarstern* cruise ANT-XI/3 (Rottmann et al., 1996) in the area near Ronne Entrance (Fig. 6). Bottom topography underneath George VI Ice Shelf follows the sections of Potter and Paren (1985). Bathymetry in the ice shelf cavities on the western side of Alexander Island (Bach and Wilkins Ice Shelves) is retained from ALBMAP and is used as additional information for the interpolation. In contrast to the representations in BEDMAP, ALBMAP and S-2004 (Fig. 6, top row), these data taken together clearly indicate the existence of a 700-900 m deep trough that extends all the way from Marguerite Bay through George VI Sound (i.e. the cavity under George VI Ice Shelf) to Ronne Entrance. From here, further channels provide connections to the continental shelf break. Note that we had to make assumptions for bathymetry in the data gap in the northern part of George VI Sound, but these are fully consistent with the Potter and Paren (1985) plumb line profile along the northern ice shelf front.

Surface elevation and draft of George VI Ice Shelf is derived from the Humbert (2007) thickness dataset. Given that melt ponds occupy large parts of George VI Ice Shelf and that a distinguished firn layer is virtually absent, we convert thickness to draft assuming an ice density of 910 kg m^{-3} and do not apply any firn correction.




Figure 6: Different representations of bathymetry (bedrock topography) in the Bellingshausen Sea. Top row: BEDMAP (left), S-2004 (middle), and ALBMAP (right). Bottom row: Multibeam track data (left), and the RTopo-1 product (right). Note the differences for Pine Island Glacier (PIG) and George VI sub-ice bathymetries. Black line represents the coastline or ice shelf front, dark grey line the grounding line.

2.2.8 Amery Ice Shelf

For Amery Ice Shelf and the Prydz Bay region, we use the ice draft and ocean bathymetry data of Galton-Fenzi et al. (2008), who combine radar and seismic surveys, ice thickness estimates from satellite altimetry, borehole and ship-based observations, and insight obtained from tidal modelling. This dataset is substantially improved over BEDMAP and ALBMAP and features a much deeper ice draft (maximum ≈ 2500 m) and bathymetry (maximum bottom depth in the cavity ≈ 3000 m). It also introduces five grounded-ice regions (i.e. rumples and islands), and yields substantial corrections for grounding line and ice front location.

2.2.9 Fimbul Ice Shelf

For Fimbul Ice Shelf draft and sub-ice bathymetry we use the topography dataset of the regional model of Smedsrud et al. (2006). Smedsrud et al. (2006) interpolated original seismic data from Nøst (2004) covering the central and outer parts of the ice shelf. For the deeper parts of the cavity towards the grounding line where no seismic data is available, ice shelf draft and bathymetry were interpolated along the flow line of the Jutulstraumen ice stream. This interpolation leads to a channel connecting the 1100 m deep Jutul basin with the deepest grounding line at the Fimbul ice shelf at 875 m depth.

We have retained the ALBMAP grounding line location here, and ice extent has been carefully reduced to match the more recent data.

2.3 Water column thickness, masks and coastline

As already discussed, we provide a global mask that discriminates between grounded and floating ice, open ocean, and bare bedrock (Fig. 7). For Antarctica, the mask largely follows ALBMAP. However, as already mentioned, coast and grounding line locations in the Larsen B Ice Shelf area are corrected based on ship tracks. Modifications based on ice front or grounding line locations in the newly integrated local datasets are applied to Filchner-Ronne Ice Shelf (ice front; grounding line in the Support Force Glacier region), George VI ice shelf (ice front location in Ronne Entrance), Larsen C ice shelf (ice front and grounding line), Amery Ice Shelf (ice front, grounding line, ice rumples) and Fimbul Ice Shelf (ice front). Ice caps not connected to the Antarctic ice sheet have been removed from the mask and are now classified as bedrock (with the ice surface

height adopted as the bedrock surface height). Subglacial lakes are ignored.

While lakes and enclosed seas outside Antarctica are still present in the bathymetry dataset (adopted from S-2004), they are marked as “continent” in the mask. Using the topography map(s) together with the mask thus allows for an easy generation of global or regional ocean model grids without the need to manually remove features with a topography below mean sea level and no connection to the world ocean.




Figure 7: Global surface type mask in RTopo-1. Blue = ocean, white = grounded ice, orange = ice shelf, grey = bare land surface.

Water column thickness needed for the generation of grids for ocean models with a terrain-following coordinate can be computed as the difference between ice bottom surface height (which is zero in the absence of ice shelves) and ocean bedrock topography (Fig. 8). Locations of nonzero entries in the resulting field are consistent with the area denoted as “ocean” in the mask.

Last but not least, the dataset contains position data for coast and grounding lines that are consistent with the mask and all other datasets. These can be used for an easy and overlap-free plotting of maps in any desired projection (and have been used for all the maps in this paper).

Water column thickness




Figure 8: Water column thickness in the Southern Ocean sector of RTopo-1.

3 Summary and Outlook

We have presented a global 1-minute data set for World Ocean bathymetry and Antarctic ice sheet/shelf topography that compiles high-resolution data for the Amery, Fimbul, Filchner-Ronne, Larsen C and George VI Ice Shelves, and for Pine Island Glacier into a synthesis of the S-2004 global 1-minute bathymetry with a BEDMAP-derived panantarctic topography dataset. Wherever maps were derived from original bathymetry surveys (namely in the Larsen A/B Ice Shelf area and the southeastern Bellingshausen Sea), we presented data coverage and the resulting gridded fields. Next to maps for bedrock topography and the upper and lower surface heights of the Antarctic ice sheet/ice shelf system, the dataset contains consistent masks for open ocean, grounded ice, floating ice, and bare land surface.

Naturally, this kind of dataset can hardly be complete. We already mentioned that bathymetry under Larsen C Ice Shelf is not more than an educated guess; contributions to this topic are more than welcome. Further developments in the future might be related to using alternative interpolation schemes to fill the data gaps in Antarctic bedrock topography maps, like for example the streamline-following interpolation scheme presented by Warner and Roberts (2010). Any other input regarding local ice shelf/cavity geometry will be more than welcome and used to update the dataset as soon as possible.

4 Data access

The RTopo-1 dataset is available in NetCDF format in two flavours at [doi:10.1594/pangaea.741917](https://doi.org/10.1594/pangaea.741917):

1. The complete global 1-minute dataset has been split into two files:
 - RTopo<version>_data.nc (2.8 GB file size) contains the digital maps for bedrock topography, ice bottom topography, and surface elevation.
 - RTopo<version>_aux.nc (700 MB file size) contains the auxiliary maps for data sources and the surface type mask.
2. A regional subset that covers all variables for the region south of 50°S is available in RTopo<version>_50S.nc (780 MB file size).

Datasets for the location of grounding line (RTopo<version>_gl.asc, 1.7 MB) and coast line (RTopo<version>_coast.asc, 17.5 MB) are prepared in ASCII format and simply contain two columns for longitude and latitude, separated by blanks.

Acknowledgements

We would like to thank Povl Abrahamsen, Mike Dinniman, Dorothea Graffe, Hannes Grobe, Klaus Grosfeld, Verena Haid, Stanley S. Jacobs, Laura Jensen, Uta Menzel, Eric Rignot, and David Vaughan for their rapid-response help and support. AGASEA maps were obtained from <http://www.ig.utexas.edu/research/projects/agasea>. Galton-Fenzi (2008) topography data were provided by the Antarctic Climate and Ecosystems CRC. Lambrecht et al. (2007) thickness data and ALBMAP-v1 were obtained from the PANGAEA data supplements doi:10.1594/PANGAEA.615277 and doi:10.1594/PANGAEA.734145, respectively. All other digital data were received directly from the authors. This work was supported by funding to the ice2sea programme from the European Union 7th Framework Programme, grant number 226375. Ice2sea contribution number ###.

References

- Assmann, K., Hellmer, H. H., and Beckmann, A.: Seasonal variation in circulation and water mass distribution on the Ross Sea continental shelf, *Antarctic Science*, 15(1), 3-11, doi:10.1017/S0954102003001007, 2003.
- Beckmann, A., Hellmer, H.H., and Timmermann, R.: A numerical model of the Weddell Sea: Large-scale circulation and water mass distribution. *Journal of Geophysical Research*, 104(C10), pp. 23375-23391, 1999.
- Deen, T.J. (Editor): The British Antarctic Survey Geophysics Data Portal. British Antarctic Survey, Cambridge, UK. Available online at: <http://geoportal.nerc-bas.ac.uk/GDP>, 2009.
- Dinniman, M.S., Klinck, J.M., and Smith Jr, W. O.: Influence of Sea Ice Cover and Icebergs on Circulation and Water Mass Formation in a Numerical Circulation Model of the Ross Sea, Antarctica. *Journal of Geophysical Research*, 112, C11013, doi:10.1029/2006JC004036, 2007.
- Domack, E., Duran, D., Leventer, A., Ishman, S., Doane, S., McCallum, S., Amblas, D., Ring, J., Gilbert, R., and Prentice, Mi.: Stability of the Larsen B ice shelf on the Antarctic Peninsula during the Holocene epoch. *Nature* 436, 681-685, doi:10.1038/nature03908, 2005.
- Galton-Fenzi, B.K., Maraldi, C., Coleman, R., and Hunter, J.: The cavity under the Amery Ice Shelf, East Antarctica. *Journal of Glaciology*, 54 (188), 881-887, 2008.
- Gavahan, K., and Domack, E.: Paleohistory of the Larsen Ice Shelf Phase II, Year 3. NBP0603 Multibeam End of Cruise Report. 10 pp., 2006.
- Gerdes, R., Determann, J., and Grosfeld, K.: Ocean circulation beneath Filchner-Ronne-Ice-Shelf from three-dimensional model results. *Journal of Geophysical Research*, 104 (C7), 15827-15842, 1999.
- Grosfeld, K., Schröder, M., Fahrbach, E., Gerdes, R., and Mackensen, A.: How iceberg calving and grounding change the circulation and hydrography in the Filchner Ice Shelf-Ocean System. *Journal of Geophysical Research*, 106, C5, 9039-9055, doi:2000JC000601, 2001.
- Heidrich, B., Sievers, J., Schenke, H.W., and THIEL, M.: Digitale Topographische Datenbank Antarktis - Die Küstenregionen vom westlichen Neuschwabenland bis zum Filchner-Ronne Schelfeis interpretiert aus Satellitenbilddaten. *Nachrichten aus dem Karten- und Vermessungswesen*, I(107), 127-140, 1992.
- Hellmer, H.H., and Jacobs, S.S.: Seasonal circulation under the eastern Ross Ice Shelf, Antarctica. *Journal of Geophysical Research*, 100(C6), 10873-10885, 1995.

- Holt, J. W., Blankenship, D.D., Morse, D.L., Young, D.A., Peters, M.E., Kempf, S.D., Richter, T.G., Vaughan, D.G., and Corr, H.F.J.: New boundary conditions for the West Antarctic Ice Sheet: Subglacial topography of the Thwaites and Smith glacier catchments. *Geophys. Res. Lett.*, 33, L09502, doi:10.1029/2005GL025561, 2006.
- Humbert, A.: Numerical simulations of the ice flow dynamics of the George VI Ice Shelf, Antarctica. *Journal of Glaciology*, 53(183), 659-664, 2007.
- Jansen, D., Kulesa, B. ., Sammonds, P.R., Luckman, A.J., King, E.C., and Glasser, N.F.: Present Stability of the Larsen C Ice Shelf. *Journal of Glaciology*, in press, 2010.
- Jenkins, A., Dutrieux, P., Jacobs, S.S., McPhail, St.D., Perrett, J.R., Webb, A.T., and White, D.: Observations beneath Pine Island Glacier West Antarctica and implications for its retreat, *Nature Geoscience*, in press, 2010.
- Johnson, M.R. and Smith, A.M.: Seabed topography under the southern and western Ronne Ice Shelf, derived from seismic surveys. *Antarctic Science*, 9(2), 201-208, 1997.
- Lambrecht, A., Sandhäger, H., Vaughan, D.G., and Mayer, C.: New ice thickness maps of Filchner-Ronne Ice Shelf, Antarctica, with specific focus on grounding lines and marine ice. *Antarctic Science*, 12 pp, doi:10.1017/S0954102007000661, 2007.
- Le Brocq, A.M., Payne, A.J., and Vieli, A.: An improved Antarctic dataset for high resolution numerical ice sheet models (ALBMAP v1). *Earth System Science Data*, submitted, 2010.
- Lythe, M.B., Vaughan, D.G., and the BEDMAP Consortium: BEDMAP - A new ice thickness and subglacial topographic model of Antarctica. *Journal of Geophysical Research*, 106, pp. (B6),11335-11351, 2001.
- Makinson, K., and Nicholls, K.W.: Modeling tidal currents beneath Filchner-Ronne Ice Shelf and on the adjacent continental shelf: their effect on mixing and transport. *Journal of Geophysical Research*, 104(C6), 13449-13465, 1999.
- Marks, K.M., and Smith, W.H.F.: An evaluation of publicly available global bathymetry grids. *Marine Geophysical Researches*, 27, 19-34, DOI 10.1007/s11001-005-2095-4, 2006.
- Mayer, C., Lambrecht, A., and Oerter, H.: Glaciological investigations on the Foundation Ice Stream. Filchner Ronne Ice Shelf Programme Report No 9. H. Oerter. Bremerhaven, Germany, Alfred-Wegener-Institut für Polar und Marine Research: 57-63, 1995.
- Nicholls, K.W., Padman, L., Schröder, M., Woodgate, R.A., Jenkins, A., and Østerhus, S.: Water mass modification over the continental shelf north

- of Ronne Ice Shelf, Antarctica. *J. Geophys. Res.*, 108, C8, 3260, doi:10.1029/2002JC001713, 2003.
- Nitsche, F.O., Jacobs, S., Larter, R.D. and Gohl, K.: Bathymetry of the Amundsen Sea Continental Shelf: Implications for Geology, Oceanography, and Glaciology. *Geochemistry, Geophysics, Geosystems*, 8: Q10009, doi:10.1029/2007GC001694, 2007.
- Nøst, O. A.: Measurements of ice thickness and seabed topography at Fimbul Ice Shelf, Dronning Maud Land, Antarctica, *J. Geophys. Res.*, 109, C10010, doi:10.1029/2004JC002277, 2004.
- Potter, J.R., and Paren, J.G.: Interaction between ice shelf and ocean in George VI Sound, Antarctica. In: S.S. Jacobs (editor): *Oceanology of the Antarctic Continental Shelf*, Antarctic Research Series, 43, 35-58, American Geophysical Union, 1985.
- Pugacheva, E. and Lott, J.-H.: Bathymetry. In: Gutt, J. (ed), *The expedition ANTARKTIS-XXIII/8 of the research vessel "Polarstern" in 2006/2007*. *Berichte zur Polar- und Meeresforschung*, 569, 153 pp., 2008.
- Rignot, E., and Jacobs, S.S.: Rapid bottom melting widespread near Antarctic ice sheet grounding lines. *Science*, 296, 2020-2023, 2002.
- Rottmann, E., Grünwald, T., Niederjasper, F., and Weigelt, M.: Bathymetrie. In: *The Expedition Antarktis-XI/3 of RV "Polarstern" in 1994*, Reports on Polar Research, 188, 22-28, AWI, Bremerhaven, 1996.
- Scambos, T., Bohlander, J., Raup, B., and Haran, T.: Glaciological characteristics of Institute Ice Stream using remote sensing. *Antarctic Science* 16 (2), 205-213, doi:10.1017/S0954102004001919, 2004.
- Schenke, H.W., and Ott, N.: Report on the International Bathymetric Chart of the Southern Ocean (IBCSO). IHO Hydrographic Committee on Antarctica 9th Meeting, Simon's Town, 12-14 October 2009, available from <http://www.ibcso.org/history.html>, 2009.
- Smedsrud, L.H., Jenkins, A., Holland, D.M., and Nøst, O.A.: Modeling ocean processes below Fimbulisen, Antarctica. *Journal of Geophysical Research*, Volume 111, C01007, doi:10.1029/2005JC002915, 2006.
- Smith, W.H.F., and Sandwell, D.T.: Global sea floor topography from satellite altimetry and ship depth soundings. *Science*, **277**, 1956-1962, 1997.
- Thoma, M., Grosfeld, K., and Lange, M. A.: The impact of the Eastern Weddell Ice Shelves on water masses in the eastern Weddell Sea, *J. Geophys. Res.* 111, C12010, doi:10.1029/2005JC003212, 2006.
- Timmermann, R., Beckmann, A., Hellmer, H.H.: The role of sea ice in the fresh water budget of the Weddell Sea. *Annals of Glaciology*, 33, pp. 419-424, 2001.

- Vaughan, D.G., Sievers, J., Doake, C.S.M., Hinze, H., Mantripp, D.R., Pozdeev, V.S., Sandhäger, H., Schenke, H.W., Solheim, A., and Thyssen, F.: Subglacial and Seabed Topography, Ice Thickness and Water Column Thickness in the Vicinity of Filchner-Ronne-Schelfeis, Antarctica. *Polarforschung*, 64, 2, 75-88, hdl:10013/epic.29728.d001, 1995.
- Vaughan, D.G., Corr, H.F.J., Ferraccioli, F., Frearson, N., O'Hare, A., Mach, D., Holt, J., Blankenship, D., Morse, D., and Young, D.A.: New boundary conditions for the West Antarctic ice sheet: subglacial topography beneath Pine Island Glacier. *Geophys. Res. Lett.*, 33, L09501, doi:10.1029/2005GL025588, 2006.
- Warner, R., and Roberts, J.: Unveiling the Antarctic subglacial landscape. *Geophysical Research Abstracts*, 12, EGU2010-7864-2, EGU General Assembly, 2010
- Williams, M.J.M., Grosfeld, K., Warner, R., Gerdes, R., Determann, J.: Ocean circulation and ice-ocean interaction beneath the Amery Ice Shelf, Antarctica. *Journal of Geophysical Research*, 106 (C10), 22383-22399, 2001.