

Arthur et al. (1992)
Arthur, J. M. M., R. V. Tyson, J. Thomson, and D. Matthey (1992): Early diagenesis of marine organic matter: Alteration of the carbon isotopic composition: <i>Marine Geology</i> , v. 105, p. 51-61.
Abrantes (1990)
Abrantes, F (1990): Increased upwelling off Portugal during the last deglaciation: Diatom Evidence. <i>Marine Micropaleontology</i> 17: 285-310.
Abrantes (2001)
Abrantes, F (2001): unpublished data, processed and archived through EU project EUROCORE as part of the EU-SEASED initiative, Departamento de Geologia Marinha, Instituto Geológico e Mineiro, Portugal; PANGAEA - Network for Geological and Environmental Data - http://www.pangaea.de .
Abrantes et al. (1998)
Abrantes, F; Baas, J; Hafliðason, H; Rasmussen, T; Klitgaard, D; Lončarić, N & Gaspar, L (1998): Sediment fluxes along the northeastern European Margin: inferring hydrological changes between 20 and 8 kyr. <i>Marine Geology</i> 152: 7-23.
Aller (1977)
Aller, RC (1977): The influence of macrobenthos on chemical diagenesis of marine sediments. Unpublished Ph.D. Dissertation, Yale University, New Haven, Connecticut, 600.
Anderson et al. (1994)
Anderson, R F; Rowe, G T; Kemp, P F; Trumbore, S & Biscaye, P E (1994): Carbon budget for the mid-slope depocenter of the Middle Atlantic Bight. <i>Deep-Sea Research</i> 41: 669-703.
Anonymous (1995)
Anonymous (1995): ODER-Project Interim Report, European Commission Environment Program PL 910398.
Archer and Devol (1992)
Archer, D; Devol, A (1992): Benthic oxygen fluxes on the Washington shelf and slope: A comparison of in situ microelectrode and chamber flux measurements. <i>Limnol. Oceanogr.</i> 37(3): 614 - 629.
Arz (1998)
Arz, H (1998): Dokumentation von kurzfristigen Klimaschwankungen des Spätquartärs in Sedimenten des westlichen äquatorialen Atlantiks. <i>Berichte, Fachbereich Geowissenschaften, Universität Bremen</i> , 124: 96
Arz et al. (1999)
Arz, H W; Pätzold, J & Wefer, G (1999): Climatic changes during the last deglaciation recorded in sediment cores from the northeastern Brazilian Continental Margin. <i>Geo-Marine Letters</i> 19: 209-218 - SFB261 No 238
Avilov (1965)
Avilov, I K (1965): Relief and bottom sediments of shelf and continental slope of north-western part of the Atlantic Ocean (in Russian). <i>Researches on Programm of International Geophysic year. part 2. Transactions of VNIRO. Pischeprom. Moscow</i> 57: 173-234.
Bakker and Helder (1993)
Bakker, JF; Helder, W (1993): Skagerrak (northeastern North-Sea) oxygen microprofiles and porewater chemistry in sediments. <i>Marine Geology</i> 111(3-4), 299-321.

Bender et al. (1989)
Bender, M. L., R. Jahnke, R. Weiss, W. Martin, D. T. Heggie, J. Orchado, and T. Sowers (1989): Organic carbon oxidation and benthic nitrogen and silica dynamics in San Clemente basin, a continental borderland site: <i>Geochim. Cosmochim. Acta</i> , v. 53, p. 685-697.
Berelson et al. (1987)
Berelson, W M; Hammond, D E; Johnson, K S (1987): Benthic fluxes and the cycling of biogenic silica and carbon in two southern California borderland basins. <i>Geochim. Cosmochim. Acta</i> 51: 1345-1363.
Berelson et al. (1990)
Berelson, W M; Hammond, D E; Johnson, K S (1990): Benthic fluxes and pore water studies from sediments of the central equatorial north Pacific: Nutrient diagenesis. <i>Geochim. Cosmochim. Acta</i> 54: 3001-3012.
Berelson et al. (2002)
Berelson, W. M., K. Johnson, K. Coale, and H.-C. Li (2002): Organic matter diagenesis in the sediments of the San Pedro Shelf along a transect affected by sewage effluent: <i>Cont. Shelf Res.</i> , v. 22, p. 1101-1115.
Bickert and Wefer (1996)
Bickert, T., and G. Wefer (1996): Late Quaternary deep water circulation in the South Atlantic: Reconstruction from carbonate Dissolution and benthic stable isotopes, <i>in</i> G. Wefer, W. H. Berger, G. Siedler, and D. Webb, eds., <i>The South Atlantic: Present and Past Circulation</i> , v. 103:599-620, Springer Verlag Berlin.
Birgisdottir (1991)
Birgisdottir, L (1991): Die palaeo-ozeanographische Entwicklung der Islandsee in den letzten 550.000 Jahren. <i>Berichte aus dem Sonderforschungsbereich 313</i> , Christian-Albrechts-Universität Kiel 34: 186.
Bonn (1995)
Bonn, W, J (1995): Biogenes Opal und biogenes Barium als Indikatoren für spätquartäre Produktivitätsänderungen am antarktischen Kontinentalhang, atlantischer Sektor. <i>Reports on Polar Research</i> , Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 180: 186.
Boucein et al. (1999)
Boucein, B; Fahl, K; Siebold, M & Stein, R (1999): Quantity and quality of organic carbon in surface sediments of the Ob and Yenisei estuaries and adjacent coastal areas: marine productivity vs. terrigenous input. <i>In: Matthiessen, J; Stepanets, O V; Stein, R; Fütterer, D K & Galimov, E M (eds), The Kara Sea Expedition of RV Akademik Boris Petrov 1997: First Results of a Joint Russian-German Pilot Study</i> , <i>Rep on Polar Research</i> , 300: 116-126.
Braun (1997)
Braun, B (1997): Rekonstruktion glaziomarer Sedimentationsprozesse am Kontinentalrand des westlichen Bellingshausenmeeres, Master thesis, Alfred .Wegener Institut for Polar and Marine Research, Bremerhaven & Geologisches Institut der Bayerischen Julius-Maximilians Universität, Würzburg, Germany, unpublished: 83.
Brehme (1992)
Brehme, I (1992): Sedimentfazies und Bodenwasserstrom am Kontinentalhang des nordwestlichen Weddelmeeres, <i>Reports on Polar Research</i> , Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 110: 127.

Bremner and Willis (1993)
Bremner, J. M., and P. C. Willis (1993): Mineralogy and geochemistry of the clay fraction of sediments from the Namibian continental margin and the adjacent hinterland: <i>Marine Geology</i> , v. 115, p. 85-116.
Cai and Reimers (1995)
Cai, W-J and Reimers, C E (1995): Benthic oxygen flux, bottom water oxygen concentration and core top organic carbon content in the deep northeast Pacific Ocean. <i>Deep Sea Research</i> 42(10): 1681-1699.
Calvert and Price (1983)
Calvert, S. E., and N. B. Price (1983): Geochemistry of Namibian shelf sediments, <i>in</i> J. Thiede, and E. Suess, eds., Coastal upwelling: Its sedimentary record: New York, Plenum Press, 337-375.
CLIMAP (1976):
CLIMAP, P M (1976): The surface of the ice-age earth. <i>Science</i> 191: 1131 - 1137.
Clough et al. (1997)
Clough, L M; Ambrose jr, W G; Cochran, J K; Barnes, C; Renaud, P E & Aller, R C (1997): Infaunal density, biomass and bioturbation in the sediments of the Arctic Ocean. <i>Deep-Sea Research II</i> , 44(8): 1683-1704.
Colombo et al. (1996)
Colombo, J C; Silverberg, N; Gearing, J N (1996): Biogeochemistry of organic matter in the Laurentian Trough, II. Bulk composition of the sediments and relative reactivity of major components during early diagenesis. <i>Mar. Chem.</i> 51: 295-314.
Cordes (1990)
Cordes, D (1990): Sedimentologie und Paläomagnetik an Sedimenten der Maudkuppe (Nördliches Weddellmeer). <i>Reports on Polar Research</i> , Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 71: 158.
Correns (1937)
Correns, C W (1937): Die Sedimente des Äquatorialen Atlantischen Ozeans. <i>Notgemeinschaft der Deutschen Wissenschaft</i> , Verlag Walter de Gruyter & Co, Berlin und Leipzig: 298.
Cough et al. (1993)
Cough, M A; Fauzi, R; Mantoura, C; Preston, M (1993): Terrestrial plant biopolymers in marine sediments. <i>Geochim.Cosmochim.Acta</i> 57: 945-964.
Cowie et al. (1999)
Cowie, G. L., S. E. Calvert, T. F. Petersen, H. Schulz, and U. von Rad (1999): Organic content and preservational controls in surficial shelf and slope sediments from the Arabian Sea (Pakistan Margin): <i>Marine Geology</i> , v. 161, p. 23-38.
Cranston (1997)
Cranston, R. E. (1997) Organic carbon burial rates across the Arctic Ocean from 1994 Arctic Ocean Section expedition: <i>Deep-Sea Res. II</i> , v. 44, p. 1705-1723.
Crawford et al. (1996)
Crawford, K; Kuhn, G & Hambrey, M J (1996): Changes in the character of glaciomarine sedimentation in the southwestern Weddell Sea, Antarctica: evidence from the core PS1423-2. <i>Ann. Glaciol.</i> , 22: 200-204.
Curry and Lohmann (1983)
Curry, W B & Lohmann, G P (1983): Carbon isotopic changes in benthic foraminifera from the Western South Atlantic: Reconstruction of glacial abyssal circulation patterns. <i>Quaternary Research</i> , 18: 218-235.

Cwienk (1986)
Cwienk, D S (1986): Recent and Glacial Age Organic Carbon and Biogenic Silica Accumulation in Marine Sediments. Master of Science Thesis in Oceanography, University of Rhode Island: 237.
De Rijk et al. (1999)
De Rijk, S; Troelstra, S R & Rohling, E J (1999): Benthic foraminiferal distribution in the Mediterranean Sea. <i>Journal of Foraminiferal Research</i> , 29: 93-103.
Devol and Christensen (1993)
Devol, A H and Christensen, J P (1993): Benthic fluxes and nitrogen cycling in sediments of the continental margin of the eastern North Pacific. <i>J. Mar. Res.</i> 51: 345-372.
De Wit et al. (1997)
De Wit, R. d., J.-C. Relexans, T. Bouvier, and D. J. W. Moriarty (1997): Microbial respiration and diffusive uptake of deep-sea sediments in the Southern Ocean (ANTARES-I cruise): <i>Deep-Sea Research II</i> , v. 44, 5, p. 1053-1068.
DSDP; JOI; NGDC (1989)
DSDP; JOI; NGDC (1989): Marine Geological and Geophysical Data from the DEEP SEA DRILLING PROJECT, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Environmental Satellite, Data & Information Service, National Geophysical Data Center, CD-ROM Data Set.
Ebel et al. (1999)
Ebel, T; Melles, M & Niessen, F (1999): Laminated Sediments from Levinson-Lessing Lake, Northern Central Siberia - A 30,000 year record of environmental history?, In: Kassens, H; Bauch, H A; Dmitrenko, I A; Eicken, H; Hubberten, H-W; Melles, M; Thiede, J & Timokhov, L A (eds.), <i>Land-ocean systems in the Siberian Arctic: dynamics and history</i> . (Springer-Verlag Berlin Heidelberg): 425-436.
Elderfield et al. (1981)
Elderfield, H., Luedke, N A.; Mc Caffrey; RJ.; Bender, M. (1981): Benthic flux studies in Narragansett Bay, <i>American Journal of Science</i> , 281, 768-787.
Emerson et al. (1985)
Emerson, S; Fischer, K; Reimers, C; Heggie, D (1985): Organic carbon dynamics and preservation in deep-sea sediments. <i>Deep-Sea Research</i> 32: 1 - 21.
Emerson and Hedges (1988)
Emerson, S and Hedges, J I (1988): Processes controlling the oceanic carbon content of open ocean sediments. <i>Paleoceanography</i> 3: 621 - 634.
Epping et al. (2002)
Epping, E. H. G., v. d. Zee, C. Soeteart, K., Helder, W. (2002): On the oxidation and burial of organic carbon in sediments of the Iberian Margin and Nazare Canyon (NE Atlantic). <i>Progress in Oceanography</i> , 52: 399-431.
Etcheber et al. (1999)
Etcheber, H., J.-C. Relexans, M. Beliard, O. Weber, R. Buscail, and S. Heussner (1999): Distribution and quality of sedimentary organic matter on the Aquitanian margin (Bay of Biscay): <i>Deep Sea Res. II</i> , 46: 2249-2288.
Emelyanov and Romankevich (1979):
Emelyanov, E. M. and Romankevich, E A (1979): <i>Geochemistry of the Atlantic Ocean: Organic carbon and Phosphorous</i> (in Russian). Nauka, Moscow, p.220.
Fahl and Stein (1999)
Fahl, K & Stein, R (1999): Biomarkers as organic-carbon-source and environmental indicators in the Late Quaternary Arctic Ocean: Problems and Perspectives. <i>Marine Chemistry</i> , 63(3-4): 293-309

Fahl et al. (2002)
Fahl, K. C., H; Erlenkeuser, H; Hanssen, H; Hölemann, J; Kassens, H; Knickmeier, K; Kosobokova, K; Kunz-Pirrung, M; Lindemann, F; Markhaseva, E; Lischka, S; Petryashov, V; Piepenburg, D; Stein, R & Tuschling, K (2002): Sources and pathways of organic carbon in the modern Laptev Sea (Arctic Ocean): implications from biological, geochemical and geological data, <i>Berichte zur Polarforschung</i> , Alfred-Wegener Institute Bremerhaven: 193-205.
Finney et al. (1988)
Finney, M; H. Lyle; GR. Heath (1988): Sedimentation at Manop site H (eastern equatorial Pacific) over the past 400,000 years: climatically induced redox variations and their effects on transition metal cycling, <i>Paleoceanography</i> 3:169-189.
Fontugne et al. (1989)
Fontugne, M R; Paterne, M; Calvert, S E; Murat, A; Guichard, F & Arnold, M (1989): Adriatic deep water formation during the Holocene: implication for the reoxygenation of the deep eastern Mediterranean Sea. <i>Paleoceanography</i> , 4: 199-206.
Frenz et al. (2004)
Frenz, M; Henrich, R; Höppner, R; Stuut, J-B & Wagner, T (2004): Surface sediment bulk geochemistry and grain-size composition related to oceanic circulation along the South American continental margin in the Southwest Atlantic. In: Wefer, G; Mulitza, S & Rathmeyer, V (eds.), <i>The South Atlantic in the Late Quaternary: Reconstruction of Material Budgets and Current Systems</i> , Springer, Berlin, Heidelberg, New York: pp. 347-373.
Fütterer et al. (1988)
Fütterer, D K; Grobe, H & Grünig, S (1988): Quaternary sediment patterns in the Weddell Sea: relations and environmental conditions. <i>Paleoceanography</i> , 3/5: 551-561, awi55.
Gaspar et al. (1996)
Gaspar, L; Abrantes, F & Lebreiro, S (1996): Nature and origin of carbonates within the Portuguese margin sediments for the last 30 kyrs. ENAM 1 meeting, Geilo, Norway.
Gingele and Leipe (1997)
Gingele, F X & Leipe, T (1997): Distribution and enrichment of redox-sensitive metals in Baltic Sea sediments. <i>Baltica</i> , 11.
Gingele et al. (1997)
Gingele, F X; Kuhn, G; Maus, B; Melles, M & Schöne, T (1997): Holocene ice retreat from the Lazarev Sea shelf, East Antarctica. <i>Continental Shelf Research</i> , 17: 137-163, SFB261 No 106.
Giordani et al. (2002)
Giordani, P; Helder, W; Koning, E; Miserocchi, S; Danovaro, R; Malaguti, A. (2002): Gradients of benthic-pelagic coupling and carbon budgets in the Adriatic and Northern Ionian Sea. <i>Journal of Marine Systems</i> 33-34: 365-387.
Giraudeau et al. (2001)
Giraudeau, J; Meyers, P A; Christensen, B A (2001): Accumulation of organic and inorganic carbon in Pliocene-Pleistocene sediments along the SW African margin. <i>Marine Geology</i> 2966: 1-21.
Gordon et al. (2001)
Gordon, E S; Goni, M A; Roberts, Q N; Kineke, G C; Allison, M A (2001): Organic matter distribution and accumulation on the inner Louisiana shelf west of the Atchafalaya River. <i>Continental shelf research</i> 21: 1691-1721.

Gorshkova (1962)
Gorshkova, T I (1962): Organic matter in bottom sediments of Norwegian Sea and conditions of its deposition (in Russian). Vinogradova, L G & Fedosova, M (eds.), Researches on Programm of International geophysical year, Proceedings of VNIRO. Pischeprom. Moscow. L. G. Vinogradova and M. Fedosova: 38-57.
Grandel et al. (2000)
Grandel, S; Rickert, D; Schlüter, M; Wallmann, K (2000): Pore-water distribution and quantification of diffusive benthic fluxes of silicic acid, nitrate, and phosphate in surface sediments of the deep Arabian Sea. Deep Sea Research II, 47 (14): 2707-2734
Grobe (1986)
Grobe, H (1986): Spätpleistozäne Sedimentationsprozesse am antarktischen Kontinentalhang vor Kapp Norvegia, östliche Weddel See. Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 27: 121.
Grobe, H (2000)
Grobe, H (2000): Sedimentological data from high latitude oceans, PANGAEA - Network for Geological and Environmental Data (Alfred Wegener Institute for Polar and Marine Research & World Data Center for Marine Environmental Sciences, Germany), unpublished.
Grobe et al. (1993)
Grobe, H; Fütterer, D K; Hubberten, H-W; Kuhn, G & Mackensen, A (1993): Zur Entwicklung der spätquartären Sedimentfazies im Südpolarmeer, Zeitschrift der Deutschen Geologischen Gesellschaft, 144: 330-351.
Grobe and Mackensen (1992)
Grobe, H & Mackensen, A (1992): Late Quaternary climatic cycles as recorded in sediments from the Antarctic continental margin. In: Kennett, J P & Warnke, D (eds.), The Antarctic Paleoenvironment: a perspective on global change, Antarctic Research Series, 56: 349-376, SFB261 No 39, awi546.
Grundmanis and Murray (1982)
Grundmanis, V & Murray, J W (1982): Aerobic respiration in pelagic marine sediments. Geochim. Cosmochim. Acta 46: 1101-1120.
Grünig (1991)
Grünig, S (1991): Quartäre Sedimentationsprozesse am Kontinentalhang des Süd-Orkney-Plateaus im nordwestlichen Weddellmeer (Antarktis). Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 75: 196.
Haase (1986)
Haase, G M (1986): Glaciomarine sediments along the Filchner/Ronne ice shelf, southern Weddell Sea - first results of the 1983/1984 Antarktis-II/4 expedition. Marine Geology, 72: 241-258.
Hales and Emerson (1996)
Hales, B and Emerson, S (1996): Calcite dissolution in sediments of the Ontong-Java Plateau: In situ measurements of pore water O ₂ and pH. Global Biogeochemical Cycles 10(3): 527 - 541.
Hales et al. (1994)
Hales, B; Emerson, S; Archer, D (1994): Respiration and dissolution in the sediments of the western North Atlantic: estimates from models of in situ microelectrode measurements of porewater oxygen and pH. Deep-Sea Research 41(4): 695 - 719.

Harwart et al. (1999)
Harwart, S; Hagedorn, B; Melles, M & Wand, U (1999): Lithological and biochemical properties in sediments of Lama Lake as indicators for the late Pleistocene and Holocene ecosystem development of the southern Taymyr Peninsula, Central Siberia. <i>Boreas</i> , 28: 167-180.
Hathaway and Watkins (1971)
Hathaway, J C and Watkins, N D (1971): Continental Margin program, Atlantic Coast of the United States, Woods Hole, Oceanographic Institution, Ref. No. 71-15: 496.
Hays et al. (1976)
Hays, J D; Lozano, J A; Shackleton, N; Irving, G (1976): Reconstruction of the Atlantic and western Indian Ocean sectors of the 18.000 B.P. Antarctic Ocean. Investigations of Late Quaternary Paleoceanography and Paleoclimate. R. M. Cline and J. D. Hays: 336-372.
Hebbeln (1991)
Hebbeln, D (1991): Spätquartäre Stratigraphie und Paläozeanographie in der Fram-Strasse. <i>Berichte, Fachbereich Geowissenschaften, Universität Bremen</i> , 22: 174.
Hebbeln and Berner (1993)
Hebbeln, D & Berner, H (1993): Surface sediment distribution in the Fram Strait, <i>Deep-Sea Research</i> , 40: 1731-1745.
Hebbeln et al. (2001)
Hebbeln et al. (2001): PUCK: Report and preliminary results of R/V Sonne Cruise SO156, Valparaiso (Chile) - Talcahuano (Chile). March 29 - May 14, 2001. <i>Berichte, Fachbereich Geowissenschaften, Universität Bremen</i> .
Heggie et al. (1987)
Heggie, D; Maris, C; Hudson, A; Dymond, J; Beach, R & Cullen, J (1987): Organic carbon oxidation and preservation in NW Atlantic continental margin sediments, Weaver, PPE & Thomson J (eds) <i>Geology and Geochemistry of Abyssal Plains</i> . Geological Society Special Publication, 31: 215-236.
Holler (1988)
Holler, P (1988): Sedimentäre Rutschmassen in der Tiefsee. <i>Berichte-Reports, Geologisch-Paläontologisches Institut und Museum, Christian-Albrechts-Universität, Kiel</i> , 23: 141.
Holz (2001)
Holz, C (2001): Glazialmarine Sedimentationsprozesse am Kontinentalhang des westlichen Weddellmeeres. Master thesis, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven & Institut für Geowissenschaften, Christian-Albrechts-Universität, Kiel: 70 pp
Honjo et al. (1982)
Honjo, S; Manganini, S; Cole, J (1982): Sedimentation of biogenic matter in the deep ocean. <i>Deep Sea Res.</i> 29(5): 609-625.
Hübner (1995)
Hübner, M (1995): Verteilung biogeochemischer Komponenten und deren Korngrößenabhängigkeit in Oberflächensedimenten des nordostbrasilianischen Kontinentalrandes zwischen Belem und Recife. Thesis, Univ. Hamburg: 92.
Hüls (2000)
Hüls, M (2000): Millennial-scale SST variability as inferred from planktonic foraminifera sensus counts in the western subtropical Atlantic. GEOMAR Report, GEOMAR Research Center for Marine Geosciences, Christian Albrechts University in Kiel, 95: 118.

Huh et al. (1987)
Huh, C-A; Zahnke, D L; Small, L F; Noshkin, V E (1987): Budgets and behaviours of uranium and thorium series isotopes in Santa Monica Basin sediments. <i>Geochim. Cosmochim. Acta</i> 51: 1743 - 1754.
Hulth et al. (1996)
Hulth, S; Hall, P O J; Blackburn, T H & Landén, A (1996): Arctic sediments (Svalbard): pore water and solid phase distributions of C, N, P and S. <i>Polar Biology</i> , 16: 447-462.
Hurd (1973)
Hurd, D C (1973): Interactions of biogenic opal, sediment and seawater in the Central Equatorial Pacific. <i>Geochimica et Cosmochimica Acta</i> 37: 2257-2282.
Isla et al. (2002)
Isla, E; Masque, P; Palanques, A; Sanchez-Cabeza, J A; Bruach, J M; Guillen, J; Puig, P (2002): Sediment accumulation rates and carbon burial in bottom sediment in a high-productivity area: Gerlache Strait (Antarctica). <i>Deep Sea Res. II</i> , 49, 4-5 : 921-933.
Jahnke et al. (1989)
Jahnke, R. A., S. R. Emerson, C. E. Reimers, J. Schuffert, K. Ruttenger, and D. Archer (1989) Benthic Recycling of biogenic Debris in the Eastern Tropical Atlantic Ocean: <i>Geochim. Cosmochim. Acta</i> , v. 53: 2947-2960.
Jahnke et al. (1990a)
Jahnke, R A; Reimers, C E; Craven, D B (1990a): Intensification of recycling of organic matter at the sea floor near ocean margins. <i>Nature</i> 348: 50-54.
Jahnke et al. (1990b)
Jahnke, R A (1990b): Early diagenesis and recycling of biogenic debris at the seafloor, Santa Monica Basin, California. <i>Journal of Marine Research</i> 48: 413-436.
Jansen et al. (1984)
Jansen, J H F, Van Weering, TCE; Gieles, R; Van Iperen, J (1984): Middle and late quarterary oceanography and climatology of the Zaire-Congo fan and the adjacent eastern Angola Basin. <i>Neth.J.Sea Res.</i> 17, 2-4: 201-249.
Jennerjahn and Ittekkott (1997)
Jennerjahn, T C and Ittekkot, V (1997): Organic matter in sediments in the mangrove areas and adjacent continental margins of Brasil: I. Amino acids and hexosamins. <i>Oceanologica Acta</i> 20(2): 359-369.
Jeong et al. (1994)
Jeong, K S; Kang, J K; Chough S K (1994): Sedimentary processes and manganese nodule formation in the Korea Deep Odean Study (KODOS) area, western part of Clarion-Clipperton fracture zones, northeast equatorial Pacific. <i>Marine Geology</i> 122: 125-150.
Jorgensen and Des Marais (1990)
Jorgensen, B B & Des Marais, D J (1990). The diffusive boundry layer of sediments: Oxygen microgradients over a microbial mat. <i>Limnol. Oceanogr.</i> 35: 1343-1355.
Kassens (1990)
Kassens, H (1990): Verfestigte Sedimentlagen und seismische Reflektoren: Ruehdiagenese und Palaeo-Ozeanographie in der Norwegischen See. <i>Berichte aus dem Sonderforschungsbereich 313, Christian-Albrechts-Universität, Kiel</i> , 24: 117.
Kienel and Kumke (2000)
Kienel, U & Kumke, T (2000): Estimating the variation pattern of the diatom distribution at Lake Lama, Central Siberia. <i>Ecology</i> , submitted.

Klump (1999)
Klump, J (1999): Biogenic barite as a proxy of paleoproductivity variations in the southern Peru-Chile Current. Berichte, Fachbereich Geowissenschaften, Universität Bremen, 139: 107.
Knies (1994)
Knies, J (1994): Spätquartäre Sedimentation am Kontinentalhang nordwestlich Spitzbergens. Der letzte Glazial/Interglazial-Zyklus. Thesis, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, unpublished: 95
Knies (1999)
Knies, J (1999): Spätquartäre Paläoumweltbedingungen am nördlichen Kontinentalrand der Barents- und Kara-See. Eine Multi-Parameter-Analyse. Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 304: 159.
Kolla et al. (1979)
Kolla, V; Biscaye, P E; Hanley, A F (1979): Distribution of quartz in late quaternary Atlantic sediments in relation to climate. Quaternary Research 11: 261-277.
Kroencke et al.(1994)
Kroencke, I; Tan, L K; Stein, R (1994): High benthic bacteria standing stock in deep Arctic basins. Polar Biol, 14: 423-428.
Kuhn (1997)
Kuhn, T; Bau, M; Blum, N; Halbach, P (1998): Origin of negative Ce anomalies in mixed hydrothermal-hydrogenetic Fe-Mn crusts from the Central Indian Ridge. Earth and Planetary Science Letters 163: 207-220.
Kulbe (1997)
Kulbe, T (1997): Die spätquartäre Klima- und Umweltgeschichte der Bunger-Oase, Ostantarktis. Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 254: 129.
Lackschewitz (1991)
Lackschewitz, K S (1991): Sedimentationsprozesse am aktiven mittelozeanischen Kolbeinsey Rücken (nördlich von Island). GEOMAR Report, GEOMAR Research Center for Marine Geosciences, Christian Albrechts University in Kiel, 9: 133.
Lackschewitz et al. (1994)
Lackschewitz, K. S., H. J. Wallrabe-Adams, and D. Garbe-Schönberg (1994): Geochemistry of surface sediments from the mid-oceanic Kolbeinsey Ridge, north of iceland: Marine Geology, v. 121, p. 105-119.
Lebreiro et al. (1997)
Lebreiro, S M; Moreno, J C; Abrantes, F F & Pflaumann, U (1997): Productivity and paleoceanographic implications on the Tore Seamount (Iberian Margin) during the last 225 kyr: Foraminiferal evidence. Paleoceanography, 12(5): 718-727.
Ledbetter (1985)
Ledbetter, M T (1985): Tephrochronology of marine tephra adjacent to central America. Geological Society of America Bulletin 96: 77-82.
Leipe et al. (1995)
Leipe, T; Neumann, T & Emeis, K-E (1995): Schwermetallverteilung in holozänen Ostseesedimenten. Geowissenschaften, 13 (12): 470-478.
Leong (1999)
Leng, C (1999): Rekonstruktion der spätquartären Sedimentationsprozesse am Nord-Scotia-Rücken mit Hilfe sedimentologischer und sedimentphysikalischer Untersuchungen. Master thesis, Geologisches Institut der Universität zu Köln, Alfred

Wegener Institut für Polar und Meeresforschung, Bremerhaven, unpublished.
Levitan et al. (1990)
Levitan, M. A., P. G. Ditrikh, et al. (1990): Bottom sediments in transocean section (Southern part of the Atlantic Ocean) (in Russian). Transactions of IOAN. Biological and geological research of bottom of southern part of the Atlantic Ocean: 172-184.
Lisitzin et al. (1975)
Lisitzin, A P; Emelyanov, E M; Ilyin, A V; Koshelev, B A; Litvin, V M; Likoshevitchus, L S; Senin, Y M; Svirenko, Y P; Soldatov, A V & Shurko, Y.I. (1975): Distribution of bottom sediments in the Atlantic Ocean. In: Lisitzin, A (ed.), Sedimentation in the Atlantic Ocean (in Russian), Kaliningradskaya Pravda, Kaliningrad and supplement: 323.
Lochte et al. (2000)
Lochte, K; Boetius, A; Gebruk, A; Helder, W; Jahnke, R; Pfannkuche, O; Rabouille, C; Schlüter, M; Shimmield, G; Sibuet, M; Soltwedel, T; Vetrov, A & Zabel, M (2000): Atlantic data base for exchange processes at the deep sea floor (ADEPD). Data collected and published through EU-project ADEPD (MAS3-CT97-0126-ADEPD) 1998/99. Institute for Baltic Sea Research, Warnemünde, Germany.
Lowry (1994)
Lowry, R K; Machin, P & Cramer, R N (1994): BOFS North Atlantic Data Set, Natural Environmental Research Council, British Oceanographic Data Centre, Merseyside, United Kingdom, incl CD: 140.
Luff et al. (2000)
Luff, R; Wallmann, K, Grandel, S; Schlüter, M (2000): Numerical modeling of benthic processes in the deep Arabian Sea. Deep Sea Research II 47(14): 3039-3072.
Lyle (1992)
Lyle, M (1992): Composition maps of surface sediments of the eastern tropical Pacific Ocean. Proc. Ocean Drill. Program Initial Rep. 138: 101-115.
Lyle et al. (1984)
Lyle, M; Heath, G R; Robbins, J M (1984): Transport and release of transition elements during early diagenesis: Sequential leaching of sediments from MANOP Sites M and H. Part I. pH 5 acetic acid leach. Geochim. Cosmochim. Acta 48: 1705-1715.
Lyle et al. (1992)
Lyle, M., R. Zahn, F. Prahl, J. Dymond, R. Collier, N. Pisias, and E. Suess (1992): Paleoproductivity and carbon burial across the California current: The multitracers transect, 42°N, <i>Paleoceanography</i> , 7(3), 251–272.
Lyle et al. (2000)
Lyle, M; Mix, A; Ravelo, A C; Andreasen, D; Heusser, L; Olivarez, A (2000): Millennial-Scale CaCO ₃ and Corg Events Along the Northern and Central California Margins: Stratigraphy and Origins. Proc. of ODP, Sc. Res. 167: 163-182.
Mackensen (1985)
Mackensen, A (1985): Verbreitung und Umwelt benthischer Foraminiferen in der Norwegischen See. Thesis, Christian-Albrechts-Universität, Kiel: 126.
Mackensen et al. (1994)
Mackensen, A; Grobe, H; Hubberten, H-W & Kuhn, G (1994): Benthic foraminiferal assemblages and the δ ¹³ C-signal in the atlantic sector of the Southern Ocean: glacial-to-interglacial contrasts. In: Zahn, R; Pederson, T F; Kaminiski, M A & Labeyrie, L (eds.), Carbon Cycling in the Glacial Ocean: Constraints on the Ocean's Role in Global Change, Springer-Verlag, Berlin, Heidelberg, 117: 105-144, SFB261 No 67, awi691.

Mackensen et al. (1990)
Mackensen, A; Grobe, H; Kuhn, G & Fütterer, D K (1990): Benthic foraminiferal assemblages from the eastern Weddell Sea between 68 and 73°S: distribution, ecology and fossilization potential. <i>Marine Micropaleontology</i> , 16: 241-283, SFB261 No 29, awi267.
Marienfeld (1991)
Marienfeld, P (1991): Holozäne Sedimentationsentwicklung im Scoresby Sund, Ost-Grönland. <i>Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven</i> , 96: 162.
Martens and Klump (1984)
Martens, C. S., and J. V. Klump (1984) Biogeochemical cycling in an organic-rich coastal marine basin 4. An organic carbon budget for sediments dominated by sulfate reduction and methanogenesis: <i>Geochim. Cosmochim. Acta</i> , 48: 1987 - 2004.
Maus (1993)
Maus, B (1993): Plio-Pleistozäne Sedimentation im Randbereich des Ritscher-Canyons, westlich des Gunnerus-Rückens (Ostantarktis). Thesis, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven/Freiburg, unpublished: 123.
Meinecke (1992)
Meinecke, G (1992): Spätquartäre Oberflächenwassertemperaturen im östlichen äquatorialen Atlantik. <i>Berichte, Fachbereich Geowissenschaften, Universität Bremen</i> , 29: 181.
Meggers et al. (2002)
Meggers, H; Freudenthal, T; Nave, SO; Targarona, J; Abrantes, F; Helmke, P (2002): Assessment of geochemical and micropaleontological sedimentary parameters as proxies of surface water properties in the Canary Islands region <i>Deep-Sea Research II</i> , 49 (17), 3631-3654.
Melles (1991)
Melles, M (1991): Paläoglazilogie und Paläozeanographie im Spätquartär am Kontinentalrand des südlichen Weddellmeeres, Antarktis. <i>Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven</i> , 81: 190.
Mix and Fairbanks (1985)
Mix, A C and Fairbanks, R G (1985): North Atlantic surface-ocean control of Pleistocene deep-ocean circulation, <i>Earth and Planetary Science Letters</i> . 73: 231-243.
Molina-Cruz (1976)
Molina-Cruz, A (1976): Paleo-oceanography of the subtropical southeastern Pacific during late Quaternary: a study of radiolaria, opal and quartz contents of deep-sea sediments. Corvallis, Oregon State University: 178.
Mollenhauer et al. (2004)
Mollenhauer, G; Schneider, R R; Jennerjahn, T; Müller, P J & Wefer, G (2004): Organic carbon accumulation in the South Atlantic Ocean: Its modern, mid-Holocene and Last Glacial basin-wide distribution. <i>Global and Planetary Change</i> , 40: 249-266.
Mollenhauer et al. (2002)
Mollenhauer, G; Schneider, R R; Müller, P J; Spieß, V; Wefer, G (2002): Glacial/Interglacial Variability in the Benguela Upwelling System: Spatial Distribution and Budgets of Organic Carbon Accumulation. <i>Global Biogeochemical Cycles</i> , 204: 139.
Mueller (1995)
Mueller, C (1995): Spätquartäre Sedimentationsprozesse in der östlichen Framstrasse. Master thesis, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven.

84.
Mueller et al. (1994)
Mueller, P J; Schneider, R; Ruhland, G (1994): Late Quaternary PCO ₂ Variations in the Angola Current: Evidence from Organic Carbon d ¹³ C and Alkenone Temperatures. In: Zahn, R. and et al. (eds), Carbon Cycling in the Glacial Ocean: Constraints on the Ocean's Role in Global Change, Springer-Verlag, Berlin, Heidelberg. 117: 343-361.
Mueller and Suess (1979)
Mueller, P & Suess, E (1979): Productivity, sedimentation rate, and sedimentary organic matter in the oceans. I.- Organic matter preservation. Deep-Sea Research, 26: 1347-1362.
Mueller-Lupp et al. (2000)
Mueller-Lupp, T; Bauch H A; Erlenkeuser, H; Hefter, J; Kassens, H & Thiede, J (2000): Changes in the deposition of terrestrial organic matter on the Laptev Sea shelf during the Holocene: evidence from stable carbon isotopes. International Journal of Earth Sciences, 89: 563-568.
Murray and Kuivila (1990)
Murray, J W & Kuivila, K M (1990): Organic matter diagenesis in the northeast Pacific: transition from aerobic red clay to suboxic hemipelagic sediments. Deep-Sea Res. 37: 59-80.
Murray and Leinen (1993)
Murray, R W & Leinen, M (1993): Chemical transport to the seafloor of the equatorial Pacific Ocean across a latitudinal transect at 135°W: Tracking sedimentary major, trace, and rare earth element fluxes at the Equator and the Intertropical Convergence Zone. Geochim. Cosmochim. Acta 57: 4141-4163.
Murray et al. (1995)
Murray, R. W., M. Leinen, D. W. Murray, A. C. Mix, and C. W. Knowlton (1995): Terrigenous Fe input and biogenic sedimentation in the glacial and interglacial equatorial Pacific Ocean: Global Biogeochemical Cycles, v. 9, p. 667 - 684.
Murray (1987)
Murray, D. W. (1987): Spatial and temporal variations in sediment accumulation in the central tropical Pacific: Ph.D. dissertation, 343pp., Oreg. State Univ., Covallis, Oreg.
Nam (1997)
Nam, S-II (1997): Late Quaternary glacial history and paleoceanographic reconstructions along the East Greenland continental margin: Evidence from high-resolution records of stable isotopes and ice-rafted debris. Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 241: 250.
Nothold (1998)
Nothold, H (1998): Die Auswirkungen der "NorthEastWater"-Polynya auf die Sedimentation vor NO-Grönland und Untersuchungen zur Paläo-Ozeanographie seit dem Mittelweichsel.
Prahl et al. (1989)
Prahl, F. G., L. A. Muehlhausen, and M. Lyle (1989): An organic geochemical assessment of oceanographic conditions at MANOP site C over the past 26.000 years: Paleoceanography, v. 4, p. 495 - 510.
Obdyke and Foster (1970)
Obdyke, N., and J. H. Foster (1970): Paleomagnetism of cores from the North Pacific: Geological Society of America Memior, v. 126, p. 83-119.
OMEX-I (1997)

OMEX-I (1997): Ocean Margin Exchange, OMEX-I project data set, Natural Environmental Research Council, published by British Oceanographic Data Center, Bidston Observatory, Birkenhead, Merseyside.
Ott (1997)
Ott, G (1997): Stabile Isotope im Spätquartär des Atlantisch-Indischen RYckens im Südpolarmeer Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, data report, PANGAEA - Network for Geological and Environmental Data.
Ovalle et al. (1999)
Ovalle, A.R.C., C.E. Rezende, C.E.V. Carvalho, T.C. Jennerjahn & V. Ittekkot (1999). Biogeochemical characteristics of coastal waters adjacent to small river-mangrove systems, East Brazil.- In <i>Sedimentation process at the East and Northeast Brazil Shelf</i> , (eds. B. Knoppers & W. Ekau), Geo-Mar. Lett. 19 (3): 179-185.
Paetsch (1991)
Paetsch, H (1991): Sedimentation im Europäischen Nordmeer. Berichte aus dem Sonderforschungsbereich 313, Christian-Albrechts-Universität, Kiel, 29: 102.
Pagels (1991)
Pagels, U (1991): Sedimentologische Untersuchungen und Bestimmungen der Karbonatlösung in spätquartären Sedimenten des oestlichen arktischen Ozeans. GEOMAR Report, GEOMAR Research Center for Marine Geosciences, Christian Albrechts University in Kiel, 10: 106.
Pavlidis et al. (1986)
Pavlidis, Yu A; Ionin, A S; Scherbakov, F A; Dunaev, N N & Nikiforov, S L (1986): Late Quaternary history as a predicting base of future changing (in Russian). Moscow: GEOS: 187.
Pedersen et al. (1986)
Pedersen, T F; Vogel, J S; Southon, J R (1986): Copper and manganese in hemipelagic sediments at 21°N, East Pacific Rise: Diagenetic contrasts. Geochim. Cosmochim. Acta 50: 2019 - 2031.
Pedersen et al. (1992)
Pedersen, T. F., G. B. Shimmield, and N. B. Price (1992): Lack of enhanced preservation of organic matter in sediments under the oxygen minimum on the Oman Margin: Geochim. Cosmochim. Acta, 56, 545 - 551.
Prakasch Babu et al.(2002)
Prakasch Babu, C; Brumsack, H-J; Schnetger, B; Böttcher, M E (2002): Barium as a productivity proxy in continental margin sediments:a study from the eastern Arabian Sea. Marine Geol, 162: 91-103.
Prell et al. (1980)
Prell, W L; Hutson, W H et al. (1980): Surface circulation of the Indian Ocean during the last glacial maximum, approximately 18.000 yr. B.P. Quaternary Research 14: 309-336.
Rabouille and Gaillard (1991)
Rabouille C and J.-F. Gaillard (1991): A coupled model representing deep sea organic carbon mineralization and oxygen consumption in surficial sediments. Journal of Geophysical Research, 96, 2761-2776.
Rao and Berner (1993)
Rao, J L & Berner, R A (1993): Phosphorus dynamics in the Amazon River and estuary. Chemical Geol. 107: 397-400.

Rea et al. (1991)
Rea, D K; Pisias, N G & Newberry, T (1991): Late Pleistocene paleoclimatology of the central equatorial Pacific: flux patterns of biogenic sediments. <i>Paleoceanography</i> , 6: 227-244.
Reimers et al. (1986)
Reimers, C.E. and K.L. Smith (1986): Reconciling measured and predicted fluxes of oxygen across the deep sea sediment-water interface. <i>Limnology and Oceanography</i> , 31, 305-318.
Reimers (1987)
Reimers, C E (1987): An in situ microprofiling instrument for measuring interfacial pore water gradients: methods and oxygen profiles from the North Pacific Ocean. <i>Deep-Sea Research</i> 34 (12): 2019-2035.
Reimers et al. (1992)
Reimers, C E; Jahnke, R H, McCorkle, D C (1992): Carbon fluxes and burial rates over the continental slope and rise off central California with implications for the global carbon cycle. <i>Glob. Biogeochem. Cycles</i> 6: 199-224.
Relexans et al. (1996)
Relexans, J-C; Deming, J; Dinet, A; Gaillardts, J-F; Sibuet, M (1996): Sedimentary organic matter and micro-meiobenthos with relation to trophic conditions in the tropical northeast Atlantic. <i>Deep-Sea Research</i> 43(8): 1343-1368.
Romankevich and Vetrov (2001)
Romankevich, E A and Vetrov A A (2001): Cycle of carbon in the Russian Arctic seas. <i>Nauka Press (in Russian)</i> : 302.
Rosenfeld (1981)
Rosenfeld, J K (1981): Nitrogen diagenesis in Long Island Sound sediments. <i>American Journal of Science</i> 281: 436-462.
Rosenthal et al. (1995)
Rosenthal, Y; Boyle, E A, Labeyrie, L; Oppo, D (1995): Glacial enrichments of authigenic Cd and U in subantarctic sediments: a climatic control on the elements oceanic budget? <i>Paleoceanography</i> 10(3): 395-413.
Rühlemann et al. (1999)
Rühlemann, C; Müller, P J; Schneider, R R (1999): Organic Carbon and Carbonate as Paleoproductivity Proxies: Examples from High and Low Productivity Areas of the Tropical Atlantic. In: G. Fischer and G. Wefer (eds.), <i>Use of Proxies in Paleoceanography: examples from the South Atlantic</i> , Springer Verlag Berlin Heidelberg New York: 315-344.
Sanders et al. (1965)
Sanders, H. L; Hessler, R R et al. (1965): A introduction to the study of deep-sea benthic faunal assemblages along the Gay Head-Bermuda transect. <i>Deep Sea Res.</i> 12: 845-867.
Sarkar et al.(1993)
Sarkar, A; Bhattacharya, S K et al. (1993): Geochemical evidence for anoxic deep water in the Arabian Sea during the last glaciation. <i>Geochim. Cosmochim. Acta</i> 57: 1009-1016.
Sarnthein et al.(1994)
Sarnthein, M; Winn, K; Jung, S J A; Duplessy, J C; Labeyrie, L D; Erlenkeuser, H & Ganssen, G (1994): Changes in east Atlantic deepwater circulation over the last

30,000 years: Eight time slice reconstructions. <i>Paleoceanography</i> , 9 (2): 209-267.
Schlünz and Schneider (2000)
Schlünz, B S; Schneider, R R (2000): Transport of terrestrial organic carbon to the oceans by rivers: re-estimating flux- and burial- rates. <i>International Journal of Earth Sciences</i> 88: 599-606.
Schlüter (1990)
Schlüter, M (1990): Zur Frühdiagenese von organischem Kohlenstoff und Opal in Sedimenten des südlichen und östlichen Weddellmeeres. <i>Geochemische Analyse und Modellierung. Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven</i> , 73: 156.
Schmiedl (1990)
Schmiedl, G (1990): Quartäre Sedimentationsprozesse in der Tiefsee des Riiser-Larsenmeeres westlich des Gunnerus-Rückens, Ost-Antarktis. Thesis, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, unpublished: 113.
Schmiedl et al. (1997)
Schmiedl, G; Mackensen, A & Müller, P (1997): Recent benthic foraminifera from the eastern South Atlantic Ocean: dependence on food supply and water masses. <i>Marine Micropaleontology</i> , 32: 249-287, SFB261 No 148.
Schubert (1995)
Schubert, C J (1995): Organischer Kohlenstoff in spätquartären Sedimenten des Arktischen Ozeans: Terrigener Eintrag und marine Produktivität. <i>Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven</i> , 177: 178.
Seibold (1972)
Seibold, E (1972): Cruise 25/1971 of R.V." Meteor": Continental Margin of West-Africa. General Report and preliminary results, "Meteor" Forschungsergebnisse C: 17-38.
Sholkovitz (1973)
Sholkovitz, E (1973): Interstitial water chemistry of the Santa Barbara Basin sediments. <i>Geochim. et Cosmochim. Acta</i> 37: 2043-2073.
Sirocko (1989)
Sirocko, F (1989): Zur Akkumulation von Staubsedimenten im nördlichen Indischen Ozean; Anzeiger der Klimageschichte Arabiens und Indiens (Accumulation of ealioan sediments in the northern Indian Ocean; record of the climatic history of Arabia and India). <i>Berichte-Reports, Geologisch-Paläontologisches Institut und Museum, Christian-Albrechts-Universität, Kiel (in german)</i> , 27: 185.
Sirocko (1994)
Sirocko, F (1994): Abrupt change in monsoonal climate: evidence from the geochemical composition of Arabian Sea sediments. <i>Habilitation Thesis, Kiel University</i> .
Sirocko et al. (2000)
Sirocko, F; Garbe-Schönberg, D; Devey, C (2000): Processes controlling trace element geochemistry of Arabian Sea sediments during the last 25,000 years. <i>Global and Planetary Change</i> 26: 217-303.
Smith (1987)
Smith, K L j. (1987): Food energy supply and demand: A discrepancy between particulate organic carbonflux and sediment community oxygen consumption in the

deep ocean. <i>Limn.Oceanogr.</i> 32(1): 201-220.
Smith and Baldwin (1984)
Smith Jr, K L & Baldwin, R J (1984): Seasonal fluctuations in deep-sea sediment community oxygen consumption: central and eastern North Pacific. <i>Nature</i> 307 (5952): 624-626.
Smith et al. (1978)
Smith, K L J; White, G A; Laver, M B; Haugsness, J A (1978): Nutrient exchange and oxygen consumption by deep-sea benthic communities: Preliminary in situ measurements. <i>Limnol. Oceanogr.</i> 23(5): 997-1005.
Sokolowski (1986)
Sokolowski, B F (1986): Vergleichende sedimentologisch-geochemische Untersuchungen an marinen Oberflächensedimenten der Deutschen Bucht (Nordsee), der Lübecker-Mecklenburger Bucht (Ostsee) und des Weddell-Meer (Antarktis). Beiträge zur Meerestechnik, Arbeitsgruppe Meerestechnik und Marine Mineralstoffe, Technische Universität Clausthal, 10: 192.
Stein (1996)
Stein, R (1996): Organic-carbon and carbonate distribution in surface sediments from the eastern central Arctic Ocean and the Eurasian continental margin: sources and pathways. In: Stein, R; Ivanov, G I; Levitan, M A & Fahl, K (eds), Surface-sediment composition and sedimentary processes in the central Arctic Ocean and along the Eurasian Continental Margin, Rep on Polar Research, 212: 243-267.
Stein and Fahl (2000)
Stein, R & Fahl, K (2000): Holocene accumulation of organic carbon at the Laptev Sea continental margin (Arctic Ocean): sources, pathways, and sinks. <i>Geo-Marine Letters</i> , 20(1): 27-36.
Stein (1999)
Stein, R (1999): Zur neogenen Klimaentwicklung in Nordwest-Afrika und Paläo-Ozeanographie im Nordost-Atlantik, Berichte-Reports, Geologisch-Paläontologisches Institut und Museum, Christian-Albrechts-Universität, Kiel, 4,120.
Stein et al. (1994)
Stein, R; Grobe, H & Wahsner, M (1994): Organic carbon, carbonate, and clay mineral distributions in eastern central Arctic Ocean surface sediments. <i>Marine Geology</i> , 119: 269-285, awi702.
Steinberg (1987)
Steinberg, S M; Venkatesau, M I et al. (1987): Organic geochemistry of sediments from the continental margin off southern New England, USA. Part 1. Amino acids, carbohydrates and lignin. <i>Mar. Chem.</i> 21(3): 249-265.
Stevenson and Cheng (1972)
Stevenson, F J & Cheng, C N (1972): Organic geochemistry of the Argentine Basin in sediments; carbon-nitrogen relationships and Quaternary correlation. <i>Geochim. and Cosmochim.</i> 36(6): 653-671.
Stock (1997)
Stock, O (1997): Rekonstruktion der holozänen Entwicklungsgeschichte des Pjassinosees mit Hilfe von Seesedimenten. unpubl. Master Thesis, University Potsdam: 67.
Suits and Arthur (2000)
Suits, N. S., and M. A. Arthur (2000): Sulfur diagenesis and partitioning in Holocene Peru shelf and upper slope sediments: <i>Chemical Geology</i> , v. 163, p. 219-234.

Suthof et al. (2000)
Suthof, A; Jennerjahn, T C; Schäfer, P; Ittekkot, V (2000): Nature of organic matter in surface sediments from the Pakistan continental margin and the deep Arabian Sea: amino acids. <i>Deep Sea Research II</i> 47: 329-351.
Thiede (1979)
Thiede, J (1979): Wind regimes over the late Quaternary southwest Pacific Ocean. <i>Geology</i> 7: 259-262.
Thiedemann (1986)
Thiedemann, R. (1986): Verteilung von organischem Kohlenstoff in Oberflächensedimenten und örtliche Primärproduktion im äquatorialen Ostatlantik, 0-20°N, 15-25°W: Diplom thesis, Christian-Albrechts-Universität zu Kiel.
Thomson (1982)
Thomson, J (1982): Holocene sedimentation rates on the Hellenic outer Ridge: a comparison by ¹⁴ C and ²³⁰ Th excess methods. <i>Sedimentary geology</i> , 32: 99-110.
Thomson et al. (1984)
Thomson, J., T. R. S. Wilson, et al. (1984): "Non-steady state diagenetic record in eastern equatorial Atlantic sediments." <i>Earth Planet. Sci. Let.</i> 71: 23-30.
Tietjen (1971)
Tietjen, J H (1971): Ecology and distribution of deep-sea meiobenthos off North Carolina. <i>Deep Sea Res.</i> 18: 941-957.
Trotsuk and Marina (1988)
Trotsuk, V Y & Marina, M M (1988): Organic carbon in bottom sediments of the World Ocean (in Russian). Nauka, Moscow, : 176.
Van Cappellen and Qiu (1997)
Van Cappellen, P V and Qiu, L (1997): Biogenic silica dissolution in sediments of the Southern Ocean. I. Solubility. <i>Deep-Sea Research II</i> 44 (5): 1109-1128.
Vetrov and Korneeva (1998)
Vetrov, A & Korneeva, G (1998): This data set has been compiled for the ADEPD project. For detailed relation between data and reference see parameter 'Sample comment/reference' with this data set. For full text of citation see table 'Reference' in the data model of PANGAEA.
Vetrov et al. (1997)
Vetrov, A A; Romankevitch, E A; Benenson, M A (1997): Computer compilation of organic carbon distribution maps for bottom sediments. <i>Oceanology</i> 37(5): 651-656.
Vleet and Quinn (1979)
Vleet, E. S. v., and J. G. Quinn (1979): Diagenesis of Marine lipids in ocean sediments: <i>Deep Sea Res.</i> , v. 26, p. 1225-1236.
Vogt (1997)
Vogt, C (1997): Zeitliche und räumliche Verteilung von Mineralvergesellschaftungen in spätquartären Sedimenten des Arktische Ozeans und ihre Nützlichkeit als Klimaindikatoren während der Glazial/Interglazial-Wechsel. <i>Reports on Polar Research, Bremerhaven</i> , 251: 309.
Wagner (1993)
Wagner, T (1993): Organisches Material in pelagischen Sedimenten: Glaziale/interglaziale Variationen im Europäisches Nordmeer. <i>Berichte aus dem Sonderforschungsbereich 313, Christian-Albrechts-Universität, Kiel</i> , 41: 138.
Wagner (2000)
Wagner, B (2000): Holocene environmental history of East Greenland - evidence from lake sediments. <i>Reports on Polar Research, Alfred Wegener Institute for Polar and</i>

Marine Research, Bremerhaven, 358: 93.
Wagner and Dupont (1999)
Wagner, T & Dupont, L M (1999): Terrestrial Organic Matter in Marine Sediments: Analytical Approaches and Eolian-Marine Records in the Central Equatorial Atlantic. In: Fischer, G & Wefer, G (eds.), Use of Proxies in Paleoceanography - Examples from the South Atlantic, Springer, Berlin, Heidelberg: 547-574, SFB261 No 183.
Wagner et al. (2003)
Wagner, T., M. Zabel, L. Dupont, J. Holtvoeth, and C. J. Schubert (2003): Terrigenous Signals in Sediments of the Low Latitude Atlantic- Implications for Environmental Variations during the Late Quaternary, PartI: Organic Carbon, in G. Wefer, S. Mulitza, and V. Rathmeyer, eds., The South Atlantic in the Late Quaternary: Reconstruction of Material Budget and Current Systems, Springer.
Wagner and Melles (2000)
Wagner, B & Melles, M (2000): Holocene climatic and oceanic changes at East Greenland - evidences from seabird affected lake sediments on Raffles island. Boreas, in revision
Wallmann (2001)
Wallmann, K (2001): unpublished data, variabilities and outliers of data from the same location (i.e. between different cruises or different cores) are considered to be due to analyses, contact: kwallmann@geomar.de
Weber (1992)
Weber, M (1992): Spätquartäre Sedimentation am Kontinentalrand des südöstlichen Weddellmeeres, Antarktis. Reports on Polar Research, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, 109: 165.
Weber et al. (2000)
Weber, M E; von Stackelberg, U; Marchig, V; Wiedicke, M & Grupe, B (2000): Variability of surface sediments in the Peru Basin: dependence on water depth, productivity, bottom water flow, and seafloor topography. Marine Geology, 163 (1-4): 169-184.
Weber et al. (1995)
Weber, M E; Wiedicke, M; Riech, V & Erlenkeuser, H (1995): Carbonate preservation history in the Peru Basin: Paleoceanographic implications. Paleoceanography, 10(4): 775-800.
Van der Weijden et al. (1999)
Van der Weijden C H; Reichart, G J; Visser, H J (1999): Enhanced preservation of organic matter in sediments deposited within the oxygen minimum zone in the northeastern Arabian Sea. Deep-Sea Research I 46: 807-830.
Wenkam (1976)
Wenkam, C (1976): Late Quaternary changes in the oceanography of the eastern tropical Pacific, Corvallis, Oregon state University: 143.
Wessels (1989)
Wessels, M (1989): Sedimente vor dem Filchner-Schelfeis, Weddellmeer, Antarktis. Thesis, Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, unpublished thesis, unpublished: 34.
Westall (2000)
Westall, F (2000): 'Principal investigator' (0): unpublished data, Source: PANGAEA-

Network for Geological and Environmental Data, Alfred Wegener Institute for Polar and Marine Research (AWI), Center for Marine Environmental Sciences (MARUM).
Westerhausen et al. (1993)
Westerhausen, L; Poynter, J; Eglinton, H; Erlenkeuser, H; Sarnthein, M (1993): Marine and terrigenous origin of organic matter in modern sediments of the equatorial East Atlantik: the d13C and molecular record. <i>Deep Sea Res. I</i> 40(5): 1087-1121.
Winters et al. (1984)
Winters, G. V., D. E. Buckley, R. A. Cranston, R. A. Fitzgerald, M. Stoffyn, and P. Stoffyn-Egli (1984): Geological and geochemical data for sediment and pore water samples from the Sohm Abyssal Plain in the North Western Atlantic Ocean: Geological Survey of Canada, Open File Report 1082: 55.
Wollenburg and Mackensen (1998)
Wollenburg, J & Mackensen, A (1998): Living benthic foraminifers from the central Arctic Ocean: faunal composition, standing stock and diversity. <i>Marine Micropaleontology</i> , 34 (3-4): 153-185.
Yang et al. (1990)
Yang, Y-L; Elderfield, H; Ivanovich, M (1990): Glacial to holocene changes in carbonate and clay sedimentation in the Equatorial Pacific Ocean estimated from Thorium 230 profiles. <i>Paleocean.</i> 5(5): 789-809.
Yingst and Aller (1982)
Yingst, J Y and Aller, R C (1982): Biological activity and associated sedimentary structures in Hebble-Area deposits, Western North Atlantic. <i>Marine Geology</i> 48: