

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Ocean & Coastal Management

journal homepage: www.elsevier.com/locate/ocecoaman

Nonnative macrobenthos in the Wadden Sea ecosystem

Christian Buschbaum*, Dagmar Lackschewitz, Karsten Reise

Alfred Wegener Institute for Polar and Marine Research, Wadden Sea Station Sylt, Hafenstrasse 43, D-25992 List, Germany

ARTICLE INFO

Article history:
Available online 4 January 2012

ABSTRACT

An inventory in the Dutch-German-Danish Wadden Sea revealed a total of 66 nonnative (alien) taxa including 17 tentative cryptogenics in the brackish-marine macrobenthos until 2010, which is close to average compared with similar inventories from other coasts. Although the Wadden Sea is known for the largest sedimentary tidal flats in the world, most aliens encountered were fouling at harbor walls, pontoons in marinas, at hard structures for coastal defense but also in epibenthic mussel and oyster beds. Recent qualitative rapid assessments focusing on port localities have substantially improved knowledge on introduced species in the Wadden Sea. Nonnative species have pervaded algal and invertebrate communities, and the guild of suspension feeders became particularly strengthened by aliens which took advantage of a recent phase with relatively warm years. Most alien species were not directly introduced into the Wadden Sea but have spread secondarily from adjacent coasts with more active harbors or shellfish cultures. We suggest that aliens which have already established in the Wadden Sea ecosystem should be tolerated to avoid ongoing manipulations in eco-evolutionary developments in this protected nature area. Mitigation of the advancing tide of invasive aliens should focus on vector control (shipping and open aquacultures) as well as on early detection with attempts of eradication during the initial phase of invasions on a scale of the entire European Atlantic coast of which the Wadden Sea ecosystem is an integral part.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

In the Wadden Sea, species richness of macrobenthic organisms is comparatively low because this shallow coastal region of the North Sea is young with respect to geological history and has been above sea level during glacial periods. When 7,500 years ago sea level rise slowed down, barrier islands, extensive sediment flats, salt marshes and estuaries did evolve in roughly the present position (Reise et al., 2010). During the last thousand years, most salt marshes have gradually been converted into arable land, small estuaries into freshwater streams, large estuaries into shipping canals, and many soft shores into artificial hard shores (Lotze et al., 2005). Still, this coastal region is characterized by the largest coherent tidal flats on earth, attracting huge flocks of foraging birds and young fish from the North Sea. Therefore, the Wadden Sea has come under joint nature protection by The Netherlands, Germany and Denmark, and most of the area is listed as a World Heritage Site, based on a unique and outstanding geomorphology, ecological and biological processes and biodiversity (CWSS, 2008). Success in

environmental management includes a ban to the hunting of seals and birds, restrictions to the harvest of shellfish, restoration of salt marshes, and reductions in the discharge of nutrients and hazardous substances.

On the other hand, the introduction of alien species proceeds almost unchecked and undermines the conservation target of the trilateral Wadden Sea Plan (CWSS, 2010) to keep the ecosystem as natural as possible. Global trade by shipping is accelerating the rate of introduction of nonnative species in marine communities of European coasts (Leppäkoski et al., 2002; Steftaris et al., 2005), and is in particular transforming estuarine ecosystems (Campbell et al., 2007; Carlton, 1989; Ruiz et al., 1997; Williams and Grosholz, 2008; Wolff, 1999). It is widely assumed that invasions by alien species are a leading cause of recent species extinctions which is mainly concluded from studies on oceanic islands and freshwater lakes (Fritts and Rodda, 1998; Gurevitch and Padilla, 2004; Lockwood, 2004). However, there is no evidence for extinctions of native species in marine environments caused by invaders (Wolff, 2000a; Gurevitch and Padilla, 2004; Briggs, 2010; Craig, 2010). Nevertheless, it is feared that regionally the biota may lose their integrity by the chronic infiltration of alien species, resulting in a global biotic homogenization of ecosystems (Lövei, 1997; McKinney and Lockwood, 1999; Olden et al., 2004).

* Corresponding author. Tel.: +49 (0) 46519564228; fax: +49 (0) 4651956200.

E-mail addresses: Christian.Buschbaum@awi.de (C. Buschbaum), Dagmar.Lackschewitz@awi.de (D. Lackschewitz), Karsten.Reise@awi.de (K. Reise).

Furthermore, marine alien species are able to profoundly alter native communities, ecosystem functions and services (Ruiz et al., 1999; Occhipinti-Ambrogi and Galil, 2010; Ruesink et al., 2005), and this holds true for the European Wadden Sea as well (Reise et al., 2005; Nehring et al., 2009; Wolff et al., 2010). So far, no inventory of alien species has been accomplished for the trilateral Wadden Sea area, which is needed for joint management efforts. Up to now, only lists for national waters are available (Nehring and Leuchs, 1999; Jensen and Knudsen, 2005; Wolff, 2005; Gollasch and Nehring, 2006; Gittenberger et al., 2010) or summaries for the entire North Sea region (Reise et al., 1999; Gollasch et al., 2009). We here describe a rapid assessment at suspected hotspots of introductions in the German sector, combine this with results from a comparable survey in the Dutch sector (Gittenberger et al., 2010), as well as critically examined published records of alien species in order to compile a first inventory of nonnative macrobenthic species, comprising marine and brackish-water macroflora and macrofauna of the entire Wadden Sea area. Based on recent research, the integration of alien species in the ecosystem and options for their management in this protected nature area are discussed. We confine this study to macrobenthos because nonnative micro- and meiobenthos is not known for the Wadden Sea. For phyto- and zooplankton, coastal waters flowing through

the Wadden Sea are not a meaningful subregion and lists for the entire North Sea should be consulted (i.e., Reise et al., 1999; Gollasch et al., 2009), and permanent resident and seasonally visiting fish of nonnative origin have not yet been recorded in the waters of the Wadden Sea.

2. Methods

2.1. Rapid assessment

To investigate the occurrence and distribution of nonnative macrobenthos along the German Wadden Sea coast, we visited in 2009 eight locations between July and October once and repeated the investigation in 2010 (Fig. 1). At each location three different habitats were examined resulting in a total number of 24 sites (Table 1). Despite a variety of habitats, we particularly focused in our study on artificial structures in harbors and marinas since these are known to be gateways for alien newcomers with suitable habitats for establishment (Arenas et al., 2006; Ashton et al., 2006; Cohen et al., 2005; Minchin, 2007a; Gittenberger et al., 2010). Search and sampling by one to two experts lasted between 1 and 2 h at each site and 2–4 h per locality, depending on habitat size and heterogeneity. Most species were detected within the first




Fig. 1. The European Wadden Sea Area (surrounded by a red line) according to Marencic (2009) and investigated locations (see Table 1) of the rapid assessment of neobiota along the German Wadden Sea coast.

Table 1

Information on the locations and sites surveyed during the rapid assessment study at the German Wadden Sea coast in 2009 and 2010. The surface salinities represent the maximum values measured during investigations.

Locations (8)	Location coordinates		Salinity 2009/2010	Sites surveyed (24)
	Latitude	Longitude		
1. List	54°59.41'	8°23.05'	30/31 psu	Oyster farm and shallow sediments south of List Harbor pontoons for leisure boats Harbor mole and embankments
	55°00.59'	8°26.25'		
	55°02.00'	8°26.22'		
2. Hörnum	54°45.37'	8°17.45'	30/30 psu	Harbor pontoons for leisure boats Harbor mole and embankments Sandy beach sediment north of harbor
	54°45.38'	8°17.48'		
	54°45.57'	8°17.59'		
3. Büsum	54°07.24'	8°51.52'	25/24 psu	Harbor pontoons for leisure boats Harbor mole and embankments Mud and sand tidal area in northeastern harbor edge
	54°07.27'	8°52.02'		
	54°07.31'	8°52.09'		
4. Brunsbüttel	53°53.17'	9°07.17'	10/6 psu	Breakwater into the river Elbe Tidal mud flat along the river bank Embankment of the Kiel-Canal behind the sluice
	53°53.21'	9°07.20'		
	53°54.00'	9°09.12'		
5. Cuxhaven	53°53.35'	8°41.09'	20/18 psu	Breakwaters into the Elbe estuary at the 'Kugelbake' Sandy tidal flats at the northern tip ('Kugelbake') Harbor pontoons for leisure boats
	53°53.33'	8°41.05'		
	53°52.32'	8°42.16'		
6. Wilhelmshaven	53°30.35'	8°07.45'	30/30 psu	Breakwaters into the Jadebusen Sandy tidal areas south of the city Harbor pontoons for leisure boats (Nassauhafen)
	53°30.39'	8°07.54'		
	53°30.55'	8°08.59'		
7. Bensersiel	53°40.40'	7°34.16'	30/30 psu	Harbor pontoons for leisure boats Sandy tidal flats and beach Breakwater along the ferry canal
	53°40.47'	7°33.58'		
	53°40.56'	7°33.59'		
8. Emden	53°20.36'	7°11.23'	17/10 psu	Harbor pontoons for leisure boats Breakwaters and dike revetment at the river Ems Muddy and sandy areas along the northern river bank
	53°20.05'	7°11.12'		
	53°20.05'	7°11.15'		

15 min, and a survey was considered as complete when all sub-habitats had been visited and no more than one additional species were found for at least 30 min. Organisms attached to artificial structures such as floating pontoons (being independent of the tidal cycle) or harbor walls were sampled with scrapers. Sediments were dug up with a spade during low tide and sieved through a 1 mm mesh. In our experience, the focus on the intertidal as well as the shallow subtidal zone accessible to wading during low tide, together with the underside of floating objects, captures the spectrum of alien macrobenthic species in the Wadden Sea fairly well. Most organisms were identified in the field. Unknown species and minute organisms were preserved in 75% ethanol or transported alive to the laboratory and identified using optical appliance. Salinity of surface waters was measured once during sampling to get a rough estimate. Similar to other rapid assessments with a non-quantitative search-and-sample strategy (cited above), we aimed to detect as many alien species as possible, combining an efficient use of given resources in manpower and available time with the highest gain of information.

2.2. Compiling an overview of alien macrobenthos in the Wadden Sea

To achieve a valid and comprehensive list of nonnative and cryptogenic species (*sensu* Carlton, 1996) for the entire Wadden Sea, we performed an extensive literature study focusing on recent records to ensure the current distribution in the area. The distinction between nonnative and cryptogenic is often confounded by uncertainties. Except for the most recent records we followed Wolff (2005) in assigning taxa to these tentative categories. Only species assumed to originate outside the European Atlantic coasts are considered as introduced nonnative organisms. We included species whose status is still debated, but rejected conspicuous species with single records only, assuming these not (yet) to be established, i.e., the crabs *Limulus polyphemus* and *Callinectes sapidus* or the flatfish *Trinectes maculatus*. Likewise we did not list species above mean high-tide line, such as the saltmarsh spermatophyte *Cotula coronopifolia*.

The Wadden Sea can neither geographically nor ecologically strictly be separated from the North Sea and the incoming rivers.

We adopted the geographical definition of the Wadden Sea Area given by Marencic (2009) in Fig. 1. Boundaries in rivers are the transitions between freshwater and brackish water. Seaward the boundary is at three nautical miles offshore or at the outer limit of the conservation area in two cases. Although islands are part of the Wadden Sea Area, our assessment does not cover their freshwater and terrestrial habitats. Likewise, pelagic species are not included because these are better treated at the scale of the entire North Sea.

3. Results

3.1. Rapid assessment survey

The rapid assessment of alien species along the German Wadden Sea coast at eight locations comprising 24 sites yielded 25 and 30 species in 2009 and 2010, respectively. In total, we identified 34 nonnative or cryptogenic macrobenthic species covering 7 plants and 27 invertebrates (Table 2). Five species were encountered which have not been reported from the area so far. We found the cute tanaid crustacean *Sinelobus stanfordi* at the entrance of the Kiel-Canal in 2009 and 2010, and occurring also in the harbor of Emden in 2010. In 2009, we detected the bryozoan *Tricellaria inopinata*, new to the northern and southern German Wadden Sea, and in 2010 *Bugula neritina* in the harbor of Hörnum/Sylt. Two new alien red algae were recognized in 2010, namely *Ceramium cimbriicum* and *Antithamnionella ternifolia*. Only the latter was found in natural habitats.

The highest number of alien species (26) was found at two locations near the island of Sylt where salinity is high (Tables 1 and 2: location 1 + 2), whereas only few occurred at the entrance of the Kiel-Canal (Brunsbüttel) and in the Ems estuary where salinity is low (Tables 1 and 2: location 4 + 8). An oyster farm with annual imports of *Crassostrea gigas* and mussel cultures with imported *Mytilus edulis* from the British Isles may contribute to the amount of aliens near Sylt. The barnacle *Balanus improvisus* was observed at all locations and not more than 9 species were found at more than half of the locations. The majority of species (22) was found at 1–3 locations only, and ten were found either in 2009 or 2010. Thus,

Table 2

Alien and cryptogenic species recorded at 8 locations during the rapid assessment surveys on neobiota along the German Wadden Sea coast in 2009 and 2010. Location 1 List/Sylt, 2 Hörnum/Sylt, 3 Büsum, 4 Brunsbüttel, 5 Cuxhaven, 6 Wilhelmshaven, 7 Benseniel, 8 Emden (for further details see Fig. 1 and Table 1). Absent (–)/present (+) in 2009/2010, respectively. No entry: not found at locality.

Location	1	2	3	4	5	6	7	8
Phaeophyceae								
<i>Sargassum muticum</i>	+/+	+/+						
Chlorophyceae								
<i>Codium fragile tomentosoides</i>	+/-							
Rhodophyceae								
<i>Antithamionella ternifolia</i>	-/+							
<i>Ceramium cimbricum</i>		-/+						
<i>Gracilaria vermiculophylla</i>	+/+	+/+	+/+		-/+			+/+
<i>Polysiphonia harveyi</i>		-/+					-/+	
Tracheophyta								
<i>Spartina anglica</i>			+/+				+/+	
Cnidaria								
<i>Cordylophora caspia</i>				-/+				
Mollusca								
<i>Crepidula fornicata</i>	+/+	+/+				+/+	-/+	
<i>Crassostrea gigas</i>	+/+	+/+	+/+		+/+	+/+	+/+	+/+
<i>Ensis americanus</i>	-/+	+/+			-/+		+/+	
<i>Mya arenaria</i>	+/+	+/-	-/+		+/+	-/+	-/+	+/+
<i>Mytilopsis leucophaeta</i>				-/+				
<i>Petricola pholadiformis</i>							-/+	
<i>Teredo navalis</i>	+/+				+/+			
Polychaeta								
<i>Nereis virens</i>	+/-	+/-	+/+		+/-			
<i>Tharyx killariensis</i>	-/+							
Crustacea								
<i>Austrominius modestus</i>	+/+	+/+	+/+		+/+	+/+	+/+	
<i>Balanus improvisus</i>	-/+	-/+	+/+	+/+	+/+	-/+	-/+	+/+
<i>Sinelobus stanfordi</i>				+/+				-/+
<i>Caprella mutica</i>	+/+	+/+	-/+					
<i>Gammarus tigrinus</i>				+/+				
<i>Eriocheir sinensis</i>				+/+				
<i>Hemigrapsus takanoi</i>	+/+	-/+	-/+		+/+	+/+	+/+	
<i>Hemigrapsus sanguineus</i>	+/+	+/+	+/-		+/-	+/-	+/-	
<i>Palaemon macrodactylus</i>			+/-			-/+		-/+
Insecta								
<i>Telmatogeton japonicus</i>	-/+		+/+	+/+	+/+		+/+	-/+
Bryozoa								
<i>Bugula neritina</i>		-/+						
<i>Bugula stolonifera</i>		-/+					-/+	
<i>Tricellaria inopinata</i>	+/-	+/-					+/-	
Tunicata								
<i>Aplidium</i> c.f. <i>glabrum</i>	+/+	+/-					+/-	
<i>Botryllus schlosseri</i>	+/+	+/+					-/+	
<i>Molgula manhattensis</i>	+/+	+/+	+/+				-/+	+/+
<i>Styela clava</i>	+/+	+/+					+/+	
Species in total	22	21	13	7	11	16	12	5

alien species tend to show a rather uneven spatiotemporal distribution along the coast.

3.2. Alien species inventory

Up to 2010, a total of 66 nonnative (49) and cryptogenic (17) species of the macrobenthos have been recorded for the entire Wadden Sea (Table 3). The most diverse higher taxa are crustaceans (18 species), followed by macroalgae (14), polychaetes (8), mollusks (7), tunicates (6) and cnidarians (5). In the Dutch and German sectors 57 and 52 species respectively, have been recorded. In the Danish sector no specific assessment has been conducted and only 19 alien species are on the record there. The majority of the species presented in Table 3 can be regarded as established, however, for 8 of them the records are still very recent (2009 and/or 2010). Since the closure of the Zuiderzee, the nonnative snail *Corambe obscura*

has not been recorded anymore (Swennen and Dekker, 1995) and should be regarded as extinct for the Wadden Sea.

Most aliens in the Wadden Sea are marine species. Sixteen are either brackish water species or are regularly present under brackish conditions in addition to marine or freshwater. Two species cover the entire range from freshwater to marine (the eel parasite *Anguillicola crassus* and the crab *Eriocheir sinensis*). The most important donor regions are the western Pacific (28 species) followed by the western Atlantic (16 species). The origin of at least 25% is unknown or uncertain.

Most nonnatives found in the Wadden Sea have been encountered at other European coasts, mostly southwest of the Wadden Sea such as France (13 species), British Isles (16) and the Dutch Delta area (15) before arriving in the Wadden Sea. Only 8 are known to have entered directly this region and from there spread or were transported to adjacent coasts. Examples of the latter are the intentionally introduced cordgrass *Spartina anglica* and the Pacific oyster *C. gigas*, both imported from the British Isles but natural spread along coasts occurred in both species as well (Reise et al., 2005). The Chinese crab *E. sinensis* was first recorded in Europe in a tributary of the river Weser which debouches into the Wadden Sea. From there this crab has spread to other parts of Europe, although there is also genetic evidence for multiple introductions (Wang et al., 2009). The American razor clam *Ensis (directus) americanus* presumably arrived with ballast water released in the German Bight in front of the Wadden Sea or near the western Dutch Wadden Sea (Wolff, 2005). The mussel parasite *Mytilicola intestinalis* and the eel parasite *A. crassus* were first encountered near the Elbe estuary in the central Wadden Sea (Elsner et al., 2011; Koops and Hartmann, 1989). Three other species were first found in the former Zuiderzee which was an embayment of the Wadden Sea until its conversion into land and a freshwater reservoir since 1932 (Wolff, 2005). Such direct introductions account for only about 12% of the nonnative and cryptogenic species.

Only 9 of the macrobenthic aliens recorded in the Wadden Sea are dwelling in sediments. All others are fouling on artificial hard substrates of harbors and coastal defense, attach to biogenic hard structures such as mussel or oyster beds or are closely associated with these for hiding and feeding. Often first observations are from artificial structures and from there a spread to natural substrates has followed, i.e., *Caprella mutica*, *Styela clava*, *Aplidium* cf. *glabrum* and *Hemigrapsus* spp. (own observations).

4. Discussion

Rapid qualitative assessments of alien species with a focus on harbors, marinas and shellfish farming have substantially improved the knowledge on the occurrence and spatial distribution of nonnatives and cryptogenics in the Wadden Sea. In the discussion below we first compare the inventory to those of other coasts, consider management implications of secondary introductions and alien fouling organisms which prevail in the Wadden Sea, and suggest improvements for rapid assessments. Concerning nature management in the protected Wadden Sea we argue to tolerate the already established alien species, and to focus on vector control and eradications prior to a firm establishment of an alien species.

4.1. Inventory of alien species in the Wadden Sea

The inventory of nonnative and presumably nonnative (cryptogenic) macrobenthic organisms in the brackish and marine parts of the Wadden Sea comprised a total of 66 taxa until 2010.

As in other coastal studies, the macrobenthos is the group comprising most nonnative species. In addition, the North Sea coastal waters flowing through the Wadden Sea contain about 14

Table 3
 Alien and cryptogenic benthic macroalgae and -invertebrates and one vascular plant recorded until 2010 in the Dutch (NL), German (GE) and Danish (DK) sectors of the Wadden Sea. The status 'recent finding' refers to records from 2009 to 2010. Status or origin under discussion are denoted by '?'.

Species	Occurrence	Status	Salinity	Origin	Selected Ref.	Comments
Rhodophyta						
<i>Acrochaetium densum</i>	NL	Established	Marine	Pacific?	Wolff 2005	Recently found in the Wadden Sea (but established in the Delta region since longer)
<i>Anthamionella spirographidis</i>	NL	Established	Marine	N Pacific	Gittenberger et al., 2010	See Eno et al., 1997 for 'origin'
<i>Anthamionella ternifolia</i>	GE	Recent finding	Marine	S Pacific	This study	Acc. to Gittenberger et al., 2010
<i>Ceramium cimbriicum</i>	GE	Cryptogenic/established	Marine	Unknown	This study, Gittenberger et al., 2010	established in NL, only recently found in GE
<i>Ceramiales</i> sp.	NL	Cryptogenic/recent finding	Marine	Unknown	Gittenberger et al., 2010	Only recently found in the NL
<i>Gracilaria vermiculophylla</i>	NL	DK	Marine	W Pacific	Buschbaum et al., 2008 Gittenberger et al., 2010	
<i>Polysiphonia harveyi</i>	NL	GE	Marine	W Pacific	Thomsen et al., 2007a This study, Gittenberger et al., 2010	Syn. <i>Neosiphonia harveyi</i>
Chlorophyta						
<i>Codium fragile atlanticum</i>	NL	Recent finding	Marine	NW Pacific	Gittenberger et al., 2010	Recently found in the Wadden Sea area
<i>Codium fragile tomentosoides</i>	NL	DK	Marine	Pacific	Schories et al., 1997 Wolff 2005	Acc. to Nielsen 2005 the species is distributed at the Danish North Sea coast but where exactly remains unclear. Thomsen et al., 2007b list only drift specimen
<i>Ulva pertusa</i>	NL	Established	Marine	N Pacific	Gittenberger et al., 2010	
Phaeophyta						
<i>Boryella</i> sp.	NL	Established	Marine	Pacific?	Wolff 2005	A rare species but acc. to Wolff 2005 it seems to be established
<i>Colpomenia peregrina/C. sinuosa</i>	NL	Established	Marine	W Atlantic	Wolff 2005	In Wolff (2005) the two names were used synonymously while Gittenberger et al., 2010 consider them to be two valid species
<i>Sargassum muticum</i>	NL	GE	Marine	NW Pacific	Kremer et al., 1983 Schories and Albrecht 1995	Acc. to Nielsen 2005 only drifting along the Danish North Sea coast
<i>Undaria pinnatifida</i>	NL	Established	Marine	NW Pacific	Gittenberger et al., 2010	
Tracheophyta						
<i>Spartina anglica</i>	NL	DK	Marine	England	Loebj et al., 2006 Wolff 2005	Fertile hybrid of <i>S. alternifolia</i> and <i>S. maritima</i>
Porifera						
<i>Haliclona xena</i>	NL	Cryptogenic/established	Marine	Unknown	van Soest et al., 2007 Gittenberger et al., 2010	Syn. <i>Soestella xena</i>
Cnidaria						
<i>Bimeria francisiana</i>	GE	Uncertain	Marine	Unknown	Dittmer 1981	Syn. <i>Garveia francisiana</i> . <i>Perigoninus megas</i> ; a rare species in the Ems estuary (1960s), current status uncertain
<i>Cordylophora caspia</i>	NL	Established	Brackish	Ponto-Caspian	Dittmer 1981, this study Vervoort 1964	
<i>Nemopsis bachei</i>	NL	GE	Brackish to marine	W Atlantic	Frost et al., 2010 Wolff 2005	2011 also found in the harbors of Büsum and Benseniel by the authors (continued on next page)

Table 3 (continued)

Species	Occurrence	Status	Salinity	Origin	Selected Ref.	Comments
<i>Diadumene cincta</i>	NL	Established	Marine	Pacific?	Gittenberger et al., 2010	Presumed origin acc. to Nehring and Leuchs 1999
<i>Haliplanella lineata</i>	NL GE	Established	Marine	Pacific	Nehring and Leuchs 1999 Wolff 2005	Acc. to Nehring and Leuchs 1999 this species is probably extinct in the German Wadden Sea
Nematoda						
<i>Anguillicola crassus</i>	NL GE	Established	Freshwater to marine	W Pacific	Koops and Hartmann 1989 Wolff 2005	A parasitic nematod on fish (eel)
Mollusca						
<i>Corambe obscura</i>	NL	Extinct	Marine	W Atlantic	Wolff 2005	Extinct according to Wolff (2005)
<i>Crepidula fornicata</i>	NL GE	Established	Marine	NW Atlantic	Nehring and Leuchs 1999 Nehls et al., 2006	
<i>Crassostrea gigas</i>	NL GE	Established	Marine	W Pacific	Jensen and Knudsen 2005 Reise et al., 2005 Nehls and Büttger 2007 Gittenberger et al., 2010	
<i>Ensis americanus</i>	NL GE	Established	Marine	NW Atlantic	Wirange et al., 2010 Reise et al., 2005 Nehring and Leuchs 1999 Wolff 2005	Syn. <i>E. directus</i>
<i>Mya arenaria</i>	NL GE	Established	Marine	NW Atlantic	Jensen and Knudsen 2005 Strasser et al., 1999 Wolff 2005	Introduced before 1500
<i>Mytilopsis leucophaeta</i>	NL GE	Established	brackish	NW Atlantic	Smidt 1951 Jensen and Knudsen 2005 Post and Landmann 1994 This study	Syn. <i>Congeria leucophaeta</i>
<i>Petricola pholadiformis</i>	NL GE	Established	Marine	NW Atlantic	Wolff 2005 Nehring and Leuchs 1999 Wolff 2005	
<i>Teredo navalis</i>	NL GE	Cryptogenic/established	Marine	Unknown	Smidt 1951 Gollasch et al., 2009 Wolff 2005	
Polychaeta						
<i>Aphelochaeta marioni/Tharyx killariniensis</i>	NL GE	Cryptogenic/established	Marine	Unknown	Reise 1990 Wolff 2005 Jensen 1992	Obviously these two species have often been confused; not possible to evaluate the identification in retrospect
<i>Boccardia ligetica</i>	NL GE	Cryptogenic/established	Brackish	Unknown	Dittmer 1981 Osterkamp and Schirmer 2000 Wolff and Dankers 1981	Syn. <i>Polydora redeki</i> , <i>Polydora ligetica</i>
<i>Ficopomatus enigmaticus</i>	NL GE	Established	Brackish to marine	SW Pacific	Nehring and Leuchs 1999 Nehring and Leuchs 1999 Post and Landmann 1994	Syn. <i>Mercierella enigmatica</i> ; reported to be locally abundant in the harbor of Emden (GE)
<i>Marenzelleria neglecta</i>	NL GE	Established	Brackish	NW Atlantic	Gittenberger et al., 2010 Baštop and Blank 2006	
<i>Marenzelleria viridis</i>	NL GE	Established	Marine	NW Atlantic	Essink and Kleef 1993 Nehring et al., 2009 van der Graaf et al., 2009 Westheide 1966	Delefosse (in prep.)
<i>Microphthalamus similis</i>	GE	Cryptogenic/established	Marine	Unknown		
cf. <i>Neodexiospira brasiliensis</i>	NL	Recent finding	Marine	SW Atlantic?	Gittenberger et al., 2010	Syn. <i>Janua brasiliensis</i> ; recently found in the Dutch Wadden Sea but known already from elsewhere in the NL

<i>Nereis virens</i>	NL	GE	DK	Cryptogenic/ established	Marine	Unknown	Reise et al., 1994 Wolff 2005 Smidt 1951	Syn. <i>Alitta virens</i>
Crustacea								
<i>Mytilicola intestinalis</i>	NL	GE		Established	Brackish to marine	Mediterranean?	Elsner et al., 2011 Wolff 2005	A parasitic copepod in mussels; first record 1902 from the Adria
<i>Mytilicola orientalis</i>		GE		Recent finding	Marine	NW Pacific	Elsner et al., 2011	A parasitic copepod in oysters and mussels, recently found near Sylt
<i>Austrominius modestus</i>	NL	GE	DK	Established	Marine	S Pacific	Witte et al., 2010 Gittenberger et al., 2010 Jensen and Knudsen 2005	Syn. <i>Elminius modestus</i>
<i>Balanus improvisus</i>	NL	GE	DK	Cryptogenic/ established	Brackish to marine	Unknown	This study Gittenberger et al., 2010 Smidt 1951	
<i>Sinealobus stanfordi</i>	NL	GE		Recent finding	Brackish	Unknown	This study Gittenberger et al., 2010	Probably a native species
<i>Limnoria lignorum</i>	NL	GE		Cryptogenic/ established	Marine	Unknown	Wolff and Dankers 1981 Wolff 2005	
<i>Proasellus coxalis</i>	GE	GE		Established	Freshwater to brackish	Mediterranean	Post and Landmann 1994 Osterkamp and Schirmer 2000	An estuarine species, locally established (Ems and Weser estuary)
<i>Caprella mutica</i>	NL	GE	DK	Established	Marine	NW Pacific	Buschbaum and Gutow 2005 Gittenberger et al., 2010 Buschbaum pers. obs. 2010, harbour of Fanó (DK)	
<i>Chelicorophium curvispinum</i>		GE		Established	Freshwater to brackish	Ponto-Caspian	Nehring and Leuchs 1999	
(Mono-)Chorophium sextonae	NL	GE		Cryptogenic/ uncertain	Marine	Unknown	Nehring and Leuchs 1999 Wolff 2005	
<i>Gammarus tigrinus</i>	NL	GE		Established	Freshwater to brackish	NW Atlantic	This study, Klein 1969 Wolff 2005	Mainly a freshwater species, locally penetrating into estuaries (Elbe)
<i>Jassa marmorata</i>	NL	GE		Cryptogenic/ established	Marine	NW Atlantic	Gittenberger et al., 2010, own record	Only records from the Ems estuary and the harbour of List/Sylt but established in the North Sea
<i>Platorchestia platensis</i>	NL	GE		Established	Brackish	SW Atlantic	Witt and Krummwiede 2006 Rüppell 1967	Mainly a freshwater species penetrating into estuaries
<i>Eriocheir sinensis</i>	NL	GE	DK	Established	Freshwater to marine	NW Pacific	Den Hartog 1963 Jensen and Knudsen 2005 Gollasch 2006	While this amphipod is mostly regarded as native, Gittenberger et al., 2010 consider it to be introduced Semiterrestrial
<i>Hemigrapsus sanguineus</i>	NL	GE	DK	Established	Marine	NW Pacific	Gittenberger et al., 2010 Obert et al., 2007 This study Gittenberger et al., 2010	
<i>Hemigrapsus takanoi</i>	NL	GE		Established	Marine	NW Pacific	Obert et al., 2007 This study	For DK: pers. comm. (Reise); despite a recent appearance of <i>Hemigrapsus</i> sp. in the Wadden Sea high abundances and wide distribution refer to a successful establishment
<i>Palaemon macrodactylus</i>	NL	GE		Established	Marine	NW Pacific	Gittenberger et al., 2010 Gonzalez-Ortegon et al., 2007 Ruijter 2008 This study	Syn. <i>Hemigrapsus penicillatus</i> ; for 'status' see above
<i>Rhithropanopeus harrisi</i>	NL	GE		Established	Freshwater to brackish	W Atlantic	Nehring 2000 Wolff 2005	Acc. to Türkay (pers. comm. in Dürr 2010) stable populations exist in the Jadebusen (GE) (and North Sea)
Insecta								
<i>Telmatogiton japonicus</i>	GE	GE		established	marine	NW Pacific	This study	Mainly a freshwater to brackish species in canals and ditches

(continued on next page)

Table 3 (continued)

Species	Occurrence	Status	Salinity	Origin	Selected Ref.	Comments
Bryozoa						
<i>Bugula neritina</i>	GE	Recent finding	Marine	W Atlantic?	This study	2010 found in the harbor of Hörnum/Sylt (see Ryland et al., 2011 for 'origin') For discussion of status see Wolff 2005, Ryland et al., 2011 and Gittenberger et al., 2010
<i>Bugula stolonifera</i>	NL GE	Cryptogenic/ established	Marine	NW Atlantic	This study Gittenberger et al., 2010	
<i>Tricellaria inopinata</i>	GE	Recent finding	Marine	N Pacific	This study	found in 3 harbors in 2009
Tunicata						
<i>Aplidium</i> cf. <i>glabrum</i>	NL GE	Cryptogenic/ established	Marine	Unknown	Reise 1998 Gittenberger et al., 2010	Misidentified as <i>A. nordmanni</i> by Reise (1998); occurrence associated with Pacific oysters
<i>Botrylloides violaceus</i>	NL	Established	Marine	NW Pacific	Gittenberger et al., 2010	Probably a native species (Gittenberger et al., 2010), however other authors regard it as cryptogenic (Gollasch et al., 2009); Ben-Shlomo et al., 2006 suggest global distribution by ships
<i>Botryllus schlosseri</i>	NL NL	Cryptogenic/ established	Marine	Mediterranean?	This study Gittenberger et al., 2010 Smidt 1951	
<i>Didemnum vexillum</i>	NL	Established	Marine	NW Pacific	Gittenberger et al., 2010	
<i>Molgula manhattensis</i>	NL	Cryptogenic/ established	Marine	Unknown	This study Haydar et al., 2011	
<i>Styela clava</i>	NL GE	Established	Marine	NW Pacific	Reise 1998 Gittenberger et al., 2010 Lützen 1999	

phytoplankton and two zooplankton nonnative taxa (Gollasch et al., 2009; Reise et al., 1999) while permanent populations of alien fish are not yet known. The number of alien species in the Wadden Sea is close to average when compared to similar assessments in other coastal regions (Table 4). However, coastal regions with more active ports and/or shellfish farming (i.e., San Francisco Bay, Port Phillip Bay) or Mediterranean coasts adjacent to the Suez Canal (Galil, 2009) surpass the Wadden Sea in the number of nonnative taxa. In a comprehensive inventory of nonnative species in the coastal Netherlands, Wolff (2005) already pointed out that more aliens occurred in the small Oosterschelde estuary which is the centre of the Dutch shellfish culture than in the much larger Dutch Wadden Sea.

For managing the advancing flood of alien invasions it is an important finding that only a few nonnative species were directly introduced into the Wadden Sea, while most have arrived secondarily, either by natural dispersal from sites of primary introduction, by shellfish translocations, by regional shipping along the coasts of Western Europe or through inland canals and rivers. Wasson et al. (2001) observed a high number (56) of introduced invertebrate species in Elkhorn Slough, a small estuary lacking international shipping and located about 150 km down the shore from San Francisco Bay, which is the estuary with the highest load of alien invertebrate species, worldwide. Similar to the Wadden Sea, secondary immigrations have occurred from nearby hotspots of transoceanic shipping and shellfish farming, while direct introductions from overseas played a minor role in the invasion history of Elkhorn Slough. Consequently, effective prevention of further introductions into the Wadden Sea which is much larger than Elkhorn Slough but is also located downstream of coasts receiving more introductions, calls for concerted controls of overseas trans-ports on a larger scale. Thus, management of alien species requires a coordinated strategy for the entire European Atlantic coast.

Also important to the prevention of alien invasions is the observation that man-made hard structures constitute a major gateway and may provide stepping stones for further spread along the sedimentary coast of the Wadden Sea. The almost unlimited sprawl of artificial hard substrates (i.e., harbor walls, pilings, pontoons in marinas, buoys, breakwaters, dams, petrified shore-lines, garbage) offers plenty of habitats for fouling nonnative organisms introduced either on ship hulls or by transfers of cultured shellfish (Reise and Buschbaum, 2007). Prospects to restrict this highway for the spread of aliens are rather dim with the current proliferation of offshore wind parks in the North Sea. However, coastal defense in the Wadden Sea offers a large potential for substituting or covering hard structures by sand nourishments (Reise and Lackschewitz, 2003).

4.2. Rapid assessments focused on alien species

To effectively manage the issue of nonnative marine species, sound knowledge on the alien species already present is required (Campbell et al., 2007). Otherwise the success of precautionary measures to prevent further introductions cannot be measured. In the Wadden Sea, such surveys focused on harbors and marinas with their surroundings (Gittenberger et al., 2010; this study) which in the past have been neglected in benthic studies. This effort has complemented previous lists of alien species which were merely based on chance observations. Both rapid assessments of alien species (RAAs) performed at the German coast in 2009 and 2010 revealed a rather patchy occurrence of aliens among sampling sites and years. Thus, a necessary improvement would be to increase the number of sites to be investigated and to continue such surveys on an annual basis.

Table 4

Nonnative (including cryptogenics) macrobenthic invertebrate and algal species found in nine selected coastal regions. Species numbers were obtained from the lists of nonnatives provided in the literature.

Coastal region	Invertebrates	Algae	Total	Source
Wadden Sea (North Sea)	51	14	65	This study (without cordgrass <i>Spartina anglica</i>)
Dutch Delta area (North Sea)	55	20	75	Wolff (2005); Faasse and Lighthart (2009)
Baltic Sea	30	9	39	Paavola et al. (2005)
Ireland (NE-Atlantic)	28	15	43	Minchin (2007b)
Northern Adriatic (Mediterranean Sea)	37	12	49	Occhipinti-Ambrogi et al. (2011)
Chesapeake Bay (NW-Atlantic)	37	4	41	Ruiz et al. (2000)
Puget Sound and Straits (NE-Pacific)	65	3	68	Wonham and Carlton (2005)
San Francisco Bay (NE-Pacific)	105	5	110	Cohen and Carlton (1995); Wasson et al. (2001)
Port Phillip Bay (S-Australia)	90	67	157	Hewitt et al. (2004); Ruiz et al. (2000)

Relatively little time is required for field campaigns. However, identification of new alien species is often a time consuming endeavor and depends on taxonomical experience and the cooperation of specialists. A potential bottleneck for rapid assessments is to find scientists sufficiently trained in species identifications, additionally having knowledge on alien species spreading in other regions of Europe and thus are likely to appear sooner or later in the Wadden Sea as well. If RAAS are conducted to detect introductions of nonnatives at an early stage of invasion which would still allow effective eradication or control (see Simberloff, 2009), it would be necessary to perform such regular surveys in a well coordinated program along all European Atlantic coasts with a focus on ports and shellfish culture sites. As mentioned above, such a wide scale effort is essential because of the prevalence of secondary alien arrivals in the Wadden Sea.

4.3. How to manage alien species in the Wadden Sea?

Several species from native oysters to gray whales have been lost in the past from the Wadden Sea due to habitat loss, hunting and fishery (Reise, 1990; Wolff, 2000b), and there is consensus that environmental management should facilitate the return of lost species. Should management also try to get rid of the many species that have been intentionally or unintentionally introduced from overseas? Is this desirable and is it feasible? Since the issue of species introductions has advanced to an active discipline of research and management in the 1990s, the importance of regulating alien species invasions has been fiercely debated (for a recent example see Davis et al. (2011) with subsequent correspondence and letters). While one side points out that species of a nonnative origin should not be generally dismissed because not all constitute a threat and many may be beneficial as a resource or in the ecosystem, the other side insists on “aggressive intervention” against the “most severe and fastest growing threat to biological diversity” (Lambertini et al., 2011).

Although the political stage has been set by Article 8(h) of the Rio Convention on Biological Diversity adopted in 1992 (“... as far as possible and as appropriate, to prevent the introduction of, control and eradicate those alien species which threaten ecosystems, habitats or species.”) and further specified by *Guiding Principles on Invasive Alien Species* in 2002 (CBD, www.biodiv.org), regional management should be adjusted to the specific conditions of the Wadden Sea. In spite of the numerous alien species already accumulated in the Wadden Sea, we know of no native species which have gone extinct because of these invaders (Wolff, 2000a; own observations). The scenario of a few alien winners replacing many native losers (McKinney and Lockwood, 1999) does not fit. We suggest that extinctions by competitive exclusion are rather unlikely at a coast with high habitat heterogeneity and gradients in depth, salinity and substrates as well as with temporarily empty habitats due to frequent disturbances (Armonies and Reise, 2003). In addition,

a rich natural supply of nutrients and food from rivers and the North Sea, enhanced by anthropogenic eutrophication, may ease the accommodation of alien species. Particularly inviting are ‘alien’ substrates introduced for coastal defense, shipping or recreational purposes for introduced alien species (see chap. 4.1). Taken together, these natural characteristics and man-made changes may explain why the Wadden Sea ecosystem readily accommodated alien species without a concomitant loss of natives.

The course of previous alien species invasions into the Wadden Sea may not justify treating aliens as an urgent threat to native biodiversity. On the other hand, we observe an increasing number of species that also occur at other coastal regions. This advancing globalization entails a fading degree of biological uniqueness, and such a process is in conflict with the aim of nature conservation to sustain natural conditions. Management cannot reverse history but could implement measures which mitigate this development, i.e., by further reducing eutrophication, covering hard shores with sand and trying to minimize introducing alien substrates.

A problem for management arises from the limited predictability whether an alien will be harmless or turn into a threat before it has attained a wide spread and high abundance in a recipient region, and then it is – except for islands and lakes – not feasible anymore to reverse the invasion, particularly in the case of aquatic invertebrates and plants as in the Wadden Sea. Species differ in invasibility. However, a species-by-species approach cannot cope with cumulative effects and in protected nature areas as in the Wadden Sea (see the trilateral Wadden Sea Plan (CWSS, 2010)) alien introductions are generally not wanted. This justifies a generic attitude of preventing, eradicating or controlling nonnative species.

The Wadden Sea ecosystem is pervaded by wide-spread and abundant alien species such as the hybrid grass *S. anglica* forming a coherent belt at the outer edge of salt marshes, the Pacific oyster *C. gigas* replacing former mussel beds or the high biomass of the American razor clam *Ensis (directus) americanus* in the subtidal zone where native clams are scarce (Nehring et al., 2009). Many abundant aliens in the Wadden Sea belong to the functional group of suspension feeders, i.e., *C. gigas*, the American slipper limpet *Crepidula fornicata*, *E. americanus* (Reise and van Beusekom, 2008; Tulp et al., 2010). Presumably, the reinforced guild of benthic suspension feeders has increased its pressure on small pelagic organisms such as phytoplankton and larvae of benthic invertebrates. In addition, more fecal material and shells accumulate in the sediment. Together, it constitutes an ongoing change in the functioning of the Wadden Sea ecosystem, and this development is in conflict with the aim of protecting natural processes, and justifies management to stop the flood of new invaders.

The interactions of aliens with natives and among each other reveal an extraordinary complexity of competitive, predator-prey as well as parasite-host relationships with inhibiting and facilitating effects (Reise et al., 2006). For example, the slipper limpet *C. fornicata* affects mussels to which it is attached negatively but




Fig. 2. Prominent species occurring from the mid intertidal to the subtidal zone in the northern Wadden Sea. Intertidally, native species such as the lugworm *Arenicola marina* (1) and mussels *Mytilus edulis* (2) are abundant while the shallow subtidal zone is characterized by alien habitat engineers such as *Crassostrea gigas* (3), *Crepidula fornicata* (4), *Sargassum muticum* (5) which provide habitat for many mobile and fouling nonnatives including *Caprella mutica* (6), the tunicates *Styela clava* (7), *Molgula manhattensis* (8) and the barnacle *Austrominius modestus* (9). The American razor clam *Ensis americanus* (10) is the dominant bivalve in the subtidal zone. LWTL: Low Water Tide Line. Modified after Buschbaum and Reise, 2010.

also protects them against predators and parasite infections (Thieltges et al., 2006, 2009). Within oyster reefs, the native mussels are relegated to dwell in niches between the much larger oysters which results in decreased growth but increased survival due to refuges from crabs and other predators (Eschweiler and Christensen, 2011). The razor clam *Ensis americanus* has become one of the most abundant shellfish and also an important prey for fish and birds (Freundenthal et al., 2010; Tulp et al., 2010) but the effect on its planktonic food is still unknown and has not been studied yet. Alien plants and large benthic invertebrates function as ecosystem engineers, creating new habitat structures (Wallentinus and Nyberg, 2007; Bouma et al., 2009; Tang and Kristensen, 2010). This may facilitate native and alien species alike in a habitat otherwise poor in epibenthos (Fig. 2; Buschbaum et al., 2006; Polte and Buschbaum, 2008) but may also turn out to inhibit some species (Lang and Buschbaum, 2010).

Alien species in the Wadden Sea have not only increased regional species richness and ecological complexity but also evolutionary change is inevitably entailed in mixed native-nonnative species assemblages. Such eco-evolutionary dynamics between non natives and the recipient system have indelible results which cannot be simply reversed (Carroll, 2011). Invading populations will tend to differ more and more from their source populations and co-evolution will ultimately soften the distinction between nonnatives and natives in the invaded region. This is an argument to refrain from eradicating already established alien species.

There are several unanswered questions. Is the clam *Mya arenaria* still to be regarded as an alien although present in the Wadden Sea since several centuries? Does this old alien species deserve more rights to remain than the more recently introduced razor clam *E. americanus*? Both have attained an essential role in

the food web and it would be impossible to remove them without strong effects and presumable damage to the entire ecosystem. Clearly this is neither desirable nor feasible. When the native barnacle *Semibalanus balanoides* begins to decline in the wake of climatic warming, would it then be better to have an empty niche on intertidal hard substrates or to have this niche filled by the Australian barnacle *Austrominius modestus*, which has been introduced several decades ago and now benefits from milder winters and warmer summers (see Witte et al., 2010)?

We suggest that in the open coastal ecosystem of the protected Wadden Sea, the already established alien species deserve the same treatment as do the native species. Without such a decision, environmental management would embark on an endless chain of manipulating species compositions and species interactions, which would be in conflict with the aim to let natural processes proceed as far as possible in this protected nature area. On the other hand, mitigating the ongoing flood of alien invasions by controlling the vectors and attempting eradications before a firm establishment has occurred (Williams and Grosholz, 2008) should deserve priority in the nature management of the Wadden Sea and beyond.

5. Conclusions

- (1) With a total of 49 nonnative and 17 cryptogenic species of marine-to-brackish macrobenthic organisms established in the Wadden Sea, this shallow sedimentary coast has received an average amount of aliens compared to inventories of other coastal regions.
- (2) The prevalence of secondary introductions in the Wadden Sea necessitates that regular assessments of alien species with a focus on harbors, marinas and shellfish farms should be

performed in concert with similar surveys all along the European Atlantic coast.

- (3) Artificial hard structures play a key role in the establishment and spread of introduced alien species and should be minimized in the sedimentary Wadden Sea.
- (4) The Wadden Sea ecosystem is heavily pervaded by nonnative species. However, the resulting eco-evolutionary development is irreversible and the already established alien species should be tolerated and not treated differently from native species.
- (5) For mitigating the ongoing flood of alien invasions, concerted management over the entire Atlantic coast of Europe should focus on vector control and eradication prior to firm establishment at gateway sites of alien introductions rather than on the scale of the Wadden Sea alone.

Acknowledgments

We thank Rolf Karez and the State Agency for Agriculture, Environment and Rural Areas, Flintbek (LLUR-SH) for supervising and funding our projects on alien species in German coastal waters, and also Jan Witt (Lower Saxony Water Management, Coastal Defence and Nature Conservation Agency, NLWKN) for project support. Advice from Wolfgang Groepler and Ralph Kühlenkamp on taxonomic questions has been extremely helpful. We thank three anonymous reviewers for identifying omissions, for constructive criticism and their encouragement.

References

- Arenas, F., Bishop, J.D.D., Carlton, J.T., Dyrnynda, P.J., Farnham, W.F., Gonzalez, D.J., Jacobs, M.W., Lambert, C., Lambert, G., Nielsen, S.E., Pederson, J.A., Porter, J.S., Ward, S., Wood, C.A., 2006. Alien species and other notable records from a rapid assessment survey of marinas on the south coast of England. *J. Mar. Biol. Assoc. U.K.* 86, 1329–1337.
- Armonies, W., Reise, K., 2003. Empty habitat in coastal sediments for populations of macrozoobenthos. *Helgol. Mar. Res.* 56, 279–287.
- Ashton, G., Boos, K., Shucksmith, R., Cook, E., 2006. Rapid assessment of the distribution of marine non-native species in marinas in Scotland. *Aquat. Invasions* 1 (4), 209–213.
- Bastrop, R., Blank, M., 2006. Multiple invasions – a polychaete genus enters the Baltic Sea. *Biol. Invasions* 8, 1195–1200.
- Ben-Shlomo, R., Paz, G., Rinkevich, B., 2006. Postglacial-period and recent invasions shape the population genetics of botryllid ascidians along European Atlantic coasts. *Ecosystems* 9, 1118–1127.
- Bouma, T.J., Olenin, S., Reise, K., Ysebaert, T., 2009. Ecosystem engineering and biodiversity in coastal sediments: posing hypotheses. *Helgol. Mar. Res.* 63, 95–106.
- Briggs, J.C., 2010. Marine biology: the role of accommodation in shaping marine biodiversity. *Mar. Biol.* 157, 2117–2126.
- Buschbaum, C., Gutow, L., 2005. Mass occurrence of an introduced crustacean (*Caprella cf. mutica*) in the south-eastern North Sea. *Helgol. Mar. Res.* 59, 252–253.
- Buschbaum, C., Reise, K., 2010. Neues Leben im Weltnaturerbe Wattenmeer. Globalisierung unter Wasser. *Biol. Unserer Zeit* 40 (3), 202–210.
- Buschbaum, C., Chapman, A.S., Saier, B., 2006. How an introduced seaweed can affect epibiota diversity in different coastal systems. *Mar. Biol.* 148, 743–754.
- Buschbaum, C., Mayr, T., Scheuer, K., 2008. Vorkommen der invasiven Rotalge *Gracilaria vermiculophylla* und ihre ökologischen Effekte im Schleswig-Holsteinischen Wattenmeer. *Landesamt für Natur und Umwelt des Landes Schleswig-Holstein*, Flintbek, pp. 1–50.
- Campbell, M.L., Gould, B., Hewitt, C.L., 2007. Survey evaluations to assess marine bioinvasions. *Mar. Poll. Bull.* 55, 360–378.
- Carlton, J.T., 1989. Man's role in changing the face of the ocean: biological invasions and implications for conservation of near-shore environments. *Conserv. Biol.* 3, 265–273.
- Carlton, J.T., 1996. Biological invasions and cryptogenic species. *Ecology* 77, 1653–1655.
- Carroll, S.P., 2011. Conciliation biology: the eco-evolutionary management of permanently invaded biotic systems. *Evol. Appl.* 4, 184–199.
- Cohen, A.N., Carlton, J.T., 1995. Nonindigenous Aquatic Species in a United States Estuary: A Case Study of the Biological Invasions of the San Francisco Bay and Delta. U.S. Fish and Wildlife Service, Arlington, Virginia, USA.
- Cohen, A.N., Harris, L.H., Bingham, B.L., Carlton, J.T., Chapman, J.W., Lambert, C.C., Lambert, G., Ljubenkov, J.C., Murray, S.N., Rao, L.C., Reardon, K., Schwindt, E., 2005. Rapid assessment survey for exotic organisms in southern California bays and harbors, and abundance in port and non-port areas. *Biol. Invasions* 7, 995–1002.
- Craig, M.T., 2010. Pattern versus process: broadening the view of marine invasive species. *Mar. Biol.* 157, 2127–2128.
- CWSS, 2008. Nomination of the Dutch–German Wadden Sea as World Heritage Site. WaddenSea Ecosystem No. 24. Common Wadden Sea Secretariat, Wilhelmshaven, Germany, pp. 1–200.
- CWSS, 2010. Wadden Sea Plan 2010. Common Wadden Sea Secretariat, Wilhelmshaven, Germany. www.waddensea-secretariat.org, pp. 1–88.
- Davis, M., et al., 2011. Don't judge species on their origins. *Nature* 474, 153–154 (see also correspondence in *Nature* 475, 36–37 and *Science* 333, 404–405).
- Den Hartog, C., 1963. The amphipods of the Deltaic region of the rivers Rhine, Meuse and Scheldt in relation to the hydrography of the area. *Neth. J. Sea Res.* 2, 40–67.
- Dittmer, J.D., 1981. The distribution of subtidal macrobenthos in the estuaries of the rivers Ems and Weser. In: Dankers, N., Köhl, H., Wolff, W. (Eds.), *Invertebrates of the Wadden Sea*, Report 4. Balkema, Rotterdam, pp. 188–206.
- Dürr, A., 2010. Untersuchungen zur Populations- und Fortpflanzungsbiologie von Garnelen in der westlichen Ostsee. Diplom-thesis. Univ. Kiel, p. 160.
- Elsner, N.O., Jacobsen, S., Thielges, D.W., Reise, K., 2011. Alien parasitic copepods in mussels and oysters of the Wadden Sea. *Helgol. Mar. Res.* 65, 299–307.
- Eno, N.C., Clark, R.A., Sanderson, W.G., 1997. Non-native Marine Species in British Waters: A Review and Directory. Joint Nature and Conservation Committee, Peterborough, p. 152.
- Eschweiler, N., Christensen, H.T., 2011. Trade-off between increased survival and reduced growth for blue mussels living on Pacific oyster reefs. *J. Exp. Mar. Biol. Ecol.* 403, 90–95.
- Essink, K., Kleef, H.L., 1993. Distribution and life cycle of the North American spionid *Marenzelleria viridis* (Verrill, 1873) in the Ems estuary. *Neth. J. Aquatic Ecol.* 27, 237–246.
- Faasse, M., Ligthart, M., 2009. American (*Urosalpinx cinerea*) and Japanese oyster drill (*Ocenebrellus inornatus*) (Gastropoda: Muricidae) flourish near shellfish culture plots in The Netherlands. *Aqua. Invasions* 4, 321–326.
- Freudenthal, A.S.L., Nielsen, M.M., Jensen, T., Jensen, K.T., 2010. The introduced clam *Ensis americanus* in the Wadden Sea: field experiment on impact of bird predation and tidal level on survival and growth. *Helgol. Mar. Res.* 64, 93–100.
- Fritts, T.H., Rodda, G.H., 1998. The role of introduced species in the degradation of island ecosystems: a case history of Guam. *Annu. Rev. Ecol. Syst.* 29, 113–140.
- Frost, J.R., Jacoby, C.A., Youngbluth, M.J., 2010. Behavior of *Nemopsis bachei* L. Agassiz, 1849 medusae in the presence of physical gradients and biological thin layers. *Hydrobiologia* 645, 97–111.
- Galil, B.B., 2009. Taking stock: inventory of alien species in the Mediterranean Sea. *Biol. Invasions* 11, 359–372.
- Gittenberger, A., Rensing, M., Stegenga, H., Hoeksema, B., 2010. Native and non-native species of hard substrata in the Dutch Wadden Sea. *Nederlandse Faunistische Medelingen* 33, 21–75.
- Gollasch, S., Nehring, S., 2006. National checklist for aquatic alien species in Germany. *Aqua. Invasions* 1 (4), 245–269.
- Gollasch, S., Haydar, D., Minchin, D., Wolff, W.J., Reise, K., 2009. Introduced aquatic species of the North Sea coasts and adjacent brackish waters. In: Rilov, G., Crooks, J.A. (Eds.), *Biological Invasions in Marine Ecosystems*. *Ecol. Stud.*, vol. 204, pp. 507–528.
- Gollasch, S., 2006. NOBANIS – Invasive Alien Species Fact Sheet – *Eriocheir sinensis*. From: Online Database of the North European and Baltic Network on Invasive Alien Species – NOBANIS. www.nobanis.org 2011.
- González-Ortegón, E., Cuesta, J.A., Schubart, C.D., 2007. First report of the oriental shrimp *Palaeomon macrrodactylus* Rathbun, 1902 (Decapoda, Caridea, Palaemonidae) from German waters. *Helgol. Mar. Res.* 61, 67–69.
- Gurevitch, J., Padilla, D.K., 2004. Are invasive species a major cause of extinctions? *Trends Ecol. Evol.* 19 (9), 470–474.
- Haydar, D., Hoarau, G., Olsen, J.L., Stam, W.T., Wolff, W., 2011. Introduced or glacial relict: phylogeography of the cryptogenic tunicate *Molgula manhattensis* (Ascidacea, Pleurogona). *Divers. Distrib.* 17, 68–80.
- Hewitt, C.L., Campbell, M.L., Thresher, R.E., et al., 2004. Introduced and cryptogenic species in Port Phillip Bay, Victoria, Australia. *Mar. Biol.* 144, 183–202.
- Jensen, K.R., Knudsen, J., 2005. A summary of alien marine benthic invertebrates in Danish waters. *Oceanol. Hydrobiol. Stud.* 24 (1), 137–162.
- Jensen, K.T., 1992. Macrozoobenthos on an intertidal mudflat in the Danish Wadden Sea: comparisons of surveys made in the 1930s, 1940s and the 1980s. *Helgoländer Meeresunters* 46, 363–376.
- Klein, G., 1969. Amphipoden aus der Wesermündung und der Helgoländer Bucht, mit Beschreibung von *Talorchestia frisiae* n. sp. *Veröff. Inst. Meeresforsch. Bremerhaven* 11, 173–194.
- Koops, H., Hartmann, F., 1989. *Anguillicola*-infestations in Germany and in German eel imports. *J. Appl. Ichthyol.* 1, 41–45.
- Kremer, B.P., Kubbier, H., Michaelis, H., 1983. Die Ausbreitung des Brauntanges *Sargassum muticum* in der Nordsee. *Natur und Museum* 113, 125–130.
- Lambertini, M., et al., 2011. Invasives: a major conservation threat. *Science* 333, 404–405.
- Lang, A.C., Buschbaum, C., 2010. Facilitative effects of introduced Pacific oysters on native macroalgae are limited by a secondary invader, the seaweed *Sargassum muticum*. *J. Sea Res.* 63, 119–128.
- Leppäkoski, E., Gollasch, S., Olenin, S. (Eds.), 2002. *Invasive Aquatic Species of Europe. Distribution, Impacts and Management*. Kluwer Academic Publishers, Dordrecht, p. 583.

- Lockwood, J.L., 2004. How do biological invasions alter diversity patterns? A biogeographic perspective. In: Lomolino, M.V., Heaney, L.R. (Eds.), *Frontiers of Biogeography*. Sinauer Associates, Sunderland, MA, pp. 271–310.
- Loebel, M., Van Beusekom, J.E.E., Reise, K., 2006. Is spread of the neophyte *Spartina anglica* recently enhanced by increasing temperatures? *Aqua. Ecol.* 40, 315–324.
- Lotze, H.K., et al., 2005. Human transformations of the Wadden Sea ecosystem through time: a synthesis. *Helgol. Mar. Res.* 59, 84–95.
- Lövei, G.L., 1997. Global change through invasion. *Nature* 388, 627.
- Lützen, J., 1999. *Styela clava* (Urochordata, Ascidiacea), a successful immigrant to North West Europe: ecology, propagation and chronology of spread. *Helgoländer Meeresunters* 52, 383–391.
- Marencic, H. (Ed.), 2009. The Wadden Sea – Introduction. Thematic Report No. 1. In: Marencic, H. & 1de Vlas, J. (Eds.), 2009. Quality Status Report 2009. Wadden Sea Ecosystem No. 25. Common Wadden Sea Secretariat, Trilateral Monitoring and Assessment Group, Wilhelmshaven, Germany, pp. 3–24.
- McKinney, M.L., Lockwood, J.L., 1999. Biotic homogenization: a few winners replacing many losers in the next mass extinction. *Trends Ecol. Evol.* 14 (11), 450–452.
- Minchin, D., 2007a. Rapid coastal survey for targeted alien species associated with floating pontoons in Ireland. *Aqua. Invasions* 2 (1), 63–70.
- Minchin, D., 2007b. A checklist of alien and cryptogenic aquatic species in Ireland. *Aqua. Invasions* 2, 341–366.
- Nehls, G., Büttger, H., 2007. Spread of the Pacific Oyster *Crassostrea gigas* in the Wadden Sea – Causes and Consequences of a successful invasion. Trilateral Workshop on Pacific Oyster Invasion in the Wadden Sea, vol. 22. Common Wadden Sea Secretariat, Wilhelmshaven, Germany, p. 54.
- Nehls, G., Diederich, S., Thielges, D.W., Strasser, M., 2006. Wadden Sea mussel beds invaded by oysters and slipper limpets: competition or climate control? *Helgol. Mar. Res.* 60, 135–143.
- Nehring, S., Hesse, K.-J., 2008. Invasive alien plants in marine protected areas: the *Spartina anglica* affair in the European Wadden Sea. *Biol. Invasions* 10, 937–950.
- Nehring, S., Leuchs, H., 1999. Neozoa (Makrobenthos) an der deutschen Nordseeküste. Eine Übersicht. Bundesanstalt für Gewässerkunde, Koblenz, p. 131.
- Nehring, S., Reise, K., Dankers, N., Kristensen, P.S., 2009. Alien species. Thematic report No. 7. In: Marencic, H., de Vlas, J. (Eds.), Quality Status Report 2009. Wadden Sea Ecosystem No. 25. Common Wadden Sea Secretariat, Trilateral Monitoring and Assessment Group, Wilhelmshaven, Germany.
- Nehring, S., 2000. Zur Bestandssituation von *Rhithropanopeus harrisi* (GOULD, 1841) in deutschen Gewässern: Die sukzessive Ausbreitung eines amerikanischen Neozoons. *Senckenbergiana maritima* 30, 115–122.
- Nielsen, R., 2005. Danish Seaweeds. Botanical Museum Copenhagen, Denmark.
- Obert, B., Herlyn, M., Grotjahn, M., 2007. First records of two crabs from the North West Pacific *Hemigrapsus sanguineus* and *H. takanoi* at the coast of Lower Saxony. *Germany Wadden Sea Newsl.*, 21–22. 2007-1.
- Occhipinti-Ambrogi, A., Galil, B., 2010. Marine alien species as an aspect of global change. *Adv. Oceanogr. Limnol.* 1 (1), 143–156.
- Occhipinti-Ambrogi, A., et al., 2011. Alien species along the Italian coasts: an overview. *Biol. Invasions* 13, 215–237.
- Olden, J.D., LeRoy Poff, N., Douglas, M.R., Douglas, M.E., Fausch, K.D., 2004. Ecological and evolutionary consequences of biotic homogenization. *Trends Ecol. Evol.* 19 (1), 18–24.
- Osterkamp, S., Schirmer, M., 2000. Projekt 'Klimaänderung und Unterweserregion' (Fallstudie Weserästuar). Bund-Länder-Programm 'Klimaänderung und Küste'. Universität Bremen, p. 151.
- Paavola, M., Olenin, S., Leppäkoski, E., 2005. Are invasive species most successful in habitats of low native species richness across European brackish water seas? *Est. Coast. Shelf Sci.* 64, 738–750.
- Polte, P., Buschbaum, C., 2008. Native pipefish *Entelurus aequoreus* are promoted by the introduced seaweed *Sargassum muticum* in the northern Wadden Sea. *Aquat. Biol.* 3, 11–18.
- Post, D., Landmann, M., 1994. Verbreitungsatlas der Fließgewässerfauna in Ostfriesland. Staatliches Amt für Wasser und Abfall, Aurich, p. 141.
- Reise, K., Buschbaum, C., 2007. Mehr Sand statt Stein für die Ufer der Nordseeküste. *Rostocker Meeresbiologische Beiträge* 17, 77–86.
- Reise, K., Lackschewitz, D., 2003. Combating habitat loss at eroding Wadden Sea shores by sand replenishments. In: Wolff, W.J., Essink, K., Kellermann, A., van Leeuwe, M.A. (Eds.), *Challenges to the Wadden Sea*. Proceedings of the 10th Int. Sci. Wadden Sea Symp., Groningen 2000. Ministry of Agriculture, Nature Management and Fisheries, Dept. Mar. Biol. Univ. Groningen, pp. 197–206.
- Reise, K., van Beusekom, J.E.E., 2008. Interactive effects of global and regional change on a coastal ecosystem. *Helgol. Mar. Res.* 62, 85–91.
- Reise, K., Herre, E., Sturm, M., 1994. Biomass and abundance of macrofauna in intertidal sediments of Königshafen in the northern Wadden Sea. *Helgoländer Meeresunters* 48, 201–215.
- Reise, K., Gollasch, S., Wolff, W.J., 1999. Introduced marine species of the North Sea coasts. *Helgoländer Meeresunters* 52, 219–234.
- Reise, K., Dankers, N., Essink, K., 2005. Introduced species. In: Essink, K., Dettmann, C., Farke, H., Laursen, K., Lüerßen, G., Marencic, H., Wiersinga, W. (Eds.), *Wadden Sea Quality Status Report 2004*. Wadden Sea Ecosystem No. 19. Common Wadden Sea Secretariat, Wilhelmshaven, Germany, pp. 155–161.
- Reise, K., Olenin, S., Thielges, D.W., 2006. Are aliens threatening European aquatic coastal ecosystems? *Helgol. Mar. Res.* 60, 77–83.
- Reise, K., Baptist, M., Burbridge, P., Dankers, N., Fischer, L., Flemming, B., Oost, A.P., Smit, C., 2010. The Wadden Sea – a universally outstanding tidal wetland. *Wadden Sea Ecosys.* 29, 9–23. Common Wadden Sea Secretariat, Wilhelmshaven, Germany.
- Reise, K., 1990. Historische Veränderungen in der Ökologie des Wattenmeeres. *Rhein-Westfäl. Akad. Wiss., Vorträge N 382*, 35–50.
- Reise, K., 1998. Pacific oysters invade mussel beds in the European Wadden Sea. *Senckenbergiana marit.* 28 (4/6), 167–175.
- Ruesink, J.L., Lenihan, H.S., Trimble, A.C., Heiman, K.W., Micheli, F., Byers, J.E., Kay, M.C., 2005. Introduction of non-native oysters: ecosystem effects and restoration implications. *Annu. Rev. Ecol. Evol. Syst.* 36, 643–689.
- Ruijter, R. de, 2008. Cs-verslag. *Het Zeepaard* 68 (5), 2–7.
- Ruiz, G.M., Carlton, J.T., Grosholz, E.D., Hines, A.H., 1997. Global invasions of marine and estuarine habitats by non-indigenous species: mechanisms, extent and consequences. *Am. Zool.* 37, 621–632.
- Ruiz, G.M., Fofonoff, P., Hines, A.H., Grosholz, E.D., 1999. Non-indigenous species as stressors in estuarine and marine communities: assessing invasion impacts and interactions. *Limnol. Oceanogr.* 44 (3), 950–972.
- Ruiz, G.M., Fofonoff, P., Carlton, J.T., Wonham, M.J., Hines, A.H., 2000. Invasion of coastal marine communities in North America: apparent patterns, processes and biases. *Annu. Rev. Ecol. Syst.* 31, 481–531.
- Rüppell, G., 1967. Zur Lokomotionsaktivität des Amphipoden *Orchestia platensis* im Freiland und im Laboratorium. *Helgoländer Meeresunters* 15, 172–180.
- Ryland, J.S., Bishop, J.D.D., Blauwe, H. De, Nagar, A. El, Minchin, D., Wood, C.A., Yunnice, A.L.E., 2011. Alien species of *Bugula* (Bryozoa) along the Atlantic coasts of Europe. *Aqua. Invasions* 6, 17–31.
- Schories, D., Albrecht, A., 1995. *Sargassum muticum* – Der japanische Beerentang im deutschen Wattenmeer. *Natur und Museum* 125, 92–98.
- Schories, D., Albrecht, A., Lotze, H., 1997. Historical changes and inventory of macroalgae from Königshafen Bay in the Northern Wadden Sea. *Helgoländer Meeresunters* 51, 321–341.
- Simberloff, D., 2009. We can eliminate invasions or live with them. *Successful management projects*. *Biol. Invasions* 11, 149–157.
- Smidt, E.L.B., 1951. Animal production in the Danish Waddensea. *Meddelelser fra Kommissionen for Danmarks Fiskeri- og Havundersøgelser Serie: Fiskeri XI* (6).
- Soest, R.W.M. van, Kluijver, M.J. de, Bragt, P.H. van, Faasse, M., Nijland, R., Beglinger, E., Weerdt, W.H. de, Voogd, N.J. de, 2007. Sponge invaders in Dutch coastal waters. *J. Mar. Biol. Ass. U.K.* 87, 1733–1748.
- Strasser, M., Walensky, M., Reise, K., 1999. Juvenile-adult distribution of the bivalve *Mya arenaria* on intertidal flats in the Wadden Sea: why are there so few year classes? *Helgol. Mar. Res.* 53, 45–55.
- Streftaris, N., Zenetos, A., Papanassasiou, E., 2005. Globalisation in marine ecosystems: the story of non-indigenous marine species across European seas. *Oceanogr. Mar. Biol. Ann. Rev.* 43, 419–454.
- Swennen, C., Dekker, R., 1995. *Corambe batava* Kerbert, 1886 (Gastropoda, Opisthobranchia), an immigrant in The Netherlands, with a revision of the family Corambidae. *J. Molluscan Stud.* 61, 97–107.
- Tang, M., Kristensen, E., 2010. Associations between macrobenthos and invasive cordgrass, *Spartina anglica*, in the Danish Wadden Sea. *Helgol. Mar. Res.* 64, 321–329.
- Thielges, D.W., Strasser, M., Reise, K., 2006. How bad are invaders in coastal waters? The case of the American slipper limpet *Crepidula fornicata* in western Europe. *Biol. Invasions* 8, 1673–1680.
- Thielges, D.W., Reise, K., Prinz, K., Jensen, K.T., 2009. Invaders interfere with native parasite-host interactions. *Biol. Invasions* 11, 1421–1429.
- Thomsen, M.S., Staehr, P.A., Nyberg, C., Schwaerter, S., Krause-Jensen, D., Silliman, B.R., 2007a. *Gracilaria vermiculophylla* (Ohmi) Papenfuss, 1967 (Rhodophyta, Gracilariaceae) in northern Europe, with emphasis on Danish conditions, and what to expect in the future. *Aqua. Invasions* 2, 83–94.
- Thomsen, M.S., Wernberg, T., Staehr, P.A., Krause-Jensen, D., Risgaard-Petersen, N., Silliman, B.R., 2007b. Alien macroalgae in Denmark – a broad-scale national perspective. *Marine Biol. Res.* 3, 61–72.
- Tulp, I., Craeymeersch, J., Leopold, M., van Damme, C., Fey, F., Verdaat, H., 2010. The role of the invasive bivalve *Ensis directus* as food source for fish and birds in the Dutch coastal zone. *Est. Coast. Shelf Sci.* 90, 116–128.
- Vervoort, W., 1964. Note on the distribution of *Garveia franciscana* (Torrey, 1902) and *Cordylophora caspia* (Pallas, 1771) in the Netherlands. *Zool. Meded* 39, 125–146.
- Graaf, S., van der Vlas, J., de Herlyn, M., Voss, J., Heyer, K., Drent, J., 2009. Macrozoobenthos. Thematic report no. 10. In: Marencic, H., Vlas, J. de (Eds.), Quality Status Report 2009. Wadden Sea Ecosystem No. 25. Common Wadden Sea Secretariat, Trilateral Monitoring and Assessment Group Wilhelmshaven, Germany.
- Wallentinus, I., Nyberg, C.D., 2007. Introduced marine organisms as habitat modifiers. *Mar. Pollut. Bull.* 55, 323–332.
- Wang, C., Li, S., Fu, C., Gong, X., Huang, L., Song, X., Zhao, Y., 2009. Molecular genetic structure and evolution in native and colonized populations of the Chinese mitten crab, *Eriocheir sinensis*. *Biol. Invasions* 11, 389–399.
- Wasson, K., Zabin, C.J., Bedinger, L., Diaz, M.C., Pearce, J.S., 2001. Biological invasions of estuaries without international shipping: the importance of intraregional transport. *Biol. Conservation* 102, 143–153.
- Westheide, W., 1966. Zur Polychaetenfauna des Eulitorals der Nordseeinsel Sylt. *Helgoländer wiss. Meeresunters* 13, 2003–2009.
- Williams, S.L., Grosholz, E.D., 2008. The invasive species challenge in estuarine and coastal environments: marrying management and science. *Estuaries and Coasts* 31, 3–20.

- Witt, J., Krumwiede, A., 2006. Fahrrinnenanpassung der Unterweser an die Entwicklungen im Schiffsverkehr. Entwicklung eines Bewertungskonzeptes und Bewertung des Makrobenthos der Unterweser. Im Auftrag des Wasser- und Schiffsamtes Bremerhaven (WSA).
- Witte, S., Buschbaum, C., Beusekom, J.E.E. van, Reise, K., 2010. Does climatic warming explain why an introduced barnacle finally takes over after a lag of more than 50 years? *Biol. Invasions* 12, 3579–3589.
- Wolff, W.J., Dankers, N., 1981. Preliminary checklist of the zoobenthos and nekton species of the Wadden Sea. In: Dankers, N., Kühl, H., Wolff, W. (Eds.), *Invertebrates of the Wadden Sea, Report 4*. Balkema, Rotterdam, pp. 24–60.
- Wolff, W.J., Bakker, J.P., Laursen, K., Reise, K., 2010. The Wadden Sea Quality Status Report – Synthesis Report 2010. Wadden Sea Ecosystem No. 29. Common Wadden Sea Secretariat, Wilhelmshaven, Germany, pp 27–71.
- Wolff, W.J., 1999. Exotic invaders of the meso-oligohaline zone of estuaries in The Netherlands: why are there so many? *Helgol. Meeresunters* 52, 393–400.
- Wolff, W.J., 2000a. Causes of extirpations in the Wadden Sea, an estuarine area in The Netherlands. *Conserv. Biol.* 14, 876–885.
- Wolff, W.J., 2000b. The south-east North Sea: losses of vertebrate fauna during the past 2000 years. *Biol. Cons.* 95, 209–217.
- Wolff, W.J., 2005. Non-indigenous marine and estuarine species in the Netherlands. *Zoologische Mededelingen* 79 (1), 1–116.
- Wonham, M.J., Carlton, J.T., 2005. Trends in marine biological invasions at local and regional scales: the Northeast Pacific Ocean as a model system. *Biol. Invasions* 7, 369–392.
- Wrangle, A.-L., Valero, J., Harketstad, L.S., Strand, Ø., Lindegarth, S., Christensen, H.T., Dolmer, P., Kristensen, P.S., Mortensen, S., 2010. Massive settlements of Pacific oyster, *Crassostrea gigas*, in Scandinavia. *Biol. Invasions* 12, 1453–1458.