



The Occurrence of Phycotoxins in Coastal Areas of South America

Bernd Krock¹, Urban Tillmann¹, Nicole Trefault², Gustavo A. Lovrich³, C. Marcela Borel⁴

¹ Alfred Wegener Institute for Polar and Marine Research,
Am Handelshafen 12, D-27570 Bremerhaven, Germany.
E-Mail: bernd.krock@awi.de


² Universidad Mayor, Centro de Genómica y Bioinformática, Instituto de Biotecnología,
Camino a la Pirámide 5750, Huechuraba, Santiago de Chile

³ Centro Austral de Investigaciones Científicas (CADIC-CONICET),
Houssay 200, Ushuaia - Tierra del Fuego, Argentina

⁴ Instituto Geológico del Sur (CONICET-UNS),
San Juan 670, Bahía Blanca, Argentina



1. Study areas



March 2009
Hespérides



Nov/Dec 2010
Abate Molina

March/April 2012
Puerto Deseado



2. The Hespérides Expedition Mar 2009

Water pumping, 25 – 100 µm size fraction, only lipophilic toxins and domoic acid





Trefault N., Krock B., Delherbe N., Cembella A.D., Vásquez M., 2011. Toxicon 58, 389-397.



2. The Hespérides Expedition Mar 2009

Álvarez G., Uribe E., Ávalos P., Mariño C., Blanco J., 2010. First identification of azaspiracid and spirolides in *Mesodesma donacium* and *Mulinia edulis* from Northern Chile. *Toxicon* 55, 638-641.



López-Rivera A., O'Callaghan K., Moriarty M., O'Driscoll D., Hamilton B., Lehane M., James K.J., Furey A., 2010. First evidence of azaspiracids (AZAs): A family of lipophilic polyether marine toxins in scallops (*Argopecten purpuratus*) and mussels (*Mytilus chilensis*) collected in two regions of Chile. *Toxicon* 55, 692-701.





2. The Hespérides Expedition Mar 2009

Water pumping, 25 – 100 µm size fraction, only lipophilic toxins and domoic acid





Trefault N., Krock B., Delherbe N., Cembella A.D., Vásquez M., 2011. Toxicon 58, 389-397.



2. The Hespérides Expedition Mar 2009


Different *Dinophysis* species in Chilean coastal waters



Trefault N., Krock B., Delherbe N., Cembella A.D., Vásquez M., 2011. *Toxicon* 58, 389-397.





3. The Abate Molina Expedition Nov/Dec 2010







3. The Abate Molina Expedition 2010

DTX-1 found in Nov/Dec 2009








4. The Puerto Deseado Expedition Mar/Apr 2012



San Jorge Gulf



4. The Puerto Deseado Expedition Mar/Apr 2012





The overlay of nutrient-rich, cold Beagle-Magellan water with warm, nutrient-depleted shelf water results in a high primary production



4. The Puerto Deseado Expedition Mar/Apr 2012





Enormous dinoflagellate bloom dominated by non-toxic *Ceratium* spp.







4. The Puerto Deseado Expedition Mar/Apr 2012






4. The Puerto Deseado Expedition Mar/Apr 2012



PSP toxin profiles of two *Alexandrium tamarensis* isolates from Station C43



4. The Puerto Deseado Expedition Mar/Apr 2012





5. Conclusions

AZAs are present in Chilean coastal waters, but have not been detected in Argentinean Shelf waters

PTXs occur throughout South American waters with a relatively regular distribution; PTX-11 could only be detected in Pacific but not in Atlantic waters

In the Chilean southern Patagonian Fjords DTX-1 and DTX-2 were detected, but no OA; in the years 2009/2010 DTX-1 and -2 were separated geographically and temporally

The San Jorge Gulf is characterized by cold, nutrient rich Beagle-Magellan water overlaid by warmer nutrient depleted shelf water; this situation causes high density dinoflagellate blooms including toxic species.

C1/C2 and GTX1/4 are converted to GTX2/3 abiotically within 3 months under elevated ambient temperatures




Nicole Trefault



Gustavo Lovrich



Urban Tillmann



Marcela Borel






Funding:

This work was partly financed by the HGF Programme in Earth and Environment under the research initiative PACES of the Alfred-Wegener Institute (Germany)

2009: Fondef MR07I-1005, Fondecyt 1080075 and LinkGlobal (PUC-CSIC) (Chile)

2010: Proyecto CONA C16F 10-13 (Chile)

2012: MINCyT-BMBF (AL/11/03-ARG 11/021)
(Argentina & Germany)



Thank you for your attention

