

This discussion paper is/has been under review for the journal The Cryosphere (TC).  
Please refer to the corresponding final paper in TC if available.

# SMOS derived sea ice thickness: algorithm baseline, product specifications and initial verification

X. Tian-Kunze<sup>1</sup>, L. Kaleschke<sup>1</sup>, N. Maaß<sup>1</sup>, M. Mäkynen<sup>2</sup>, N. Serra<sup>1</sup>, M. Drusch<sup>3</sup>, and T. Krumpen<sup>4</sup>

<sup>1</sup>Institute of Oceanography, University of Hamburg, Bundesstraße 53, 20146 Hamburg, Germany

<sup>2</sup>Finnish Meteorological Institute, P.O. Box 503, 00101 Helsinki, Finland

<sup>3</sup>European Space Agency, ESA-ESTEC, 2200 AG Noordwijk, the Netherlands

<sup>4</sup>Alfred Wegener Institute for Polar and Marine Research, Bussestrasse 24, 27570 Bremerhaven, Germany

Received: 21 November 2013 – Accepted: 26 November 2013 – Published: 6 December 2013

Correspondence to: X. Tian-Kunze (xiangshan.tian-kunze@zmaw.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Following the launch of ESA's Soil Moisture and Ocean salinity (SMOS) mission it has been shown that brightness temperatures at a low microwave frequency of 1.4 GHz (L-band) are sensitive to sea ice properties. In a first demonstration study, sea ice thickness has been derived using a semi-empirical algorithm with constant tie-points. Here we introduce a novel iterative retrieval algorithm that is based on a sea ice thermodynamic model and a three-layer radiative transfer model, which explicitly takes variations of ice temperature and ice salinity into account. In addition, ice thickness variations within a SMOS footprint are considered through a statistical thickness distribution function derived from high-resolution ice thickness measurements from NASA's Operation IceBridge campaign. This new algorithm has been used for the continuous operational production of a SMOS based sea ice thickness data set from 2010 on. This data set is compared and validated with estimates from assimilation systems, remote sensing data, and airborne electromagnetic sounding data. The comparisons show that the new retrieval algorithm has a considerably better agreement with the validation data and delivers a more realistic Arctic-wide ice thickness distribution than the algorithm used in the previous study.

## 1 Introduction

Satellite-based observation of ice thickness is still very challenging. The first satellite borne observations of ice thickness were conducted with satellite radar altimeters carried on European Remote Sensing satellites (ERS-1 and ERS-2) (Laxon et al., 2003) and thermal imagery from Advanced Very High Resolution Radiometer (AVHRR) (Yu and Rothrock, 1996; Drucker et al., 2003). The altimeter observations were followed by the ICESat laser altimeter from 2003 to 2009 (Kwok and Cunningham, 2008) and since 2011 by the CryoSat-2 radar altimeter (Laxon et al., 2013). The radar and laser altimeters have large uncertainties for ice thickness less than 1 m

TCD

7, 5735–5792, 2013

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Laxon et al., 2003; Kwok and Cunningham, 2008). Therefore, they are more suitable for the detection of thick ice. The altimeter ice thickness charts typically have a one month temporal resolution and a 100 km spatial resolution.

Thin ice thickness up to around 0.5 m with 1 km spatial resolution can be estimated with thermal imagery based ice surface temperature ( $T_s$ ) together with atmospheric forcing data through ice surface heat balance equation (Yu and Rothrock, 1996; Maeky-nen et al., 2013). The major drawback with the  $T_s$  based thickness retrieval is the requirement for cloud-free conditions, and thus, there may be long temporal gaps in the thickness chart coverage over a region of interest. In addition, discriminating clear-sky from clouds is difficult in winter night-time conditions (Frey et al., 2008).

Passive microwave radiometer data from Special Sensor Microwave Imager (SSM/I) (37 and 85.5 GHz channels) and Advanced Microwave Scanning Radiometer-Earth Observing System (AMSR-E) (36.5 and 89 GHz channels) sensors have been used to estimate thickness of thin ice up to 10–20 cm (Martin et al., 2005; Tamura et al., 2007; Nihashi et al., 2009; Tamura and Ohshima, 2011; Singh et al., 2011). The spatial resolution of the radiometer thin ice thickness charts (6.25 to 25 km) is much coarser than that from thermal imagery, but daily Arctic and Antarctic coverage is possible. The thin ice thickness retrieval algorithms are linear or exponential regression equations between polarization ratios (PR) or V- to H-polarization ratio (R) and AVHRR or Moderate Resolution Imaging Spectroradiometer (MODIS) thickness charts. Naoki et al. (2008) suggested that the observed decrease of near ice surface salinity as a function of ice thickness, which results in modification of the ice dielectric properties and further ice emission, is the main reason for the observed brightness temperature vs. ice thickness relationship. In addition, the brightness temperature vs. ice thickness relationship is more pronounced for H-polarization and for a lower frequency (e.g. 10.7 GHz). Nihashi et al. (2009) found out that PR at 37 GHz cannot detect thin ice when it is covered with snow. An analysis of ship-borne radiometer data at 19, 37 and 85 GHz over various thin ice types indicated that a limitation in the thin ice thickness estimation can be attributed

**SMOS derived sea ice thickness**

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the presence of snow or dense frost flower coverage (> 60 %) on the ice surface (Hwang et al., 2007).

The Soil Moisture and Ocean Salinity (SMOS) mission of the European Space Agency (ESA), which was launched in November 2009, measures for the first time globally Earth's radiation at a frequency of 1.4 GHz in the L-band (Mecklenburg et al., 2012). The footprint varies from about 35 km to more than 50 km. Besides soil moisture and ocean salinity information, for which SMOS was originally designed, L-band radiometry on SMOS can also be used to obtain the sea ice thickness, which is due to the large penetration depth in sea ice (Kaleschke et al., 2010, 2012). In contrast to IceSat and CryoSat-2 measurements, SMOS-derived ice thickness has less uncertainty in the thin ice range, but an exponentially increasing uncertainty for ice thickness thicker than 0.5 m. In our study we consider ice thickness less than 50 cm as thin ice. SMOS-derived ice thickness can thus complement the measurements from CryoSat-2 to achieve Arctic-wide sea ice thickness estimations (Kaleschke et al., 2010, 2012).

The semi-empirical SMOS ice thickness retrieval algorithm applied previously in Kaleschke et al. (2012) (hereinafter Algorithm I) is

$$TB(d_{ice}) = T_1 - (T_1 - T_0)e^{-\gamma d_{ice}}, \quad (1)$$

where  $d_{ice}$  is the ice thickness,  $T_1$  and  $T_0$  are two constant tie points which were estimated from the observed SMOS brightness temperatures over open water and thick first year ice during the freezing period of 2010 in the Arctic, and  $\gamma$  is a constant attenuation factor which was derived from a sea ice radiation model (Menashi et al., 1993) for a representative bulk ice temperature and bulk ice salinity in the Arctic.

The advantage of Algorithm I is the retrieval of ice thickness from the brightness temperature (TB) without any auxiliary data set. However, the TB measured by a L-band radiometer over sea ice depends on the dielectric properties of sea ice which are functions of ice temperature and ice salinity (Kaleschke et al., 2010). Although the change of TB caused by the sea ice thickness variation is much larger than that caused by the variation of ice temperature and ice salinity, the typical variability of these two

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



parameters in the Arctic can induce up to 30 K difference in TB (Kaleschke et al., 2012). This means the assumption of constant retrieval parameters could cause considerable errors in the regions where these parameters strongly differ from the assumed constant values.

5 Ice temperature and ice salinity measurements are rare and they are not continuously available on a daily basis in the Arctic. An alternative solution is therefore to derive these two parameters from auxiliary data during the sea ice thickness retrieval. Under the assumption of thermal equilibrium, the ice surface temperature can be estimated from the surface air temperature. Therefore, we use a heat flux balance equation and use the surface air temperature from atmospheric reanalysis data as a boundary condition. Ice salinity can be estimated from the underlying sea surface salinity (SSS) with an empirical function (Ryvlin, 1974). With these two parameters we can calculate brightness temperature with the sea ice radiation model (Menashi et al., 1993). However, both ice temperature and ice salinity are in turn functions of ice thickness. Thus, we need to apply a linear approximation method to simultaneously retrieve ice thickness and estimate suitable ice temperature and salinity values. This algorithm is called Algorithm II hereinafter.

10 In the radiation model of Menashi et al. (1993) a plane ice layer is assumed. However, due to sea ice deformation in natural sea ice, a broad scale of ice thicknesses occurs within one footprint of SMOS. The brightness temperature measured by SMOS at each footprint is a mixture of brightness temperatures from different ice thicknesses, and possibly open water. As SMOS brightness temperature is more sensitive to ice thicknesses less than 0.5 m (Kaleschke et al., 2012), SMOS-derived ice thickness under the assumption of a plane ice layer tends to represent the lower end of the ice thickness distribution within the footprint. A possible solution for the corresponding underestimation of ice thickness is to correct the retrieved ice thickness using an ice thickness distribution function. The correction of ice thickness retrieved from Algorithm II using this function is called Algorithm II\* in this study.

---

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**SMOS derived sea ice thickness**

X. Tian-Kunze et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Here we compare the three different SMOS ice thickness retrieval algorithms in the Arctic. The plane layer ice thicknesses retrieved from Algorithm I and II are compared with independent data to examine if the method that considers variable ice temperature and ice salinity improves the accuracy of the ice thickness retrieval. Thereafter, sea ice thickness uncertainty is estimated with the better algorithm on a daily basis. The growth of the sea ice cover as seen by SMOS during a freezing period in the Arctic is also discussed.

The paper is structured as follows. In Sect. 2 we describe the SMOS brightness temperature and the auxiliary data sets. The baseline of Algorithm II is described in Sect. 3. In Sect. 4 we discuss the uncertainties and bias of the retrieved ice thickness. After that we present in Sect. 5 our method to correct the retrieved ice thickness, which is based on the assumption of a plane ice layer, with an empirically determined ice thickness distribution function. The comparison of ice thicknesses retrieved from different algorithms is discussed in Sect. 6. Ice thickness growth and distribution as seen by SMOS during the freeze-up period in the Arctic are shown in Sect. 7. A further comparison of SMOS-derived ice thickness with that derived from MODIS in the Kara Sea is presented in Sect. 8. Finally, summaries and discussions are given in Sect. 9.

## 2 Data

Three different data sets are used for the retrieval of sea ice thickness in Algorithm II. The basis of the retrieval is the brightness temperature measured by the SMOS L-band radiometer. This data set is described in Sect. 2.1. For the estimation of bulk ice temperature ( $T_{ice}$ ) we use surface air temperature ( $T_a$ ) from Japanese 25 yr Reanalysis (JRA-25) data which are described in Sect. 2.2. The SSS climatology, which is used for the calculation of bulk ice salinity ( $S_{ice}$ ), is presented in Sect. 2.3. Finally, MODIS ice thickness charts over the Kara Sea for the verification of the SMOS ice thickness are presented in Sect. 2.4.

## 2.1 SMOS brightness temperature data

### 2.1.1 L1C data

The SMOS payload Microwave Imaging Radiometer using Aperture Synthesis (MI-RAS) measures in L-band the brightness temperatures in full polarization with incidence angles ranging from  $0^\circ$  to  $65^\circ$ . All four Stokes parameters are obtained. It has a global coverage every three days (Kerr et al., 2001), whereas daily coverage up to  $85^\circ$  can be expected in the polar regions. Brightness temperature is taken every 1.2 s by hexagon-like, two-dimensional snapshots which have a spatial dimension of about 1200 km across (Kerr et al., 2001). The geometric distribution of incidence angles and radiometric accuracy within the alias-free areas of a snapshot is shown in Fig. 1. The footprint varies from about 35 km at nadir view to more than 50 km at incidence angles higher than  $60^\circ$ . Each snapshot measures one or two of the Stokes components in the antenna reference frame. Horizontally and vertically polarized brightness temperatures are measured by separate snapshots.

The SMOS L1C data are given on the Discrete Global Grid (DGG) system. The DGGs are fixed Earth grid coordinates of the ISEA 4–9 hexagonal grid centers which have a spatial distance of 15 km (Indra, 2010). Most of the pixels in the Arctic are covered by several overflights during one day. Therefore, for our daily product we collect at each DGG grid point all brightness temperatures measured during one day, together with other information like the incidence angles.

### 2.1.2 Radio frequency interference

SMOS measurements are partly influenced by Radio Frequency Interference (RFI) which comes from radars, TV and radio transmission (Mecklenburg et al., 2012). The detection of the RFI sources and the mitigation of RFI influence are critical steps for the further retrieval of geophysical parameters. The RFI influence depends on the incidence angle, polarization, and ascending and descending modes of the satellite

TCD

7, 5735–5792, 2013

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Camps et al., 2010). A closer look into RFI contaminated snapshots shows that RFI can either completely or partly destroy a snapshot (Camps et al., 2010). For simplification we apply a threshold value for both horizontally and vertically polarized brightness temperatures. If either of them exceeds 300 K within one snapshot, this snapshot is considered as RFI contaminated. Brightness temperatures higher than 300 K can not be expected in the Arctic and Antarctic.

According to this RFI filter, strongly RFI affected regions are the region northeast of Greenland and parts of the Canadian Arctic Archipelago. Figure 2 shows the RFI induced data loss based on our RFI filter. The data loss in the figure is defined as the ratio between the number of RFI contaminated measurements and the number of total measurements. As can be seen from Fig. 2, the RFI situation in the Arctic region has improved much since 2010.

### 2.1.3 Brightness temperature intensity

Over sea ice the first Stokes parameter (intensity) is almost independent of incidence angle in the incidence angle range of 0–40° (Fig. 3). The intensity is the average of the horizontally and the vertically polarized brightness temperatures, i.e. it is equal to  $0.5(TB_h + TB_v)$ . The intensity is independent of both geometric and Faraday rotations and robust to instrumental and geophysical errors (Camps et al., 2005). We can avoid additional uncertainties caused by the transformation from the antenna reference frame to the Earth reference frame by using the intensity. Since each snapshot measures either horizontally or vertically polarized brightness temperature, we use consecutive snapshots with an acquisition time difference of less than 2.5 s to calculate the intensity. The advantage of using near nadir view measurements is the smaller footprint associated with low incidence angles. Furthermore, by using the whole incidence angle range of 0–40° we get more than 100 brightness temperature measurements per day for most of the pixels in the Arctic, and by averaging over a lot of measurements we can significantly reduce the uncertainty of the retrieval. The daily averaged brightness temperature intensities in the Arctic and in the Antarctic are interpolated with nearest

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



neighbor algorithm and gridded into the National Snow and Ice Data Center (NSIDC) polar stereographic projection with a grid resolution of 12.5 km. We use this grid resolution because other products that we use as auxiliary data in the retrieval are also given in this resolution. We call this product L3B brightness temperature. In the following we use TB to indicate the daily averaged brightness temperature intensity. The data have been processed with about 24 h latency for both hemispheres and available since January 2010. The L3B TBs are the basis of our sea ice thickness retrieval with Algorithm I and II and can be obtained from [icdc.zmaw.de](http://icdc.zmaw.de).

## 2.2 JRA-25 reanalysis data

For estimating the ice surface temperature, we extract the 2 m surface air temperature and the 10 m wind velocity data from JRA-25 atmospheric reanalysis data and interpolate them into the polar stereographic projection with 12.5 km grid resolution. JRA-25 reanalysis data provide various physical variables in 1.125° resolution every six hours. The data have been produced by the Japanese Meteorological Agency using the latest numerical analysis and prediction system. The data are available from 1979 to the present (Onogi et al., 2007). Various studies have been carried out to compare the JRA-25, ERA40 and NCEP data sets. Good agreements were found between JRA-25 and ERA40 (Onogi et al., 2007).

## 2.3 Sea surface salinity climatology

The SSS information is needed to estimate the bulk ice salinity which is an input parameter of the radiation model of sea ice. There are global ocean salinity products derived from SMOS brightness temperatures. Ocean salinity is one of the two applications SMOS was originally designed for. However, SMOS-derived ocean salinity is not available for the ice covered regions in the Arctic. Thus, we use a SSS climatology based on the outputs of an ocean-sea ice model. SSS weekly climatology is computed based on the daily output from a high-resolution numerical simulation of the Atlantic and Arctic

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oceans' circulations using the MIT general circulation ocean-sea ice coupled model (Marshall et al., 1997). The model is configured for the Atlantic Ocean north of 33° S, including all Atlantic marginal seas and the Arctic Ocean up to the Bering Strait, and is integrated at the eddy-resolving resolution of approximately 4 km. The vertical resolution of the model varies from 5 m in the upper ocean to 275 m in the deep (100 vertical levels are used). Bottom topography is interpolated from the ETOPO2 database and initial temperature and salinity conditions from a 8 km resolution integration of the same model (to achieve a good degree of spin-up), which in turn starts from the WOA09 climatology. The model is forced at the surface by fluxes of momentum, heat and freshwater computed internally in the model with the help of the 6 hourly atmospheric state from the ECMWF/ERA-Interim Reanalysis (Dee et al., 2011) and bulk formula. At the open northern (Bering Strait) and southern (33° S) boundaries, the model is forced by a 1° resolution global solution. The K-Profile Parameterisation (KPP) formulation is used for the parameterization of vertical mixing, with a background vertical viscosity coefficient of  $1 \times 10^{-4} \text{ m}^2$ . The vertical diffusion employed amounts to  $1 \times 10^{-5} \text{ m}^2 \text{ s}^{-1}$ . Unresolved horizontal mixing uses a bi-harmonic diffusion/viscosity of  $3 \times 10^9 \text{ m}^4 \text{ s}^{-1}$ . The overall good performance of the above model configuration (integrated at 8 km resolution), assessed through comparisons with in-situ measurements, can be found in Serra et al. (2010); Brath et al. (2010); Dmitrenko et al. (2012). We choose to use a model climatology and not Polar science center Hydrographic climatology (PHC) to benefit from the dynamical oceanographic structures realistically resolved in the model, which leads to spatial and seasonal variability of SSS.

Figure 4 shows the mean and standard deviation of weekly SSS from October to April, based on 8 yr of daily model output. SSS in the Laptev Sea, parts of the Kara Sea, and the Baltic Sea is much lower than in the central Arctic due to the influence of river run-offs. On the contrary, in the Baffin Bay, the Greenland Sea, and the Barents Sea the SSS is higher than in the central Arctic. The mean weekly SSS in the Baltic sea varies in the range of  $4\text{--}10 \text{ g kg}^{-1}$  which agrees well with the observed climatology given in Janssen et al. (1999). As the ice salinity is correlated with SSS, the variability of

SSS should be considered in the retrieval when we calculate Arctic-wide ice thickness distributions.

## 2.4 MODIS ice thickness charts

MODIS ice thickness charts covering an area of 1500 km × 1350 km over the Kara Sea and the eastern part of the Barents Sea have been calculated. The derivation of the charts and their uncertainty estimation are described in detail in Maekynen et al. (2013). The total number of the charts is 120 and they cover two winters (November to April) in 2009–2011. The spatial resolution of the charts is 1 km and they show ice thickness from 0 to 99 cm. The external forcing data for solving the ice thickness from the surface heat balance equation come from a numerical weather prediction (NWP) model HIRLAM (High-Resolution Limited Area Model) (Kaellen, 1996; Uden, 2002). Only night-time MODIS data are employed. Thus, the uncertainties related to the effects of solar shortwave radiation and surface albedo are excluded. For the cloud masking of the MODIS data, in addition to the different cloud tests (Frey et al., 2008), also manual methods are used in order to improve the detection of thin clouds and ice fog. The cloud masking is conducted with 10 km × 10 km blocks to identify larger cloud-free areas and to reduce errors due to the MODIS sensor striping effect. In the ice thickness chart calculation an average snow thickness ( $h_s$ ) vs. ice thickness ( $h_i$ ) relationship is used. The thickness of the snow layer is assumed to be:

$$\begin{aligned} h_s &= 0 \text{ m} && \text{for } d_{\text{ice}} < 0.05 \text{ m} \\ h_s &= 0.05 \times d_{\text{ice}} && \text{for } 0.05 \text{ m} \leq d_{\text{ice}} < 0.2 \text{ m} \\ h_s &= 0.09 \times d_{\text{ice}} && \text{for } d_{\text{ice}} \geq 0.2 \text{ m} \end{aligned}$$

This relationship is based on Doronin (1971) and the Soviet Union's Sever expeditions data (NSIDC, 2004). The typical maximum reliable ice thickness (max 50% uncertainty) is estimated to be 35–50 cm under typical weather conditions (air temperature  $T_a < -20^\circ\text{C}$ , wind speed  $V_a < 5 \text{ ms}^{-1}$ ) for the MODIS data. The accuracy is best

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for the 15–30 cm thickness range, around 38%. These figures are based on Monte Carlo method using estimated or guessed standard deviations and covariances of the input variables to the thickness retrieval. No in-situ data are available for the thickness accuracy estimation.

### 3 Sea ice thickness retrieval Algorithm II

Algorithm I is described in detail in Kaleschke et al. (2012), we will here introduce the retrieval Algorithm II. As in Algorithm I we use the daily mean brightness temperature intensity TB averaged over 0–40° incidence angle range.

#### 3.1 The sea ice radiation model

The basis of the SMOS ice thickness retrievals Algorithm I and II is the sea ice radiation model adapted from Menashi et al. (1993). While for Algorithm I the radiation model is used to calculate the constant attenuation factor  $\gamma$  for a representative  $T_{ice}$  and  $S_{ice}$  in the Arctic, in Algorithm II the model is used to calculate TB at variable  $T_{ice}$  and  $S_{ice}$ .

The sea ice radiation model consists of a plane ice layer bordered by the underlying sea water and air on the top. The model does not include a snow layer. The TB over sea ice depends on the dielectric properties of the ice layer which are a function of brine volume (Vant et al., 1978). The brine volume is a function of  $S_{ice}$  and  $T_{ice}$  (Cox and Weeks, 1983).

For a thin ice layer, the ice temperature gradient within the ice can be assumed to be linear (Maaß, 2013a). Assuming that the water under sea ice is at the freezing point, we can calculate  $T_{ice}$  with  $0.5(T_{si} + T_w)$ , where  $T_{si}$  is the snow-ice interface temperature and  $T_w$  is the freezing sea water temperature. The  $T_{si}$  is calculated with a thermodynamic model with  $T_a$  as boundary condition. The thermodynamic model is presented in the next section.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$S_{\text{ice}}$  is estimated using the empirical function of Ryvlin (1974)

$$S_{\text{ice}} = S_w(1 - S_R)e^{-a\sqrt{d_{\text{ice}}}} + S_R S_w, \quad (2)$$

where  $S_w$  is the SSS,  $d_{\text{ice}}$  is the ice thickness (here in cm),  $S_R$  is the salinity ratio of the bulk ice salinity at the end of the ice growth season and the SSS,  $a$  is the growth rate coefficient which varies from 0.35 to 0.5. Ryvlin (1974) suggests to use 0.5 for  $a$  and 0.13 for  $S_R$ . However, Kovacs (1996) compared the Ryvlin empirical equation with observed data in the Arctic and suggests to use 0.175 for  $S_R$  instead of 0.13. In our model we use 0.175 for  $S_R$  which seems to fit better to the observation data in the Arctic. Cox and Weeks (1983) give another empirical relationship between  $S_{\text{ice}}$  and  $d_{\text{ice}}$  in the Central Arctic. The two empirical relationships have similar values for first year ice and a water salinity of  $S_w = 31 \text{ g kg}^{-1}$  (Kovacs, 1996). The  $S_{\text{ice}}$  in Eq. (2) is a function of the underlying SSS, and can therefore be applied to regions outside the central Arctic where SSS is much lower.

The ice thickness retrieval with SMOS data is limited by the saturation of TB. We consider TB to reach saturation if the change of TB with  $d_{\text{ice}}$  is less than 0.1 K per cm. Thus, TB of an ice layer with a  $T_{\text{ice}}$  of  $-2^\circ\text{C}$  and a salinity of  $8 \text{ g kg}^{-1}$  reaches its saturation for ice thicknesses of less than 20 cm, for example. This means that the maximal retrievable ice thickness  $d_{\text{max}}$  under warmer conditions can be as low as a few centimeter. On the contrary, under cold conditions and a low ice salinity, which is typical for coastal regions with river run-off, L-band TB emanates from a thicker ice layer. TB reaches its saturation much more slowly and  $d_{\text{max}}$  can be as high as 1.5 m (Figs. 5 and 6). Therefore, SMOS ice thickness retrieval is more suitable for cold conditions and low ice salinity. If the ice temperature varies between  $-5^\circ\text{C}$  and  $-10^\circ\text{C}$ , which is typical for the Arctic in winter, the difference of retrieved ice thicknesses can be as high as 20 cm. The influence of ice salinity on the ice thickness retrieval increases with decreasing ice salinity (Maaß, 2013a). For example, under a  $T_{\text{ice}}$  of  $-10^\circ\text{C}$  the  $d_{\text{max}}$  at  $1 \text{ g kg}^{-1}$   $S_{\text{ice}}$  can be twice of that at  $5 \text{ g kg}^{-1}$   $S_{\text{ice}}$ .

## 3.2 The thermodynamic model

In Algorithm II  $T_{ice}$  is estimated at each step from the  $d_{ice}$  and  $T_a$ . For this purpose thermal equilibrium is assumed at the surface of the ice layer and the heat fluxes are calculated with a thermodynamic model based on Maykut (1986). Although we neglect snow layer in the sea ice radiation model, we consider its thermal insulation effect in the thermodynamic model when we calculate the  $T_{ice}$ . It is shown in Maaß et al. (2013b) that the impact of a snow layer on the TB is partly caused by its insulation effect on the ice temperature. The insulation effect of a snow layer increases with snow thickness. Linear temperature gradient profiles are assumed for the ice and snow layers in the model.

Under the assumption of thermal equilibrium, the incoming and outgoing heat fluxes compensate each other. During winter season, surface melting can be neglected. Therefore the heat balance at the surface of a slab ice layer with thickness  $d_{ice}$  and a layer of snow with thickness  $h_s$  on top can be described as

$$(1 - \alpha)F_r - I_0 + F_{Lin} - F_{Lout} + F_s + F_e + F_c = 0 \quad (3)$$

where  $F_r$  is the incoming shortwave radiation,  $\alpha$  is the albedo of the snow/ice layer,  $I_0$  is the part of the incoming shortwave radiation that is transmitted into the ice,  $F_{Lin}$  is the incoming longwave radiation,  $F_{Lout}$  is the outgoing longwave radiation,  $F_s$  is the sensitive heat flux,  $F_e$  is the latent heat flux, and  $F_c$  is the conductive heat flux.

The radiative and turbulent fluxes  $(1 - \alpha)F_r - I_0$ ,  $F_{Lin}$ ,  $F_{Lout}$ ,  $F_e$ , and  $F_s$  are calculated as in Maykut (1986). For simplification we assume constant values for the cloud cover  $C$ , the relative humidity  $r$ , and the bulk transfer coefficients for sensible and latent heat flux  $C_s$  and  $C_e$  estimated from the reanalysis data. However, these parameters can be obtained from the auxiliary data that will be delivered with SMOS L1C data in the future.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The conductive heat flux  $F_c$  is given by

$$F_c = \frac{k_i k_s}{k_i h_s + k_s d_{ice}} (T_w - T_s) \quad (4)$$

where  $k_s$  is the thermal conductivity of snow,  $T_w$  is the freezing point of sea water.  $k_s$  is set to  $0.31 \text{ W m}^{-1} \text{ K}^{-1}$  according to Yu and Rothrock (1996). The thermal conductivity of ice  $k_i$  can be expressed as (Untersteiner, 1964)

$$k_i = 2.034 + 0.13 \frac{S_{ice}}{T_{ice} - 273}, \quad (5)$$

where  $S_{ice}$  is in  $\text{g kg}^{-1}$  and  $T_{ice}$  is in K.  $T_{ice}$  can be calculated with

$$T_{ice} = 0.5(T_{si} + T_w), \quad (6)$$

where  $T_{si}$  is the snow-ice interface temperature calculated with

$$T_{si} = \frac{T_s + \frac{k_i h_s}{k_s d_{ice}} T_w}{1 + \frac{k_i h_s}{k_s d_{ice}}}. \quad (7)$$

To calculate  $T_{si}$  we need to know  $k_i$ . However,  $k_i$  is in turn a function of  $T_{ice}$ . As an approximation we first calculate  $k_i$  with  $0.5(T_s + T_w)$  instead of  $T_{ice}$  and use this  $k_i$  to calculate  $T_{ice}$  and  $T_{si}$ .  $T_s$  is estimated with leastsquare method for each  $d_{ice}$  under thermal equilibrium assumption.

### 3.3 Retrieval steps

As discussed in Sect. 3.1, the challenge of using variable  $T_{ice}$  and  $S_{ice}$  in the Algorithm II is that both of them are in turn functions of  $d_{ice}$ . The algorithm is based on the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



forward model consisting of the radiation and thermodynamic model. Therefore, we approximate  $d_{ice}$  by repeating the radiation and the thermodynamic model until a convergence point is found for the solution (Fig. 7). In this process, at each step  $T_{ice}$  and  $S_{ice}$  are calculated for the respective  $d_{ice}$  approximation. The starting point of the iteration is the  $d_{ice}$  retrieved with Algorithm I, which uses a constant  $T_{ice}$  of  $-7^{\circ}\text{C}$  and  $S_{ice}$  of  $8\text{ g kg}^{-1}$ . At each iteration step, we use  $d_{ice}$ ,  $T_{ice}$ , and  $S_{ice}$  to calculate TB with the radiation model. The calculated TB is then compared with that observed by SMOS. To minimize the difference between the observed and the calculated TBs, the new  $d_{ice}$  is estimated with a linear approximation method. We define two stopping criteria for the iteration: a brightness temperature difference of less than 0.1 K, or an ice thickness difference of less than 1 cm. The first criterion represents half of the optimal accuracy of daily averaged measurements. We apply the first criterion if the ice is thicker than 30 cm and otherwise the second criterion. The  $d_{max}$  is determined with the same criteria for the saturation of TB, i.e. that the TB change is less than 0.1 K per 1 cm  $d_{ice}$ . We define a saturation factor

$$S_{TB} = d_{ice}/d_{max}. \quad (8)$$

If the saturation factor reaches 100 %, it indicates that the  $d_{max}$  can be considered as the minimum ice thickness of the pixel.

## 4 Assessment of uncertainties

### 4.1 Systematic errors

In both algorithms we assume 100 % ice coverage for simplicity. TB over ice-sea water mixed areas can be described as

$$TB = TB_{water} \times (1 - IC) + TB_{ice} \times IC, \quad (9)$$

where IC is the ice concentration,  $TB_{\text{water}}$  and  $TB_{\text{ice}}$  are the TBs over sea water and ice, respectively.

SMOS  $TB_{\text{water}}$  shows a stable value of about 100.5 K with a standard deviation of about 1 K in the Arctic region. With this constant  $TB_{\text{water}}$ , we can calculate  $TB_{\text{ice}}$  using ice concentration charts from passive microwave radiometer data. During the winter most of the ice covered area in the Arctic has IC higher than 90 % (Andersen et al., 2007). The passive microwave radiometer IC charts have an uncertainty of 5 % in the winter time (Andersen et al., 2007). At high concentrations, correcting the retrieved ice thickness with IC data set with an uncertainty of 5 % can cause higher errors than the 100 % ice coverage assumption. Therefore, we assume in the retrievals a 100 % ice coverage. The possible underestimation of ice thickness due to this assumption is investigated with the simple semi-empirical function used in the Algorithm I. Figure 8 shows that the bias caused by this assumption increases exponentially with decreasing ice concentration. If we assume a SMOS TB of 220 K, the bias can be very high even for IC of more than 80 %. At lower brightness temperatures the bias caused by this assumption is less than a few centimeters.

## 4.2 Sea ice thickness uncertainties

There are several factors that cause uncertainties in the sea ice thickness retrieval: the uncertainty of the SMOS TB, the uncertainties of the auxiliary data sets, and the assumptions made for the radiation and thermodynamic models.

For our purpose we average TB over the incidence angle range of 0–40°. Due to the large amount of measurements in one day this can significantly reduce the uncertainty of TB by dividing standard deviation of TB with square root of the number of measurements during one day at each pixel. This uncertainty is less than 0.5 K in the Arctic except for the strongly RFI affected regions. The uncertainties of  $T_{\text{ice}}$  and  $S_{\text{ice}}$  depend on the uncertainties in the  $T_a$  and SSS. Both  $T_a$  and SSS are derived from model outputs. Due to the sparse observations in the polar regions  $T_a$  and SSS themselves contain large uncertainties.

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A first estimation of SMOS retrieved ice thickness uncertainty is made with Algorithm II based on the  $\text{std}(\text{TB})$ ,  $\text{std}(T_{\text{ice}})$  and  $\text{std}(S_{\text{ice}})$ . The  $\text{std}(\text{TB})$  is calculated at each pixel with standard deviation of all available TB measurements divided by the  $\text{sqrt}(\text{number of TB measurements})$  for each day. The  $\text{std}(S_{\text{ice}})$  is calculated based on the  $\text{std}(\text{SSS})$  chart (see Fig. 4) and  $d_{\text{ice}}$ . The estimation of  $\text{std}(T_{\text{ice}})$  is rather difficult because it depends not only on the  $T_a$  but also the assumptions made in the thermodynamic model. As a first approximation we assume 1 K for the  $\text{std}(T_{\text{ice}})$  which is estimated with the variations in  $T_a$ . More investigations should be conducted to better estimate the uncertainty in  $T_{\text{ice}}$  in the future.

In Table 1 we show an example of estimated ice thickness uncertainties for conditions where  $T_{\text{ice}}$  varies from  $-10^\circ\text{C}$  to  $-2^\circ\text{C}$  and  $S_{\text{ice}}$  varies from  $2\text{ g kg}^{-1}$  to  $8\text{ g kg}^{-1}$ . We assume a standard deviation of 0.5 K, 1 K, and  $1\text{ g kg}^{-1}$  for TB,  $T_{\text{ice}}$ , and  $S_{\text{ice}}$ , respectively. The ice thickness uncertainty caused by  $\text{std}(\text{TB})$  is rather small for thin ice less than 50 cm, and increases exponentially for thicker ice. The uncertainty caused by  $\text{std}(T_{\text{ice}})$  is higher than that caused by  $\text{std}(\text{TB})$  with an increasing trend with increasing ice thickness.  $S_{\text{ice}}$  uncertainty has little impact on the ice thickness retrieval for saline ice with a  $S_{\text{ice}}$  of more than  $5\text{ g kg}^{-1}$ . However, for less saline ice, which is typical for example in the regions with river run-off,  $\text{std}(S_{\text{ice}})$  has much more impact on the ice thickness uncertainty than the other two parameters for  $d_{\text{ice}}$  less than half a meter.

## 5 The effect of the subpixel-scale heterogeneity on the thickness retrieval (Algorithm II\* post-processing)

Natural sea ice is usually not a uniform layer of level ice with a plane geometry, as it was assumed in the emissivity model, but behaves fractally on a wide range of scales. Sea ice deformation patterns are often described using self-similar functions such as the lognormal distribution (Erlingsson, 1988; Key and McLaren, 1991; Tan et al., 2012). A theory of sea ice thickness distribution was developed by Thorndike et al. (1975). Models which include ice growth and deformation may be used to simulate the evolu-

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tion of the thickness distribution (Thorndike, 1992; Godlovitch et al., 2012). A common feature of simulations and empirical observations is the exponential tail resulting from dynamic deformation processes. The inherent skewness of the thickness distribution results in a considerable underestimation of sea ice thickness when the retrieval model is based on a plane sea ice layer. In the following we use airborne sea ice thickness measurements in order to parameterise the thickness distribution function and to investigate the effect of the subpixel-scale heterogeneity on the thickness retrieval.

NASA's Operation IceBridge (OIB) airborne campaigns obtained large scale profiles of sea ice thickness derived from a laser altimeter system (Kurtz et al., 2013). The footprint size of a single laser beam is about 1 m and the vertical accuracy is given as 6.6 cm. The sea ice thickness is estimated from the freeboard by accounting for the snow thickness and assumptions about the densities of ice and snow. The snow thickness is retrieved using a snow-depth radar simultaneously. Here we use the OIB "quicklook" data as obtained from a NSIDC website.

We assume that the sea ice thickness follows a lognormal distribution

$$p(d_{\text{ice}}, \mu, \sigma) = \frac{1}{d_{\text{ice}} \sigma \sqrt{2\pi}} e^{-\frac{(\log(d_{\text{ice}}) - \mu)^2}{2\sigma^2}} \quad (10)$$

with the two parameters logmean  $\mu$  and logsigma  $\sigma$ . Furthermore, we assume a constant logsigma value  $\sigma$  to approximate the thickness distribution function with only one independent variable. To test this assumption we split the 2012 and 2013 OIB Arctic sea ice thickness data in chunks of about 30 km length. We found that using constant values  $\sigma = 0.6 \pm 0.1$  rejects less than 15 % of the chunks tested with Kolmogorov-Smirnov statistics at a significance level of 95 %. The parameter  $\sigma$  increases with increasing length of the chunks and converges to about 0.7 for the maximal number of samples. The parameter  $\sigma$  changed only slightly from 0.692 to 0.695 while the mean thickness decreased considerably from 3.1 m to 2.2 m when considering the entire data sets of the years 2012 and 2013, respectively (Fig. 9). One percent of the 2012 thickness data ( $N_{1\%} = 3430$ ) are above 10 m, and one permil exceeds 16 m with a maximum thickness

value as large as 27.4 m which justifies the exponential tail of the distribution function. The effect of the ice thickness distribution on TB is taken into account by the integration over the thickness range according to the superposition principle

$$TB^*(d_{ice}) = \int_0^{\max(d_{ice})} TB(d_{ice})g(d_{ice})dd_{ice} \quad (11)$$

with the thickness distribution function  $g(d_{ice})$  and the brightness temperature of a single/plane-layer model  $TB(d_{ice})$ . The brightness temperature weighted with the thickness distribution  $TB^*$  suggests a sensitivity to ice thicknesses larger than  $d_{max}$ . Here  $d_{max}$  and  $d_{ice}$  both refer to the single-layer thickness. The real mean thickness denoted as  $H$  is strongly underestimated if the retrieval does not account for the thickness distribution. The overall effect can be explained as an apparently deeper penetration depth caused by the leading edge of the thickness distribution. The implementation of a radiative transfer model that includes this effect is straightforward but computationally expensive because of the integration. A post-processing look-up table for the single-layer model has been generated to estimate an approximate correction factor. This method that converts the single-layer thickness  $d_{ice}$  to the mean thickness  $H$  is called Algorithm II\* hereinafter. Figure 10 shows that the involved correction factor increases with increasing salinity and temperature.

A main uncertainty is the shape of the thickness distribution and its parameterisation with a constant  $\sigma$ . This seems to be a reasonably good representation of the IceBridge thickness data. However, the data in the important thin ice range is only sparse. More field data are required to further analyse the thickness distribution for thin ice on different scales.

SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 6 Comparison of ice thicknesses retrieved with Algorithms I, II, and II\*

In this section, we analyse the time series of ice thicknesses retrieved from the Algorithm I, II, and II\* at single grid points in the Laptev Sea and the Beaufort Sea (Point 1: 77.5° N, 137.5° E, Point 2: 71.0° N, 165.0° W, Point 3: 74.5° N, 127.0° E). The time series begin on 15 October 2011. The time series of ice thickness extracted from two different sea ice assimilation systems are included for comparison. In addition, we show time series of SMOS TB together with ice concentration and derived snow/ice surface temperature.

One of the assimilation systems is the TOPAZ system. TOPAZ is an advanced data assimilation system, using the Ocean model HYbrid Coordinate Ocean Model (HYCOM) and Elastic-Viscous-Plastic (EVP) ice rheology (Bertino and Lisæter, 2008). TOPAZ has a resolution between 18 and 36 km with 22 isopycnal layers. The assimilated observations are satellite-observed Sea Level Anomaly (SLA), Sea Surface Temperature (SST), sea ice concentrations from AMSR-E, sea ice drift products from CERSAT, and Coriolis in-situ temperature and salinity profiles. The TOPAZ system has been in operation since 1 January 2003. The major outcomes in terms of products are weekly issued short term forecasts.

The other assimilation system is the Panarctic Ice Ocean Modeling and Assimilation System (PIOMAS) (Zhang and Rothrock, 2003). It is based on a coupled ocean-ice model forced with National Centers for Environmental Prediction Atmospheric Reanalysis data. PIOMAS assimilates satellite-observed sea ice concentration and sea surface temperature data.

At Point 1, which is located in the north of the Laptev Sea, in the first 30 days Algorithm I and II show very similar  $d_{ice}$  ranging from 0 m to about 0.3 m (Fig. 11). The TB increases from about 100 K to about 230 K. In this TB range,  $d_{ice}$  is the dominant factor of TB variation (Kaleschke et al., 2012). In the next 30 days, TB increases to about 240 K, whereas  $d_{ice}$  increases from about 0.3 m to about 0.4 m in Algorithm I and to more than 0.5 m in Algorithm II. From mid-December to the end of April, the TB shows

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



little variability with a mean value of 237.4 K and a standard deviation of 1.9 K. In this period,  $d_{ice}$  from Algorithm I shows a stable value around 0.35 m with a standard deviation of 3 cm, which results from the constant parameters assumed in Algorithm I. On the contrary,  $d_{ice}$  from Algorithm II shows an average value of 0.48 m with a standard deviation of 11 cm. The strong variability in  $d_{ice}$  is mainly caused by  $T_{ice}$ . A correlation coefficient  $R$  of  $-0.7$  can be found between  $T_{ice}$  and  $d_{ice}$ . In the total time period of 200 days, the  $d_{ice}$  from Algorithm II is on average 10 cm thicker than that from Algorithm I. The ice thickness corrected with the thickness distribution function (Algorithm II\*) is about two times that of Algorithm II.

Simulated ice thicknesses from TOPAZ and PIOMAS show continuous ice growth during the time period, however, with more than half a meter span between them (shaded area in the upper panel of Fig. 11). The ice thicknesses retrieved with Algorithm II\* correspond well with those from TOPAZ and PIOMAS in the first three months. However, from March to April TOPAZ and PIOMAS show further growth in the ice thickness, whereas SMOS shows rather constant or decreasing trends. The decreasing trend in  $d_{ice}$  corresponds to the decreasing  $d_{max}$  caused by the increasing  $T_s$ .

Point 2 is located in the Beaufort Sea, near Barrow. The first sea ice occurrence happens in mid-November, one month later than at Point 1. A few days after the first occurrence of sea ice, the ice concentration rapidly reaches nearly 100 % (Fig. 12). In the following 80 days, the  $T_s$  decreases from about 270 K to 240 K,  $d_{ice}$  retrieved with Algorithm II\* increases from a few centimeters to more than 1.5 m. In this period, the ice thickness growth from SMOS Algorithm II\* agrees well with that simulated by TOPAZ and PIOMAS. Just as at point 1, after the three months freeze-up period the SMOS retrieved  $d_{ice}$  reaches its maximum with a decreasing trend in April, which corresponds to the increasing  $T_s$ .

Point 3 is located north of the Lena Delta where frequent formation of polynyas can be observed. The area is characterised by large interannual variations, being the consequence of an enormous freshwater input from the Lena river and ice formation and salt rejection processes taking place in polynyas offshore the fast ice edge. Anticyclonic

wind conditions force the riverine water northwards and result in a stronger density stratification in the eastern Laptev sea during winter. Cyclonic atmospheric circulation deflects the freshwater plume of the Lena river eastward towards the East Siberian Sea, thus causing higher salinity in the eastern Laptev Sea and the area around the West New Siberian (WNS) polynya.

The strong variability of ice thicknesses in SMOS and PIOMAS shows good correlation (Fig. 13). The decrease and increase of ice thicknesses in SMOS and in the model outputs are very likely caused by the drift of thick ice due to wind forcing and thin ice formation in the polynya areas. From March to April, there is a large discrepancy between the model outputs and the SMOS-derived ice thickness. While PIOMAS shows an ice thickness of more than 2 m in April, SMOS-derived ice thickness is less than half a meter.

Sea ice thickness measurements were carried out in this area during helicopter-borne ice thickness surveys performed in the Laptev Sea during the Transdrift (TD) XX campaign in April 2012. The helicopter-borne ice thickness measurements were made with an electromagnetic (EM)-Bird that utilizes the contrast of electrical conductivity between sea water and ice to determine the distance to the ice-water interface (Haas et al., 2009). An additional laser altimeter yields the distance to the uppermost reflecting surface. Hence, the obtained ice thickness is the ice- plus snow thickness from the difference between the laser range and the EM-derived distance. The accuracy over level sea ice is in the order of 10 cm (Pfaffling et al., 2007). Uncertainties in the ice thickness measurements may arise from the assumption that sea ice is a non-conductive medium. Over thin ice, this assumption may be invalid because the conductivity of saline young ice can be significantly higher than that of older first-year or multi-year ice. This can lead to an underestimation of ice thickness.

The survey flight made on 20 April has a length of about 200 km and covers mostly thin ice being formed in the WNS polynya and the Anabar-Lena polynya. A period of strong and consistent offshore winds led to the development of an extensive thin ice zone extending several hundred kilometres offshore. Point 3 is located in the middle of

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the flight track. Therefore, we use the EM-Bird measurements to validate the SMOS derived ice thickness. During the flights, the EM-Bird recorded a total of 46386 measurements with a mean value of 43 cm and a standard deviation of 33 cm. This agrees well with the 31 cm ice thickness from SMOS Algorithm II\*, considering that the EM-Bird derived ice thickness is the sum of the thicknesses of the ice layer and the snow layer on top of it. The comparison shows that in the polynya area SMOS estimates the ice thickness better than TOPAZ or PIOMAS.

After the time series comparison at single points, we compare the daily ice thickness distribution from the three algorithms in the Arctic on 1 February 2013. As can be seen in Fig. 14, the mean ice thickness considerably increases from Algorithm I to Algorithm II\*. In the central Arctic covered with thick multi-year ice, the TB reaches its saturation. Therefore, none of the algorithms can deliver reliable ice thickness information in the thick multi-year ice area. If we consider only the pixels where TB has not reached its saturation, ice thickness from Algorithm II\* is on average 0.82 m, which is about 40 cm thicker than that from Algorithm II and 55 cm thicker than that from Algorithm I. However, the increase of ice thickness varies from region to region, depending on SSS and weather conditions. For example, in the Laptev Sea where the SSS is much lower than that in the central Arctic, the difference between Algorithm II and Algorithm I is as large as half a meter. On the contrary, in parts of the Kara Sea and north of the Barents Sea little change can be observed between Algorithm I and II. The increase of ice thickness in Algorithm II compared to Algorithm I is caused by the deviation of estimated  $T_{ice}$  and  $S_{ice}$  from the constant values assumed in Algorithm I. To investigate the contribution of  $T_{ice}$  and  $S_{ice}$  in the thickness retrieval separately we carried out two tests with the data of 1 February 2013. In the first test  $S_{ice}$  is assumed to be  $8 \text{ g kg}^{-1}$  as in Algorithm I and we vary only  $T_{ice}$ . In the second test  $T_{ice}$  is assumed to be  $-7^\circ\text{C}$  as in Algorithm I and  $S_{ice}$  is calculated from SSS. In both tests we assume a plane ice layer. If we only consider the pixels where TB has not reached its saturation, the change of ice thickness caused by  $T_{ice}$  in Test 1 varies from  $-10 \text{ cm}$  to more than  $50 \text{ cm}$ , with an average of  $11 \text{ cm}$ . Larger change is found where cold air temperatures prevail. The ice

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



thickness change caused by  $S_{ice}$  from Test 2 is on average 3 cm. However, up to 20 cm and 60 cm difference can be found in the Laptev Sea and in the Baltic Sea.

The comparisons show that Algorithm II\* has a considerably better agreement with the model outputs and the EM-Bird validation data than Algorithm I and II. Taking into account the variability of ice temperature and ice salinity delivers better information about the Arctic-wide ice thickness distribution. Therefore, we use Algorithm II\* to retrieve ice thickness in our operational data processing.

## 7 Ice thickness growth and distribution as seen by SMOS during the freeze-up period

SMOS-derived ice thickness shows continuous growth and expansion of first year ice in the Arctic during the freeze-up period. Figure 15 shows the monthly mean sea ice thickness from October 2012 to March 2013 retrieved with Algorithm II\*. From October to November, thin first-year ice extends to most areas of the East Siberian Sea, the Laptev Sea, and the Beaufort Sea. In addition to the area expansion, also an increase of ice thickness due to the thermodynamic growth can be observed. In December, first-year ice reaches a thickness of more than 1 m in the Laptev Sea and the Beaufort Sea. In March 2013 large areas of thin ice with a thickness less than 40 cm are observed in the Beaufort Sea which is caused by the opening of leads and polynyas in this period.

## 8 Comparison of SMOS and MODIS ice thickness charts in the Kara Sea

### 8.1 Sea ice thickness derived from MODIS data

For the initial verification of SMOS-retrieved sea ice thickness we use MODIS ice thickness charts for the Kara Sea. The validation area extends over 1500 km by 1350 km. The area is suitable for SMOS ice thickness validation because even in the winter time this area is frequently covered by thin first-year ice, which SMOS can best detect. To

compare SMOS and MODIS ice thicknesses, we reduce the 1 km spatial resolution of the MODIS thickness charts to the NSIDC grid resolution of 12.5 km by spatial averaging.

We first compare ice thickness distributions from SMOS and MODIS for two selected days (26 December 2010 and 2 February 2011), on which a sufficient amount of pixels with valid MODIS data is available. After that, we collect all pixels with valid MODIS data from 30 days during the two winter seasons 2009–2010 and 2010–2011 and carry out a pixel to pixel comparison. The 30 days are selected manually. MODIS ice charts with strong cloud limitation are excluded. Similarly to Algorithm I and II, the MODIS sea ice thickness retrieval assumes a plane ice layer. Therefore, by spatial averaging of MODIS data to a grid resolution of 12.5 km we use the modal mean of the MODIS ice thickness instead of the arithmetic mean. For the comparison we use the plane layer SMOS ice thickness, not the inhomogeneous mean ice thickness of Algorithm II\*.

## 8.2 Daily comparison

Figure 16 shows the averaged MODIS ice thickness in a 12.5 km grid resolution, the SMOS ice thicknesses retrieved from Algorithm I and II, and the histogram of the three ice thickness data in the Kara Sea on 26 December 2010. Ice concentration from the same day (Fig. 17) shows near 100 % ice coverage in the ice-covered area except for the marginal ice zone. We use here the ice concentration maps derived from SSM/I with the ARTIST Sea Ice (ASI) algorithm. Both SMOS and MODIS show similar patterns of thin and thick ice distributions, whereas SMOS ice thickness from Algorithm I is considerably lower than the other two in the thicker ice range. Surface air temperature over the ice covered area varies from  $-30$  to  $-20^{\circ}\text{C}$  (Fig. 17), providing favorable conditions for both SMOS and MODIS ice thickness retrievals (Kaleschke et al., 2010; Yu and Rothrock, 1996).

The insulation effect of snow is considered in the SMOS Algorithm II and in the MODIS ice thickness retrieval, but not in the SMOS Algorithm I. Surface temperature and ice thickness are retrieved simultaneously in Algorithm II with the surface air tem-

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



perature as a boundary condition. The SMOS-derived snow surface temperature is in good agreement with that from the MODIS snow/ice surface temperature product (Hall et al., 2004) (Fig. 17). The mean surface temperatures from MODIS and SMOS are both 247 K, and the root mean square deviation (RMSD) is 4 K. Discrepancies can be seen in the marginal ice zone and in the Ob estuary where the low salinities are not well represented by the ocean model. In the marginal ice zone with lower ice concentrations, SMOS strongly underestimates ice thickness, which leads to too warm surface temperatures. The surface temperature is used in SMOS Algorithm II to calculate the bulk ice temperature, which is a variable parameter in the radiation model to calculate the emissivity of an ice layer.

In total there are 4167 pixels in 12.5 km grids with valid MODIS ice thicknesses. For these pixels MODIS has a mean thickness of 44 cm, whereas SMOS has an average of 32 cm and 47 cm from Algorithm I and II, respectively. The correlation coefficient  $R$  and RMSD between the SMOS Algorithm II and MODIS are 0.60 and 20 cm, whereas for SMOS Algorithm I and MODIS they are 0.57 and 23 cm, respectively. If we only consider the 2679 pixels with a MODIS ice thickness less than 50 cm, mean ice thicknesses of SMOS Algorithm I, SMOS Algorithm II and MODIS are 29 cm, 40 cm, and 29 cm respectively. That means in the thin ice range Algorithm II overestimates ice thickness compared to MODIS. The bulk ice temperature derived from the surface temperature in Algorithm II is on average 263.6 K, which is 2.5 K lower than that assumed in SMOS Algorithm I. This can partly explain the ice thickness difference between Algorithm I and II. The SMOS-derived ice thickness decreases with increasing ice temperature at the ice temperature as low as  $-10^{\circ}\text{C}$  (Maaß, 2013a).

Similar results can be derived from another comparison on 2 February 2011 (see Figs. 18 and 19). On this day, large areas of thin ice can be observed from SMOS and MODIS near the Kara Sea strait and in the estuaries. In both regions polynyas appear frequently due to the strong wind forcing. Under cold air temperatures, the polynyas are soon covered by thin ice. Both SMOS and MODIS show ice thicknesses in the range of 20–40 cm in the polynyas with similar distribution patterns. Ice concentration

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is normally higher than 90%, except for the marginal ice zone. Like on 26 December 2010, surface air temperature in the Kara Sea is as low as  $-30^{\circ}\text{C}$ . In total 4016 pixels have valid MODIS data. The mean ice thickness of SMOS Algorithm I, SMOS Algorithm II, and MODIS for the pixels are 33 cm, 50 cm, and 47 cm, respectively. The correlation coefficient and RMSD between the SMOS Algorithm II and MODIS are 0.61 and 21 cm, whereas between SMOS Algorithm I and MODIS they are 0.59 and 26 cm, respectively. The mean surface temperatures from MODIS and SMOS are 246 K and 245 K, with a RMSD of 4 K.

### 8.3 Comparison with 30 days data from the two winter seasons

In total 33 and 87 days of MODIS validation data are available for the winter seasons of 2009–2010 and 2010–2011. However, many of them have only small areas with usable MODIS data. Therefore, we selected out 30 days on that the data are not badly affected by cloud coverage. Altogether 81350 pixels are available in 12.5 km resolution. The histogram of the ice thicknesses (Fig. 20) shows better agreement between SMOS Algorithm II and MODIS than between SMOS Algorithm I and MODIS for these pixels. The mean ice thicknesses derived from SMOS Algorithm II and MODIS are of the same order, 44 cm and 42 cm, respectively, whereas SMOS Algorithm I shows on average 31 cm. If we restrict the comparison to the pixels with MODIS ice thicknesses less than 50 cm, the mean ice thickness from SMOS Algorithm II is about 13 cm higher than the MODIS mean value (see Table 2).

## 9 Conclusions

In this study we developed a new SMOS sea ice thickness retrieval algorithm (denoted as Algorithm II) in which we take into account variations of ice temperature  $T_{\text{ice}}$  and salinity  $S_{\text{ice}}$ .  $T_{\text{ice}}$  and  $S_{\text{ice}}$  are estimated during the ice thickness retrieval from the surface air temperature  $T_{\text{a}}$  of atmospheric reanalysis data and a model-based SSS

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



climatology as boundary conditions. Ice thicknesses retrieved from Algorithm II are compared with that from an earlier semi-empirical algorithm (Kaleschke et al., 2012) (Algorithm I) in which a constant  $T_{ice}$  ( $-7^{\circ}\text{C}$ ) and  $S_{ice}$  ( $8\text{ g kg}^{-1}$ ) are assumed. The new algorithm allows to retrieve considerably higher thickness for cold conditions and less saline ice. The maximal retrievable ice thickness  $d_{max}$  can be estimated based on the  $T_{ice}$  and  $S_{ice}$  at each pixel. In contrast, we estimated  $d_{max}$  about 0.5 m as a constant upper limit for the ice thickness retrieval with Algorithm I. In Algorithm II  $d_{max}$  varies from a few centimeters to about 1 m depending on the  $T_{ice}$  and  $S_{ice}$ . A TB saturation factor is defined as the ratio of  $d_{ice}$  to  $d_{max}$  for each pixel. A saturation ratio close to 100 % indicates that the retrieved ice thickness has to be considered as a minimum ice thickness and that the upper bounds of uncertainty cannot be constrained by the SMOS measurement alone.

Natural sea ice exhibits a broad scale of ice thicknesses within one SMOS footprint due to ice deformation. Algorithm II is based on an emissivity model including a plane layer geometry which is not an adequate assumption for natural sea ice. Therefore, Algorithm II is statistically corrected assuming that the thickness of natural sea ice follows a lognormal distribution. This version of the retrieval is denoted as Algorithm II\*. The statistical correction factor depends on  $T_{ice}$  and  $S_{ice}$ . For warm saline ice the correction factor is higher than for cold less saline ice. The corrected ice thickness is on average about twice as large compared to the plane layer assumption, which is similar to the general relation between modal and mean ice thickness.

The ice thickness from Algorithm II\* agrees well with those from the assimilation systems TOPAZ and PIOMAS in the three months after the first occurrence of sea ice. However, from March to April TOPAZ and PIOMAS have much higher ice thickness compared to the SMOS retrieval. The discrepancy goes along with the onset of surface warming and indicates a possible shortcoming of our retrieval method. We observe a strong impact of  $T_{ice}$  on the ice thickness retrieval when TB approaches saturation. The emissivity model used here does not correctly account for vertical gradients of temperature and salinity. The invalid assumption of a vertically homogeneous ice layer

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

introduces significant uncertainties because the relative brine volume and thus the permittivity depends on ice temperature and salinity (Maaß, 2013a). More work has to be done to develop and test parameterisations that could account for the effects of a vertically structured sea ice cover to further improve the emissivity model. However, a validation with EM-Bird measurements in the polynya areas of the Laptev Sea in April 2012 shows very good agreement between EM-Bird and SMOS ice thicknesses, whereas TOPAZ and PIOMAS overestimate the ice thickness by 0.5–2 m.

For further verification we compare our retrieval results with ice thickness derived from MODIS thermal infrared data in the Kara Sea for several clear sky occasions. The MODIS retrieval relies on a similar plane layer assumption as Algorithm I and II and represents the modal ice thickness. Because of the much larger footprint of SMOS (~40 km) compared to MODIS (1 km) we aggregate the MODIS retrievals on the SMOS grid by taking the modal mean. The different integration times (SMOS: daily averages vs. MODIS: single overpasses) introduce additional uncertainties. Nevertheless, the ice thicknesses retrieved from SMOS and MODIS are very similar, with a considerably better agreement between SMOS Algorithm II and MODIS. The correlation coefficient  $R$  between SMOS and MODIS data is about 0.6 for both Algorithm I and II.

The retrieval uncertainty is dominated by inaccurate assumptions and boundary conditions obtained from auxiliary data, whereas the radiometric accuracy is well constrained and sufficient except for RFI-affected areas. Factors that affect the ice thickness retrieval include the ice concentration, ice salinity, ice temperature, snow thickness as well as the statistical thickness distribution function. Sea ice concentration data available from passive microwave sensors like the Special Sensor Microwave Imager/Sounder (SSMIS) and the Advanced Microwave Scanning Radiometer 2 (AMSR2) have an inherent uncertainty of about 2.5–5 % over high-concentration ice in winter and have strong limitations in new-ice areas (Andersen et al., 2007; Ezraty, 2002). Thus, we do not correct for varying ice concentration because this would considerably increase the noise and raise so far unresolved problems in regions where new ice is not detected with traditional methods but with SMOS. The variability of the parent water

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



salinity contributes only little to the overall uncertainty except for low-salinity areas. By neglecting the growth-rate dependent salt inclusion in Ryvlin's parameterization we introduce additional errors that can not be quantified without the use of more advanced sea ice thermodynamic models (Notz and Worster, 2009; Vancoppenolle et al., 2006).

5 The radiation model used in this study is adapted from Menashi et al. (1993) which is a simple one layer model without a snow layer. Although we consider the insulation effect of snow, the radiative contribution of the snow layer to the overall brightness temperature is neglected. This effect is investigated in Maaß et al. (2013b) with another radiation model based on Burke et al. (1979).

10 A distinct advantage of the SMOS sea ice thickness retrieval is the daily coverage independent of clouds and the large sensitivity for thin ice. Thus, our SMOS product is complementary to the sea ice thickness derived from CryoSat-2 and feasible for operational usage. However, the thickness retrieval is strictly limited to cold periods and not applicable during late spring and summer. Daily SMOS ice thickness charts from 15 October to 15 April since 2010 are available via <http://icdc.zmaw.de>.

*Acknowledgements.* This work was funded through ESA's Support to Science Element Program under contract 4000101476. The authors would like to thank the members of the ESA SMOSIce project for helpful discussions. EM-Bird thickness measurements were conducted by Alfred Wegener Institute for Polar and Marine Research (Thomas Krumpfen and Valeria Sel-yuzhenok) within the Transdrift XX campaign in the Framework of a Russian-German research cooperation. JRA-25 reanalysis data are provided by Japanese Meteorological Agency (JMA), the sea ice concentration data are provided by ICDC University of Hamburg, TOPAZ data are provided by Laurant Bertino from Nansen Environmental and Remote Sensing Center in Nor-way, and PIOMAS data are provided by Jinlun Zhang from University of Washington.

## 25 References

Andersen, S., Tonboe, R., Kaleschke, L., Heygster, G., and Pedersen, L.: Intercomparison of passive microwave sea ice concentration retrievals over the high-concentration Arctic sea ice, *J. Geophys. Res.*, 112, C08004, doi:10.1029/2006JC003543, 2007. 5751, 5764

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Bertino, L. and Lisæter, K. A.: The TOPAZ monitoring and prediction system for the Atlantic and Arctic Oceans, *Journal of Operational Oceanography*, 1, 15–19, 2008. 5755
- Brath, M., Scharffenberg, M. G., Serra, N., and Stammer, D.: Altimeter-based estimates of eddy variability and eddy transports in the subpolar North Atlantic, *Mar. Geod.*, 33, 472–503, 2010. 5744
- Burke, W., Schmugge, T., and Paris, J.: Comparison of 2.8 and 21 cm microwave radiometer observations over soils with emission model calculations, *J. Geophys. Res.*, 84, 287–294, 1979. 5765
- Camps, A., Vall-llossera, N., Duffo, N., Torres, F., and Corbella, I.: Performance of sea surface salinity and soil moisture retrieval algorithms with different auxiliary datasets in 2-D L-band aperture synthesis interferometric radiometers, *IEEE T. Geosci. Remote*, 43, 1189–1200, doi:10.1109/TGRS.2004.842096, 2005. 5742
- Camps, A., Gourrion, J., Tarongi, J., Gutierrez, A., Barbosa, J., and Castro, R.: RFI analysis in SMOS imagery, in: *Geoscience and Remote Sensing Symposium (IGRASS proceedings 2010)*, 2007–2010, 2010. 5742
- Cox, G. and Weeks, W.: Equations for determining the gas and brine volumes in sea-ice samples, *J. Glaciol.*, 29, 306–316, 1983. 5746, 5747
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011. 5744
- Dmitrenko, I. A., Kirillov, S. A., Ivanov, V. V., Rudels, B., Serra, N., and Koldunov, N. V.: Modified halocline water over the laptev sea continental margin: historical data analysis, *J. Climate*, 25, 5556–5565, 2012. 5744
- Doronin, Y.: *Thermal Interaction of the Atmosphere and the Hydrosphere in the Arctic*, CornetBooks, Philadelphia, 1971. 5745
- Drucker, R., Martin, S., and Moritz, R.: Observations of ice thickness and frazil ice in the St. Lawrence Island polynya from satellite imagery, upward look-

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ing sonar, and salinity/temperature moorings, *J. Geophys. Res.-Oceans*, 108, 3149, doi:10.1029/2001JC001213, 2003. 5736

Erlingsson, B.: Two-dimensional deformation patterns in sea ice, *J. Glaciol*, 34, 301–308, 1988. 5752

5 Ezraty, R.: Detection and monitoring of new-ice in the East Greenland Sea using the SeaWinds scatterometer, *Journal of Ocean University of Qingdao*, 1, 109–118, doi:10.1007/s11802-002-0002-z, 2002. 5764

Frey, R. A., Ackerman, S. A., Liu, Y., Strabala, K. I., Zhang, H., Key, J. R., and Wang, X.: Cloud detection with MODIS – Part 1: Improvements in the MODIS cloud mask for collection 5, *J. Atmos. Ocean. Tech.*, 25, 1057–1072, 2008. 5737, 5745

10 Godlovitch, D., Monahan, A., and Flato, G.: An idealised stochastic model of sea ice thickness dynamics, *Cold Reg. Sci. Technol.*, 78, 14–30, 2012. 5753

Haas, C., Lobach, J., Hendricks, S., Rabenstein, L., and Pfaffling, A.: Helicopter-borne measurements of sea ice thickness, using a small and lightweight, digital EM system, *J. Appl. Geophys.*, 67, 234–241, 2009. 5757

15 Hall, D., Key, J., Casey, K., Riggs, G., and Cavalieri, D.: Sea ice surface temperature product from MODIS, *IEEE T. Geosci. Remote*, 42, 1076–1087, 2004. 5761

Hwang, B. J., Ehn, J. K., Barber, D. G., Galley, R., and Grenfell, T. C.: Investigations of newly formed sea ice in the Cape Bathurst polynya: 2. Microwave emission, *J. Geophys. Res.-Oceans* (1978–2012), 112, 2007. 5738

20 Indra: SMOS DPGS: SMOS Level 1 and Auxiliary Data Products Specifications, Ref. SO-TN-IDR-GS-0005, 2010. 5741

Janssen, F., Schrum, C., and Backhaus, J.: A climatological data set of temperature and salinity for the Baltic Sea and the North Sea, *Deutsche Hydrographische Zeitschrift*, 51, 5–245, 1999. 5744

25 Kaellen, E.: HIRLAM Documentation Manual, System 2.5. Swed. Meteorol., and Hydrol. Inst., Norrköeping, Sweden, 1996, 1996. 5745

Kaleschke, L., Maaß, N., Haas, C., Hendricks, S., Heygster, G., and Tonboe, R. T.: A sea-ice thickness retrieval model for 1.4 GHz radiometry and application to airborne measurements over low salinity sea-ice, *The Cryosphere*, 4, 583–592, doi:10.5194/tc-4-583-2010, 2010. 5738, 5760

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kaleschke, L., Tian-Kunze, X., Maaß, N., Mäkynen, M., and Drusch, M.: Sea ice thickness retrieval from SMOS brightness temperatures during the Arctic freeze-up period, *Geophys. Res. Lett.*, 39, L05501, doi:10.1029/2012GL050916, 2012. 5738, 5739, 5746, 5755, 5763
- 5 Kerr, Y., Waldteufel, P., Wigneron, J., Martinuzzi, J., Font, J., and Berger, M.: Soil moisture retrieval from space: the Soil Moisture and Ocean Salinity (SMOS) mission, *IEEE T. Geosci. Remote*, 39, 1729–1735, 2001. 5741
- Key, J. and McLaren, A.: Fractal nature of the sea ice draft profile, *Geophys. Res. Lett.*, 18, 1437–1440, 1991. 5752
- 10 Kovacs, A.: Sea Ice. Part 1. Bulk Salinity Versus Ice Floe Thickness, Tech. rep., DTIC Document, 1996. 5747
- Kurtz, N. T., Farrell, S. L., Studinger, M., Galin, N., Harbeck, J. P., Lindsay, R., Onana, V. D., Panzer, B., and Sonntag, J. G.: Sea ice thickness, freeboard, and snow depth products from Operation IceBridge airborne data, *The Cryosphere*, 7, 1035–1056, doi:10.5194/tc-7-1035-2013, 2013. 5753
- 15 Kwok, R. and Cunningham, G.: ICESat over Arctic sea ice: estimation of snow depth and ice thickness, *J. Geophys. Res.*, 113, C08010, doi:10.1029/2008JC004753, 2008. 5736, 5737
- Laxon, S., Peacock, N., and Smith, D.: High interannual variability of sea ice thickness in the Arctic region, *Nature*, 425, 947–950, 2003. 5736, 5737
- Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., Hendricks, S., Krishfield, R., Kurtz, N., Farrell, S., and Davidson, M.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophys. Res. Lett.*, 40, 732–737, doi:10.1002/grl.50193, 2013. 5736
- 20 Maaß, N.: Remote sensing of sea ice thickness using SMOS data, Reports on Earth System Science, available at: [http://www.mpimet.mpg.de/fileadmin/publikationen/Reports/WEB\\_BzE\\_131.pdf](http://www.mpimet.mpg.de/fileadmin/publikationen/Reports/WEB_BzE_131.pdf), 2013a. 5746, 5747, 5761, 5764
- 25 Maaß, N., Kaleschke, L., Tian-Kunze, X., and Drusch, M.: Snow thickness retrieval over thick Arctic sea ice using SMOS satellite data, *The Cryosphere Discuss.*, 7, 3627–3674, doi:10.5194/tcd-7-3627-2013, 2013. 5748, 5765
- Maekynen, M., Cheng, B., and Similae, M.: On the accuracy of thin-ice thickness retrieval using MODIS thermal imagery over Arctic first-year ice, *Ann. Glaciol.*, 54, 87–96, doi:10.3189/2013AoG62A166, 2013. 5737, 5745
- 30

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Marshall, J., Adcroft, A., Hill, C., Perelman, L., and Heisey, C.: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers, *J. Geophys. Res.*, 102, 5753–5766, 1997. 5744
- Martin, S., Drucker, R., Kwok, R., and Holt, B.: Improvements in the estimates of ice thickness and production in the Chukchi Sea polynyas derived from AMSR-E, *Geophys. Res. Lett.*, 32, L05505, doi:10.1029/2004GL022013, 2005. 5737
- Maykut, G. A.: The surface heat and mass balance, in: *Geophysics of Sea Ice*, 395–463, Plenum, New York, 1986. 5748
- Mecklenburg, S., Drusch, M., Kerr, Y. H., Font, J., Martin-Neira, M., Delwart, S., Buenadicha, G., Reul, N., Daganzo-Eusebio, E., Oliva, R., and Crapolicchio, R.: ESA's soil moisture and ocean salinity mission: mission performance and operations, *IEEE T. Geosci. Remote*, 50, 1354–1366, 2012. 5738, 5741
- Menashi, J., Germain, K., Swift, C., Comiso, J., and Lohanick, A.: Low-frequency passive-microwave observations of sea ice in the Weddell Sea, *J. Geophys. Res.*, 98, 22569–22577, 1993. 5738, 5739, 5746, 5765
- Naoki, K., Ukita, J., Nishio, F., Nakayama, M., Comiso, J. C., and Gasiewski, A.: Thin sea ice thickness as inferred from passive microwave and in situ observations, *J. Geophys. Res.*, 113, C02S16, doi:10.1029/2007JC004270, 2008. 5737
- Nihashi, S., Ohshima, K. I., Tamura, T., Fukamachi, Y., and Saitoh, S.-i.: Thickness and production of sea ice in the Okhotsk Sea coastal polynyas from AMSR-E, *J. Geophys. Res.-Oceans*, 114, C10025, doi:10.1029/2008JC005222, 2009. 5737
- Notz, D. and Worster, M. G.: Desalination processes of sea ice revisited, *J. Geophys. Res.-Oceans*, 114, C05006, doi:10.1029/2008JC004885, 2009. 5765
- NSIDC: Morphometric characteristics of ice and snow in the Arctic Basin: aircraft landing observations from the Former Soviet Union, 1928–1989, Compiled by I. P. Romanov, National Snow and Ice Data Center, Boulder, CO, Digital media, 2004. 5745
- Onogi, K., Tsutsumi, J., Koide, H., Sakamoto, M., Kobayashi, S., Hatsushika, H., Matsumoto, T., Yamazaki, N., Kaalhoru, H., Takahashi, K., Kadokura, S., Wada, K., Kato, K., Oyama, R., Ose, T., Mannoji, N., and Taira, R.: The JRA-25 reanalysis, *J. Meteorol. Soc. Jpn.*, 85, 369–432, 2007. 5743
- Pfaffling, A., Haas, C., and Reid, J. E.: Direct helicopter EM-Sea-ice thickness inversion assessed with synthetic and field data, *Geophysics*, 72, F127–F137, 2007. 5757

- Ryvlin, A. I.: Method of forecasting flexural strength of an ice cover, *Probl. Arct. Antarct.*, 45, 79–86, 1974. 5739, 5747
- Serra, N., Käse, R. H., Köhl, A., Stammer, D., and Quadfasel, D.: On the low-frequency phase relation between the Denmark Strait and the Faroe-Bank Channel overflows, *Tellus A*, 62, 530–550, doi:10.1111/j.1600-0870.2010.00445.x, 2010. 5744
- 5 Singh, R. K., Oza, S. R., Vyas, N. K., and Sarkar, A.: Estimation of thin ice thickness from the advanced microwave scanning radiometer-EOS for coastal polynyas in the Chukchi and Beaufort Seas, *IEEE T. Geosci. Remote*, 49, 2993–2998, 2011. 5737
- Tamura, T. and Ohshima, K. I.: Mapping of sea ice production in the Arctic coastal polynyas, *J. Geophys. Res.-Oceans*, 116, C05006, doi:10.1029/2008JC004885, 2011. 5737
- 10 Tamura, T., Ohshima, K. I., Markus, T., Cavalieri, D. J., Nishashi, S., and Hirasawa, N.: Estimation of thin ice thickness and detection of fast ice from SSM/I data in the Antarctic Ocean, *J. Atmos. Ocean. Tech.*, 24, 1757–1772, doi:10.1175/JTECH2113.1, 2007. 5737
- Tan, B., Li, Z.-j., Lu, P., Haas, C., and Nicolaus, M.: Morphology of sea ice pressure ridges in the northwestern Weddell Sea in winter, *J. Geophys. Res.-Oceans*, 117, C06024, doi:10.1029/2011JC007800, 2012. 5752
- 15 Thorndike, A.: Estimates of sea ice thickness distribution using observations and theory, *J. Geophys. Res.*, 97, 12601–12, 1992. 5753
- Thorndike, A., Rothrock, D., Maykut, G., and Colony, R.: The thickness distribution of sea ice, *J. Geophys. Res.*, 80, 4501–4513, 1975. 5752
- 20 Uden, P. e. a.: HIRLAM-5 scientific documentation, *Swed. Meteorol., and Hydrol. Inst., Norrköping, Sweden*, 2002, 2002. 5745
- Untersteiner, N.: Calculations of temperature regime and heat budget of sea ice in the Central Arctic, *J. Geophys. Res.*, 69, 4755–4766, 1964. 5749
- 25 Vancoppenolle, M., Fichet, T., and Bitz, C. M.: Modeling the salinity profile of undeformed Arctic sea ice, *Geophys. Res. Lett.*, 33, L21501, 2006. 5765
- Vant, M., Ramseier, R., and Makios, V.: The complex-dielectric constant of sea ice at frequencies in the range 0.1–40 GHz, *J. Appl. Phys.*, 49, 1264–1280, 1978. 5746
- Yu, Y. and Rothrock, D.: Thin ice thickness from satellite thermal imagery, *J. Geophys. Res.*, 101, 25753–25766, 1996. 5736, 5737, 5749, 5760
- 30 Zhang, J. and Rothrock, D.: Modeling global sea ice with a thickness and enthalpy distribution model in generalized curvilinear coordinates, *Mon. Weather Rev.*, 131, 845–861, 2003. 5755

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.

**Table 1.** Estimated ice thickness uncertainties caused by  $\text{std}(\text{TB})$ ,  $\text{std}(T_{\text{ice}})$ , and  $\text{std}(S_{\text{ice}})$ .

$d_{\text{ice}}$	$\text{std}(\text{TB}) = 0.5 \text{ K}$	$\text{std}(T_{\text{ice}}) = 1 \text{ K}$	$\text{std}(S_{\text{ice}}) = 1 \text{ g kg}^{-1}$
0–10 cm	less than 1 cm	less than 1 cm	less than 1 cm
10–30 cm	less than 1 cm	1–5 cm	1–13 cm
30–50 cm	1–4 cm	2–10 cm	2–22 cm
more than 50 cm	4 cm–more than 1 m	7 cm–more than 1 m	up to 40 cm

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.

**Table 2.** Comparison of SMOS and MODIS based ice thicknesses in the Kara Sea during the 30 days selected from the two winter seasons of 2009–2010 and 2010–2011.

		pixels with MODIS ice thickness < 0.5 m	all pixels
Number of pixels		51 716	81 350
mean	MODIS	0.24 m	0.42 m
	SMOS I	0.27 m	0.31 m
	SMOS II	0.37 m	0.44 m
<i>R</i>	SMOS I	0.64	0.61
	SMOS II	0.63	0.62
RMSD	SMOS I	0.11 m	0.25 m
	SMOS II	0.19 m	0.22 m

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 1.** Distribution of radiometric accuracy within a typical snapshot with incidence angles [degree] as contour lines. The snapshot is gridded with 10 km spatial grid resolution.

SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




## SMOS derived sea ice thickness

X. Tian-Kunze et al.



**Fig. 2.** The improvement of RFI-induced data loss in the Arctic from 2010 to 2012.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Fig. 3.** Vertically (V) and horizontally (H) polarized TBs and the first Stokes parameter as a function of incidence angle calculated using a three layer model for sea ice with a thickness of  $d_{\text{ice}} = 1$  m, a bulk salinity of  $S_{\text{ice}} = 8 \text{ g kg}^{-1}$ , and a bulk ice temperature of  $T_{\text{ice}} = -7^\circ\text{C}$ .

SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




## SMOS derived sea ice thickness

X. Tian-Kunze et al.



**Fig. 4.** Mean and standard deviation of weekly sea surface salinity for the winter period from October to April, based on 8 yr of daily model output.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Fig. 5.** TB as function of  $d_{ice}$  under different  $T_{ice}$ , calculated with the sea ice radiation model with a  $S_{ice}$  of  $8 \text{ g kg}^{-1}$

**SMOS derived sea ice thickness**

X. Tian-Kunze et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.






Fig. 6.  $d_{max}$  under different  $T_{ice}$  [°C] and  $S_{ice}$  [ $g\ kg^{-1}$ ].



**Fig. 7.** Schematic flow chart of the retrieval steps.  $d$  and  $d'$  are the sea ice thicknesses from the consecutive steps,  $TB$  and  $TB_{obs}$  are calculated and observed brightness temperatures,  $T_0$  is the brightness temperature of sea water assumed to be 100.5 K.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**Fig. 8.** The underestimation of ice thickness caused by the 100 % ice coverage assumption.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 9.** Sea ice thickness distribution derived from NASA's Operation IceBridge data of 2012 (upper panel,  $\sigma = 0.692$ ) and 2013 (lower panel,  $\sigma = 0.695$ ). The y axis is the number of occurrence.



**Fig. 10.** Relationship between the plane ice layer thickness  $d_{ice}$  and the mean inhomogeneous ice layer thickness  $H$  at different  $T_{ice}$  and  $S_{ice}$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.



**Fig. 11.** Time series of ice thickness derived from Algorithm I, II, and II\*, together with  $d_{max}$  and simulated ice thicknesses from TOPAZ and PIOMAS (upper panel) and time series of ice concentration, snow (or ice in case of bare ice) surface temperature and SMOS TB (lower panel) at Point 1 (77.5° N, 137.5° E).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## SMOS derived sea ice thickness

X. Tian-Kunze et al.



**Fig. 13.** Time series of ice thicknesses derived from Algorithm I, II, and II\*, together with  $d_{max}$  and simulated ice thicknesses from TOPAZ and PIOMAS (upper panel) and time series of ice concentration, snow (or ice in case of bare ice) surface temperature and SMOS TB (lower panel) at Point 3 (74.5° N, 127.0° E).

## SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶


Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 14.** SMOS ice thickness derived from retrieval algorithm I, II, and II\* in the Arctic on 1 February 2013.



**Fig. 15.** Monthly sea ice thickness derived from Algorithm II\* during the freeze-up period of October 2012 to March 2013 (from upper left to lower right) in the Arctic.

**SMOS derived sea ice thickness**

X. Tian-Kunze et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion





**Fig. 16.** The averaged MODIS ice thickness in 12.5 km grid resolution (upper left), SMOS ice thicknesses retrieved from Algorithm I (upper right) and II (lower left), and the histogram of the three ice thickness data (lower right) in the Kara Sea on 26 December 2010.



**Fig. 17.** SSM/I ice concentration (upper left), JRA-25 surface air temperature (upper right), MODIS- and SMOS-based snow/ice surface temperature (lower left and lower right) in the Kara Sea on 26 December 2010.

SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 18.** The averaged MODIS ice thickness in 12.5 km grid resolution (upper left), SMOS ice thicknesses retrieved from Algorithm I (upper right) and II (lower left), and the histogram of the three ice thickness data (lower right) in the Kara Sea on 2 February 2011.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 19.** SSM/I ice concentration (upper left), JRA-25 surface air temperature (upper right), MODIS- and SMOS-based snow/ice surface temperature (lower left and lower right) in the Kara Sea on 2 February 2011.

SMOS derived sea ice thickness

X. Tian-Kunze et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## SMOS derived sea ice thickness

X. Tian-Kunze et al.



**Fig. 20.** Histogram of SMOS (Algorithm I and II) and MODIS (in 12.5 km and 1 km grid resolution) ice thicknesses from all pixels of the 30 days.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)