

# Arctic Budget Study of Inter-member Variability using HIRHAM5 Ensemble Simulations

A. Sommerfeld (1), O. Nikiema (2), A. Rinke (1), K. Dethloff (1), R. Laprise (2)

(1) Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research in Potsdam

(2) Université du Québec à Montréal

corresponding author:

anja.sommerfeld@awi.de

## Introduction

- chaotic and non-linear nature of atmospheric dynamics [1]
- RCMs are sensitive to their initial conditions (IC)
- generation of internal variability within RCMs

1) investigation of internal variability = inter-member variance (IV) [2]

2) estimation of diabatic and dynamical contribution leading to IV

→ diagnostic potential temperature budget equation [3, 4]

3) analysing the high IV event on 5<sup>th</sup> August 2012 at 06 UTC

## Model: HIRHAM5

- hydrostatic regional atmospheric model
- driven by ERA-Interim
- integration area: Arctic
- ensemble with 20 simulations differing in their IC
- initialisation time shifts about 6 hours for each run
- analysed time period: 6<sup>th</sup> July to 30<sup>th</sup> September 2012
- runs without nudging

## 1) Inter-member Variability




Fig. 1: Domain and vertical averaged potential temperature IV




Fig. 2: Spatial distribution of the time averaged potential temperature IV at 925 hPa

## 2a) Diagnostic Potential Temperature Budget Equation

• IV is defined as the inter-member variance of the potential temperature  $\theta$  [3, 4] of the 20 ensemble-members  $n$   $\sigma_{\theta}^2 \approx \langle \theta_n'^2 \rangle$  (Eq. 1)

• emanating from the first law of thermodynamics for potential temperature and the mass-continuity equation in vertical pressure coordinates and applying the Reynolds decomposition

→ the variable  $\theta_n$  split in the ensemble mean  $\langle \theta \rangle$  and the deviation from ensemble mean  $\theta_n'$   $\theta_n = \langle \theta \rangle + \theta_n'$  (Eq. 2)

• results in a IV budget equation (Eq. 3) developed by [3, 4]

$$\frac{\partial \sigma_{\theta}^2}{\partial t} = \underbrace{-\vec{\nabla} \cdot (\langle \vec{V} \rangle \sigma_{\theta}^2)}_{A_h} - \underbrace{\frac{\partial (\langle \omega \rangle \sigma_{\theta}^2)}{\partial p}}_{A_v} - \underbrace{2 \langle \theta_n' \vec{V}_n' \rangle \cdot \vec{\nabla} \langle \theta \rangle}_{B_h} - \underbrace{2 \langle \theta_n' \omega_n' \rangle \frac{\partial \langle \theta \rangle}{\partial p}}_{B_v} + \underbrace{2 \langle \theta_n' J_n' \rangle}_{C} - \underbrace{2 \langle \theta_n' \vec{\nabla} \cdot (\theta_n' \vec{V}_n') \rangle}_{E_h} - \underbrace{2 \langle \theta_n' \frac{\partial}{\partial p} (\theta_n' \omega_n') \rangle}_{E_v} \quad (\text{Eq. 3})$$

diagnostic tendency of potential temperature IV

$A_h$  horizontal transport

$A_v$  vertical transport

$B_h$  horizontal baroclinicity


$B_v$  vertical baroclinicity

$C$  diabatic source/sink term

$E_h$  horizontal third-order term

$E_v$  vertical third-order term

## 2b) Contributions to IV



- $B_h, B_v$ : baroclinic terms most important
- $A_h$ : small on domain average, but important for specific locations
- $C$ : small on domain and vertical average

Fig. 3: Time evolution of the vertical and domain averaged contributions to potential temperature IV tendency




Fig. 4: Spatial distribution of the time averaged horizontal baroclinic term  $B_h$  (left) and vertical baroclinic term  $B_v$  (right) contributing to potential temperature IV tendency at 500 hPa

## 3) High IV event on 5<sup>th</sup> August 2012

• strongest IV event during summer 2012 on 5<sup>th</sup> August at 06 UTC

→ strong baroclinic contribution ( $B_h, B_v$ ) to IV tendency

→ coinciding with the great Arctic cyclone in the beginning of August 2012 [5]

• great Arctic cyclone leads to an intense sea ice loss in East Siberian/Chukchi Sea

→ strong diabatic contribution ( $C$ ) to IV tendency





Fig. 5: Spatial distribution of the potential temperature IV (top) and the horizontal baroclinic term  $B_h$  (left), the vertical baroclinic term  $B_v$  (middle) and the diabatic source and sink term  $C$  (right) contributing to potential temperature IV tendency on 5<sup>th</sup> August 2012 at 06 UTC at 500 hPa

## Outlook

- application of budget study for other years (summer 2006, 2007, 2009)
- investigation of more high IV events
- comparison with the results obtained with the CRCM5 over the Arctic
- dependency of the IV and its contributions on the model structure and physical parameterisations

## References

- [1]: Lorenz, E.N., 1967. *The Nature and Theory of the General Circulation of the Atmosphere*. World Meteorol. Org., 161pp.
- [2]: Alexandru, A. et al., 2007. *Internal Variability in Regional Climate Downscaling at the Seasonal Scale*. Mon Weather Rev. 135, 3221-3238
- [3]: Nikiema, O. and Laprise, R., 2010. *Diagnostic budget study of the internal variability in ensemble simulations of the Canadian RCM*. Clim. Dyn. 36, 2313-2337
- [4]: Nikiema, O. and Laprise, R., 2011. *Budget study of the internal variability in ensemble simulations of the Canadian RCM at the seasonal scale*. J. Geophys. Res. Atmos. 116(D16112)
- [5]: Zhang, J. et al., 2013. *The impact of an intense summer cyclone on 2012 Arctic sea ice retreat*. Geophys. Res. Lett. 40, 720-726