



Master Track RV Polarstern ARK-X/1

Data Processing Report

Contents

1	Introduction	1
2	Workflow	1
3	Cruise details	2
4	Sensor Layout	2
5	Processing Report	3

Contact:

Dr. Rainer Knust

Alfred Wegener Institute

Columbusstrasse, D-27568 Bremerhaven, GERMANY

Tel: +49(471)4831-1709 Fax: +49(471)4831-1918

Mail: Polarstern-Coordination@awi.de

Processing Agency:

FIELAX Gesellschaft für wissenschaftliche Datenverarbeitung mbH

Schleusenstr. 14, D-27568 Bremerhaven, GERMANY

Tel: +49 (0) 471 30015 0 Fax: +49 (0) 471 30015 22 Mail: info@fielax.de

Ref.: ARK-X	1 nav.pdf	Vers.: 1	Date: 2016/02/12	Status: final



1 Introduction

This report describes the processing of raw data acquired by position sensors on board RV Polarstern during expedition ARK-X/1 to receive a validated master track which is used as reference of further expedition data.

2 Workflow

The different steps of processing and validation are visualized in figure 1. Unvalidated data of up to three sensors are extracted from the DAVIS SHIP data base (https://dship.awi.de) in a 1-second interval or 5-second interval for cruises earlier than ARK-IX/2. They are converted to ESRI point shapefiles and imported to ArcGIS. A visual screening is performed to evaluate data quality and remove outliers manually. For all resulting position tracks, a quality check is performed using a ship's speed filter, an acceleration filter and a course-change filter. Filtered positions are flagged. In addition, a manual check is performed to flag obvious outliers. Those position tracks are combined to a single master track depending on a sensor priority list (by accuracy, reliability) and availability / applied exclusion of automatically or manually flagged of data. Missing data up to a time span of 60 seconds are linearly interpolated to achieve a master track with 1-second resolution. To reduce the amount of points for overview maps the master track is generalized by using the Ramer-Douglas-Peucker algorithm. This algorithm returns only the most significant points from the track. Full master track and generalized master track are written to text files and imported to PANGAEA (http://www.pangaea.de) for publication.




Figure 1: Workflow of master track data processing



3 Cruise details

Vessel name RV Polarstern

Cruise name ARK-X/1

Cruise start 06.07.1994 Bremerhaven

Cruise end 16.08.1994 Tromsø

Cruise duration 42 days

4 Sensor Layout

This chapter describes the position sensors mounted during this cruise.

Position sensors

Sensor name	System Position Information, short: System	
Description Position information delivered to the System		

Sensor name	Navigation Automation Control System, short: NACOS	
Description	Navigation system of the ship	

Sensor name	Ashtech Z-12, short: Ashtech	
Description	GPS-Receiver	

Position data from Parasound-surveys

Additionally to the up to three position sensors mounted, there are positions available extracted from the header-information of Parasound-surveys which are already processed and checked for quality and validity. If those data exist for this cruise, these data will be used instead of the derived master track. These data are identified as follows.

Sensor name	Corrected Parasound-Navigation, short: Parasound-NAV	
Description	Already processed Position information from Parasound navigation	



5 Processing Report

Database Extraction

Data source	DSHIP database (dship.awi.de)	
Exported values 3628799		
First dataset	1994-07-06T00:00:00 UTC	
Last dataset	1994-08-16T23:59:58 UTC	

Automatic Validation

The following thresholds were applied for the automatic flagging of the position data:

Speed Maximum 20 kn between two datapoints.	
Acceleration Maximum 1 m/s ² between two datapoints.	
Change of course Maximum 5° between two datapoints.	

Manual Validation

Obvious outliers were removed manually. For details see Processing Logbook of RV Polarstern (hdl:10013/epic.45909).

Flagging result

	System		NACOS		Ashtech	
Missing	286113	7.885%	207509	5.718%	3292790	90.740%
Speed	16692	0.460%	16543	0.456%	1742	0.048%
Acceleration	281261	7.751%	292840	8.070%	28243	0.778%
Course	959387	26.438%	624153	17.200%	42491	1.171%
Manually	0	0.000%	0	0.000%	0	0.000%

Master Track Generation

The master track is derived from the position sensors' data selected by priority.

Sensor priority used:

- 1. System
- 2. NACOS
- 3. Ashtech

Filters applied: manual, speed.

Distribution of position sensor data in master track:



Sensor	Data points	Percentage
Total	3628799	100.000%
Parasound-NAV	0	0.000%
System	3325669	91.647%
NACOS	87283	2.405%
Ashtech	15652	0.431 %
Interpolated	194799	5.368 %
Gaps	5396	0.149%

Remarks

None.

Score

For each cruise, a score is calculated ranging from 0 (no data) to 100 (only very good data). the score for the cruise ARK-X/1 is 85.

Generalization

The master track is generalized to receive a reduced set of the most significant positions of the track using the Ramer-Douglas-Peucker algorithm and allow a maximum tolerated distance between points and generalized line of 4 arcseconds.

Results:

Number of generalized points	5468 points
Data reduction	99.8493%



Result files

Report in XML format:

The XML contains all information of the master track generation in a machine-readable format. In addition a XSD schema file is provided.

Master track text file:

The format is a plain text (tab-delimited values) file with one data row in 1 second interval.

Column separator	Tabulator "\t"		
Column 1	Date and time expressed according to ISO 8601		
Column 3	Latitude in decimal format, unit degree		
Column 4	Longitude in decimal format, unit degree		
Column 5	Flag for da	ta source	
	0	Parasound-NAV	
	1	System	
	2	NACOS	
	3	Ashtech	
	INTERP	Interpolated point	
	GAP	Missing data	

Text file of the generalized master track:

The format is a plain text (tab-delimited values) file.

Column separator	Tabulator "\t"
Column 1	Date and time expressed according to ISO 8601
Column 2	Latitude in decimal format, unit degree
Column 3	Longitude in decimal format, unit degree

Processing Report:

This PDF document.



Cruise map




Figure 2: Map of the generalized master track