

# The two-faced Marginal Sea Ice Zone: A physical characterization and the link to biology

Giulia Castellani<sup>1</sup>, K.J. Müller<sup>2</sup>, F.L. Schaafsma<sup>3</sup>, S. Arndt<sup>1</sup>, B.A. Lange<sup>1,4</sup>, J. Ehrlich<sup>1,4</sup>, M. van Dorssen<sup>3</sup>, M. Nicolaus<sup>1</sup>, A. Meijboom<sup>3</sup>, A.P. Van de Putte<sup>5</sup>, M. Vortkamp<sup>1</sup>, B. Feij<sup>6</sup>, J.A. van Franeker<sup>3</sup>, H. Flores<sup>1,4</sup>

<sup>1</sup>Alfred Wegener Institute (AWI) Helmholtz-Zentrum für Polar- und Meeresforschung (Germany); <sup>2</sup>University of Oldenburg (Germany); <sup>3</sup>IMARES (The Netherlands); <sup>4</sup>University of Hamburg (Germany); <sup>5</sup>Royal Belgian Institute of Natural Sciences (Belgium); <sup>6</sup>NIOZ (The Netherlands);




@GEO Illuteam43

## INTRODUCTION


The marginal sea ice zone (MIZ) is a very dynamic and active area. In addition, the MIZ is biologically important due to an intense spring primary production bloom, which is an important carbon source for the marine food web. Here we present data collected in the Antarctic MIZ during a ship-based expedition to the Eastern Weddell Sea. The work aims to characterize the physical environment of the MIZ and identify the possible environmental drivers of ecological processes.

## 1. SAMPLING

Sea ice surveys were conducted with a Surface and Under Ice Trawl (SUIT). The SUIT is designed to sample under-ice fauna and it is equipped with a sensor array (Fig. 1) to measure the environmental properties: sea surface salinity, temperature, chl-*a*, under-ice spectral radiation, ice thickness, ice roughness and sea-ice algae content. In total, 6 profiles of ca. 2 km were collected in the MIZ in December 2014 (eastern transect, shown in Fig. 2) and January 2015 (western transect, shown in Fig. 3).




Sea ice concentration data were obtained from <http://www.meereisportal.de> (grant: REKLIM-2013-04)



## 2. SEA SURFACE PROPERTIES


Along-profile variability in sea surface properties (Fig. 4) is large. Figure 5 shows that the western stations have higher mean temperature, this agrees with the advanced melting stage as seen in the ice concentration (Fig. 3). Lower, but not significantly different, Chl-*a* concentration in surface sea water was registered in the western stations (Fig. 6).




## 4. SUIT CATCH



The variability found in sea surface and sea-ice properties is reflected in the species abundances and composition (Fig. 13 and 14).




13. Catch abundances for the dominant species *Euphausia Superba*



14. Catch abundances for other species

Species diversity also differs between transects. Krill *Euphausia superba*, *Thysanoessa macrura*, and the amphipod *Eusirus microps* were abundant in the western stations, the amphipod *Eusirus laticarpus* in the eastern ones.

## 3. SEA-ICE PROPERTIES



A clear separation between eastern and western stations is seen in all sea-ice properties. The ice is thicker (Fig. 7) in the western stations, and has more ridges (Fig. 8), detected as in Castellani et al. (2015). This is reflected in the light available under the ice (Fig. 9). Sea-ice algae (Fig. 10), retrieved as in Melbourne-Thomas et al. (2015), are more abundant in the eastern stations.

11. Sea-ice draft and sea-ice chl-*a* content for station 29\_1 (eastern transect). Shaded areas are ice covered regions

12. Sea-ice draft and sea-ice chl-*a* content for station 70-2 (western transect). Shaded areas are ice covered regions

Sea-ice algae show a large variability along profile (Fig.11 and 12). They are very low in regions where very thin ice is found. They are high in areas with thick ice and sometimes show peaks close to ridges, similar to what was observed by Lange et al. (2015) in hummocks.

## CONCLUSION

- Sea-ice physical properties in the Antarctic MIZ can vary drastically, showing two completely different environmental regimes
- The structure of under-ice communities clearly follows the strong contrast in sea-ice properties
- The general knowledge of the MIZ as a very productive area is, in half of the case studies presented here, not confirmed
- High abundances of Antarctic krill were associated with thick deformed ice and low chl-*a* in both sea ice and water, indicating that krill distribution under sea ice reflects other traits of the environment than food availability, such as predator avoidance
- Low Chl-*a* in surface water and sea ice could, on the other hand, be a consequence of grazing

## REFERENCES

- Castellani et al., (2015) Impact of sea-ice bottom topography on the Ekman pumping. Springer Earth System Science.
- Lange et al, (2015) Comparing springtime ice-algal chlorophyll *a* and physical properties of multi-year and first-year sea ice from the Lincoln Sea. PLoS One.
- Melbourne-Thomas et al. (2015) Algorithms to estimate Antarctic sea ice algal biomass from under-ice irradiance spectra at regional scales. Mar Ecol Prog Ser.

