



# Revealing Physical and Ecological Dynamics at an Ice Edge – a Robotic Approach

Wulff, T.<sup>1</sup>; Lehmenhecker, S.<sup>1</sup>; Hagemann, J.<sup>1</sup>; Busack, M.<sup>1</sup>; Tippenhauer, S.<sup>1</sup>; Strohmeier, M.<sup>2</sup>, Rothe, J.<sup>2</sup>









Polar Autonomous Underwater Laboratory















- --- Euphotic Depth
- AUV Dive Path

Polar Autonomous Underwater Laboratory







Polar Autonomous Underwater Laboratory








Chemical

Physical



Conductivity Pressure Temperature


Chlorophyll a (below vehicle)

Sample Collector

CDOM (below vehicle)















Polar Autonomous Underwater Laboratory









































# Thermal Image Sensor

- Goal: Automatic detection of ice shores
- Lepton FLIR
  - 80x60 Pixel
  - Approx. 5 Hz
  - Calibration range from -10°C to 65°C









#### Conclusion:

- AUVs are an ideal instrument to conduct high resolution water column studies
- The AUV is able to provide a comprehensive picture of the physical and ecological processes – but only on a small scale
- Flying drones can provide important environmental data, but they are not operational yet



## DGPS at high latitudes

- Carrier-phase based attitude determination relies on good GPS signal quality
- Low satellite elevation at high latitudes
   → poor SNR + high multipath
- Signal reflections in ship vicinity

