

## Arctic Ocean

# Litter Levels in the Depths of the Arctic are On the Rise

Sea ice could be responsible for transporting plastic litter

[08. February 2017] **The Arctic has a serious litter problem: in just ten years, the concentration of marine litter at a deep-sea station in the Arctic Ocean has risen 20-fold. This was recently reported in a study by researchers at the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research (AWI).**



Plastic bags, glass shards and fishing nets: despite its location, far from any urban areas, the amount of litter in the depths of the Arctic Ocean continues to rise, posing a serious threat to its fragile ecosystem. Since 2002, AWI researchers have been documenting the amount of litter at two stations of the AWI's "Hausgarten", a deep-sea observatory network, which comprises 21 stations in the Fram Strait, between Greenland and Svalbard. The results of the long-term study have now been published in the scientific journal *Deep-Sea Research I*. "Our time series confirms that litter levels in the Arctic deep sea have risen rapidly in the past few years," says first author and AWI biologist Mine Tekman.



Litter and sponges (Photo: OFOS/James Taylor)

The scientists involved in the study observed the ocean floor at a depth of 2,500 metres using the OFOS (Ocean Floor Observation System), a towed camera system. Since the start of their measurements, they have

spotted 89 pieces of litter in a total of 7,058 photographs. To enable comparison with other studies, the researchers have extrapolated the litter density to a larger area. The result: an average of 3,485 pieces of litter per square kilometre in the monitoring period (2002 to 2014). Further, there has been a clearly recognizable increase in the past few years: when the team calculated a contamination level of 4,959 pieces of litter per square kilometre for 2011 in an earlier study, they hoped it was a statistical outlier. But the

## Downloads



## Contact

### Science

 Melanie Bergmann  
 +49(471)4831-1739  


[Melanie.Bergmann@awi.de](mailto:Melanie.Bergmann@awi.de)

## Photos

[Public media](#)  
[Press media](#)

## Abo/Share




Subscribe to AWI press  
release RSS feed

[The Institute](#)

levels have continued to rise since, reaching a new peak of 6,333 pieces of litter per square kilometre in 2014.

The situation is particularly dramatic at the network's northern station, called N3. "Here the amount of litter rose more than 20-fold between 2004 and 2014," says Tekman. If we consider the findings for the northern research area in the marginal ice zone, the data for 2004 indicated 346 pieces of litter per square kilometre. Ten years later, the number had risen to 8,082. The level of contamination is similar to one of the highest litter densities ever reported from the deep seafloor, in Cap de Creus Canyon off the eastern coast of the Iberian Peninsula.

Among the litter they photographed, the researchers observed plastic and glass most frequently. As a rule, glass does not drift; it sinks straight down to the ocean floor. This indicates local sources and concurs with increasing ship traffic in the region



AWI HAUSGARTEN (Photo: Alfred Wegener Institut)

due to the receding ice. Still, it is extremely difficult to draw any firm conclusions on the origin of the plastic litter, since it often covers a considerable distance before reaching the seafloor. In most cases, the scientists can't determine the origin on the basis of photographs alone. While it is clear that the Gulf Stream transports plastic litter into the Arctic with the Atlantic water masses, the authors also have a new theory as to why litter is accumulating in the Fram Strait: their results indicate a positive relationship between litter density and the summertime expansion of sea ice. "If we're right, sea ice could entrain floating litter during ice formation. During warmer periods, the ice breaks up and is transported to the south into the Fram Strait with the Transpolar Drift, releasing entrained litter into the survey area when it melts," says deep-sea biologist Dr Melanie Bergmann, a co-author of the study. "To date we've assumed just the opposite, since we viewed the ice as a barrier to litter contamination."

The researchers are still faced with the puzzle of when and how plastic litter changes on its way to the deep sea. Over time they have observed more and more small bits of plastic, which are likely the result of larger pieces of litter fragmenting and could point to an increasing level of microplastic. This is surprising, since in the deep sea there is no UV light to break down the plastic, and the low temperatures are not conducive to disintegration. In the summer of 2016 the team rediscovered a piece of plastic, which they had first seen two years earlier. Bergmann: "Running into this same piece of plastic twice with hardly any changes to it is a vivid reminder that the depths of the Arctic are at risk of becoming a depot for plastic litter. The well-hidden accumulation of litter on the deep ocean floor could also explain why we still don't know where 99% of the marine plastic litter ends up."

#### More information on the AWI Hausgarten and the OFOS camera system:

The Hausgarten is the AWI's deep-sea observatory in the Fram Strait, and currently consists of 21 stations at depths ranging from 250 to 5,500 metres. Samples have been collected at these stations every summer since 1999. Further, the year-round operation of moorings and free-falling lander systems, which serve as observation platforms on the seafloor, makes it possible to record seasonal changes. With the help of an ROV (Remotely Operated Vehicle), samples



The Alfred Wegener Institute pursues research in the polar

regions and the oceans of mid and high latitudes. As one of the 18 centres of the Helmholtz Association it coordinates polar research in Germany and provides ships like the research icebreaker Polarstern and stations for the international scientific community.

## More information

### Topic pages

» [Marine Litter](#)

### Related news

» [Spotlight on marine litter: A new book presents the current state of research](#)  
» [Are there no European waters free of litter? A new study shows that all of Europe's deep seas investigated are polluted with litter](#)

are collected at regular intervals, automatic recording instruments are positioned and maintained, and in-situ experiments are conducted. The Hausgarten represents one of the key regions in the European Network of Excellence ESONET (European Seas Observatory Network) and is part of Germany's Long Term Ecological Research network (LTER-D).

During Polarstern expeditions to the Hausgarten, deep-sea researchers at the Alfred Wegener Institute regularly employ the remotely controlled towed camera system OFOS (Ocean Floor Observation System). At four Hausgarten stations its cameras are suspended roughly 1.5 metres above the seafloor at a depth of 2,500 metres, where they take a photograph every 30 seconds. These photographs enable deep-sea biologists to document changes in the species diversity for larger forms of sea-life, for example in sea cucumbers, sea lilies, sponges, fish and shrimps.

## Original publication

Mine B. Tekman, Thomas Krumpen, Melanie Bergmann: *Marine litter on deep Arctic seafloor continues to increase and spreads to the North at the HAUSGARTEN observatory*. Februar 2017. DOI: [10.1016/j.dsr.2016.12.011](https://doi.org/10.1016/j.dsr.2016.12.011)

