

Bonamia infection in native oysters (*Ostrea edulis*) in relation to European restoration projects

Hein Sas¹ | Brenda Deden² | Pauline Kamermans³ |
Philine S.E. zu Ermgassen⁴ | Bernadette Pogoda⁵ | Joanne Preston⁶ |
Luke Helmer⁶ | Zoë Holbrook⁷ | Isabelle Arzul⁸ | Tom van der Have⁹ |
Antonio Villalba¹⁰ | Bérenger Colsool⁵ | Alice Lown¹¹ | Verena Merk⁵ |
Nadescha Zwerschke¹² | Emilie Reuchlin²

¹Sas Consultancy, Amsterdam, Netherlands

²Oceans Unit, WWF Netherlands, Zeist, Netherlands

³Wageningen Marine Research, Wageningen University and Research, Yerseke, Netherlands

⁴Changing Oceans Group, School of Geosciences, University of Edinburgh, Edinburgh, UK

⁵Helmholtz Centre for Polar and Marine Research, Alfred Wegener Institute, Bremerhaven, Germany

⁶Institute of Marine Sciences, University of Portsmouth, Portsmouth, UK

⁷Ocean and Earth Science, University of Southampton, Southampton, UK

⁸SG2M-LGPMM, Laboratoire De Génétique Et Pathologie Des Mollusques Marins, Ifremer, La Tremblade, France

⁹Team Aquatic Ecology, Bureau Waardenburg, Culemborg, Netherlands

¹⁰Centro de Investigacións Mariñas, Consellería do Mar da Xunta de Galicia, Vilanova de Arousa, Spain

¹¹School of Life Sciences, University of Essex, Colchester, UK

¹²Marine Laboratory, Queen's University Belfast, Portaferry, Ireland

Correspondence

Hein Sas, Sas Consultancy, Danie Theronstraat 22d, Amsterdam, 1091XZ, Netherlands.
Email: hsas@xs4all.nl

Abstract

1. There is a growing effort throughout Europe to restore populations of native oysters (*Ostrea edulis*), with the ecological objective of enhancing ecosystem biodiversity and resilience.
2. The introduced parasite, *Bonamia ostreae*, caused catastrophic mortalities during the 1980s, furthering the decline of this species, and is now present throughout much of the natural range of *O. edulis*. It is therefore important that restoration attempts avoid further introduction and spread of this parasite, which can cause lethal infections of *O. edulis*.
3. This article presents a comprehensive overview of the scale and distribution of current infection, transmission pathways, and preventive measure guidelines, focusing on the seas, inlets, and estuaries of north-west Europe, where most ecological restoration attempts for the native European oyster have occurred so far.
4. This is critical information for restoration project planning in which the risk of *Bonamia* infection must be taken into account.

KEYWORDS

coastal, disease, invertebrates, restoration, subtidal

1 | INTRODUCTION

The European native (or 'flat') oyster (*Ostrea edulis*) was once abundant throughout many coastal European waters and offshore areas

of the North Sea (Figure 1), where it was found in dense aggregations (Möbius, 1877). However, *O. edulis* suffered substantial declines throughout the 19th and 20th centuries. It is now extirpated from much of its range (Beck et al., 2011) and is listed as a threatened and declining habitat by OSPAR (OSPAR Commission, 2009). There is now a growing effort throughout

Hein Sas and Brenda Deden are joint lead author.




FIGURE 1 Nineteenth century occurrence of *Ostrea edulis* in Olsen's Piscatorial Atlas of the North Sea (Olsen, 1883)

Europe to restore populations of this habitat building species, with the aim of enhancing biodiversity and ecosystem resilience (Pogoda, Brown, Hancock, & von Nordheim, 2017).

While the initial collapse of *O. edulis* populations was largely driven by overfishing (Gercken & Schmidt, 2014; Houziaux, Kerckhof, Degrendele, Roche, & Norro, 2008; Pogoda, 2019), the emergence of parasites such as *Bonamia ostreae* and *Marteilia refringens* during the 20th century resulted in substantial mortalities, furthering a renewed widespread decline of *O. edulis* (Laing, Walker, & Areal, 2006), in particular in aquaculture of this species along European coasts. These parasites are still present in several European ecoregions, with varying virulence and impact. *B. ostreae* is especially widespread in the seas and inlets of north-west Europe, posing a threat to the success of oyster restoration projects. Biosecurity relating to *B. ostreae* transmission and spread is therefore an essential consideration when planning and implementing restoration of *O. edulis*.

Bonamiosis is an oyster disease that is generally caused by parasites of the genus *Bonamia*. *Bonamia* infects immune system cells (haemocytes) of the genus *Ostrea*. *B. ostreae* is the parasite that causes the severest *O. edulis* disease in European waters; hence, it is the main subject of this article. It has been the focus of substantial research within aquaculture settings (e.g. Arzul et al., 2009, 2011; Bougrier, Tigé, Bachère, & Grizel, 1986), but the specific impact of the disease on attempts to restore high densities of *O. edulis* on the seafloor and the

appropriate management to use in this setting remains a knowledge gap. Current oyster restoration projects in Europe are seeking to increase the density and extent of *O. edulis* to levels at which the species can be considered a self-sustaining population. Since parasite prevalence probably increases with density (Engelsma et al., 2010), the risk of disease incidence may increase through restoration attempts. This should obviously be avoided.

Because of this, it is important that restoration efforts comprehensively consider the risk posed by *B. ostreae* and avoid its further spreading (Pogoda et al., 2019). This is strongly encouraged by NORA, the Native Oyster Restoration Alliance (for Europe). To avoid the risk of spreading *B. ostreae* in restoration activities, it is important to consult the best available and most up to date knowledge on *B. ostreae*. The current review presents a comprehensive overview of the current *B. ostreae* infection distribution in north-west Europe, transmission pathways and preventive measures against the disease, leading to recommendations for restoration project practices.

Many restoration projects in north-west Europe are currently being undertaken, as shown in Figure 2. There are numerous other sites where *O. edulis* are managed for aquaculture and food production, but for Figure 2, only *O. edulis* restoration projects which are being undertaken to improve biodiversity and habitat quality are selected.

2 | METHODS

The urgent need to summarize the existing information regarding the *Bonamia* infection, its potential impacts, and management strategies for *O. edulis* restoration in Europe was recognized within the NORA community with the initiation of the first European *O. edulis* restoration projects. An initial review of the existing scientific, peer-reviewed literature on the disease was presented at the 1st NORA conference, of 1–3 November 2017 in Berlin. The article was extended and refined on the basis of discussions during the conference and a second draft was presented and discussed at the 2nd NORA Conference of 21–23 May 2019 in Edinburgh. In addition, experts on specific topics were involved, resulting in the current author collective.

The basic data on geographical distribution of the *Bonamia* infection was obtained through a survey of the relevant literature and public animal disease databases, such as (WAHIS, 2020). There is a delay time between detection of the disease and publication in these sources, so that the NORA community was consulted to obtain the most up-to-date information (until January 2020).

Since various terms are adopted in the literature to indicate the disease status, potentially leading to confusion, the terminology in this article is here defined as:

- Oysters demonstrated to be infected are referred to as 'Bonamia-infected'
- Oysters originating from a region where *B. ostreae* is present, are referred to as 'Bonamia-exposed'.




FIGURE 2 Impression of current *Ostrea edulis* restoration attempts in north-west Europe (NORA, 2019). Green star denotes restoration project. See Table S1 for the corresponding restoration project information

- Oysters originating from a (also historically) *Bonamia*-free region or demonstrated to be free of the infection by adequate testing, are called 'Bonamia-free'.
- In the theoretical case that an oyster without infection is produced from a *Bonamia*-exposed population, these are called 'Bonamia-negative'.

The first three terms are also adopted to indicate the infection status of oyster growing areas.

3 | WHAT IS BONAMIOSIS AND WHICH SPECIES DOES IT AFFECT?

Bonamiosis is a disease caused by unicellular parasites of the genus *Bonamia* (Arzul & Carnegie, 2015; Culloty & Mulcahy, 2007), included in the protozoan group Haplosporida, within Ascetosporia (Bass, Ward, & Burki, 2019). Three *Bonamia* species have been characterized: *B. ostreae* (Pichot, Comps, Tige, Grizel, & Rabouin, 1980), *Bonamia exitiosa* (Berthe & Hine, 2003; Hine, Cochennec-Laureau, & Berthe, 2001), and *Bonamia perspora* (Carnegie et al., 2006). The parasite named *Bonamia*

roughleyi (Farley, Wolf, & Elston, 1988) was erroneously attributed to the genus *Bonamia* (Carnegie, Hill, Stokes, & Burreson, 2014).

The host range of *B. ostreae* and *B. exitiosa* includes multiple species of the genus *Ostrea*. Besides *O. edulis*, oyster species that are documented to be infected with *Bonamia* spp. are of the genera *Crassostrea*, *Saccostrea*, and *Dendostrea*, but with less severe consequences for the remaining populations (Laramore, Krebs, Lave, & Gallagher, 2017).

In Europe, *Bonamia exitiosa* infects *O. edulis* in Galicia (Abollo et al., 2008; Ramilo et al., 2014), and it has been detected in Catalonia (Carrasco et al., 2012), Italy (Narcisi et al., 2010), France, UK (Longshaw, Stone, Wood, Green, & White, 2013) and Portugal (Batista, López-Sanmartín, Grade, Navas, & Ruano, 2016). *B. perspora* is considered of less relevance in Europe, since it has as yet only been reported in *Ostrea stentina* in North Carolina, USA (Carnegie et al., 2006).

Several other oyster diseases such as marteiliosis, due to *Marteilia refringens*, should also be considered within the framework of oyster restoration projects, but this article focuses on *B. ostreae*, since this is currently considered to pose the most serious disease threat to *O. edulis* in north-west Europe.

4 | B. OSTREAE IN NORTH-WEST EUROPE

4.1 | Introduction of *B. ostreae* in Europe and its consequences

In Europe, oyster production and fishing activity was extensive during the 19th century. After severe declines in oyster stocks, related to unsustainable fishing pressure, large scale oyster translocations were undertaken in order to revive depleted populations. At the beginning of the 20th century, the industry started to suffer from its first disease driven mortalities.

The oldest epizootic affecting flat oysters and related in the literature took place in France, The Netherlands, and UK from 1920 to 1927 (Grizel, 1985; Héral, 1990). Although no infectious organism had really been incriminated, Orton (1924a, 1924b) described several abnormal cellular figures looking like intracellular parasites. During this period, the production drastically declined. The disease was retrospectively identified as probably caused by the flagellate protozoan *Hexamita* and associated with high laying densities as found in the managed beds (Tubbs, 1999).

In 1930s and 1940s, shell oyster disease, caused by the fungus *Ostracoblabe implexa* (Alderman, 1985; Alderman & Jones, 1970), caused severe losses to the Dutch oyster industry and, to a lesser extent, the French industry. This disease was overcome by changing some common practices in the culture procedures (Korringa, 1951; Korringa, 1976).

In 1968 in Aber Wrac'h, an inlet on the north-west coast of Brittany (France), the parasite *M. refringens* was diagnosed in oysters (Culloty & Mulcahy, 2007; Grizel et al., 1974), causing large-scale mortalities in *O. edulis*. In 1979, a second parasite – *B. ostreae* – was discovered in L'Ile Tudy, on the south-west coast of Brittany (Pichot et al., 1980), probably originating from the coastal waters of California (Elston, Kent, & Wilkinson, 1987). This infection caused additional large-scale mortality and spread rapidly following its introduction, primarily due to the movements of infected oysters to new grow-out areas, or by careless movements of infected oysters with other shellfish (Culloty & Mulcahy, 2007).

In France, a 93% reduction in yield was recorded between early 1970 and 1982 due to bonamiosis (Laing et al., 2006). Overall, European production of *O. edulis* fell from 29,595 tons in 1961 to 5,921 tons in 2000 (Culloty & Mulcahy, 2007). The impact of the diseases caused by *M. refringens* and *B. ostreae* resulted in a shift to the rearing of *Crassostrea gigas*, and for *O. edulis* production to remain low throughout the 1990s and beginning of the 21st century (Culloty & Mulcahy, 2007; Haenen, Engelsma, & Beurden, 2011).

Being a serious oyster disease, bonamiosis is notifiable to the World Organization for Animal Health (OIE, 2019) and it is included in the list of non-exotic diseases entailed in the EU Council Directive regulating aquatic animal health issues (EU, 2006). Movement of oysters from infected areas to infection-free areas poses an unacceptable biosecurity risk, yet the limited sources of *Bonamia*-free *O. edulis* spat or adults from historically *Bonamia*-free areas to be used as restoration broodstock pose a challenge to restoration efforts. Understanding

the historical spread, present infection status and current knowledge of immunological responses to this infection is imperative for sustainable restoration efforts.

4.2 | Current spread of *B. ostreae* in north-west Europe

The majority of *O. edulis* populations in Europe are now infected by *B. ostreae* (Figure 3). The database underlying Figure 3, with location names, source, and years of first recorded *B. ostreae* presence (if available) is presented in Table S2. The ultimate data underlying the map and the table are from January 2020.

B. ostreae is thought to have first spread through oyster cultures in France (Elston et al., 1987) and Spain (Cigarria & Elston, 1997) before reaching other European coastal waters within a decade (Culloty & Mulcahy, 2007). Bonamiosis reached the UK in 1982 and Ireland in 1987 (Culloty & Mulcahy, 2007). Some bays and inlets in the UK and Ireland, however, have thus far remained *Bonamia*-free (Laing, Dunn, Peeler, Feist, & Longshaw, 2014).

In Norway, oyster cultures have been regularly surveyed since 2008 and an infection detection was reported for 2009 in the Langestrand area (OIE, 2009). However, the parasite has not been detected since at this location during examinations carried out by the National Veterinary Institute (Mortensen, Sælemyr, Skår, Bodvin, & Jelmert, 2016; Mortensen, Sælemyr, Skår, Bodvin, & Jelmert, 2018). Hence, the status reported in Figure 3 is 'uncertain' for the Langestrand location. Repeated surveys at other Norwegian locations showed no *Bonamia*-infection (Mortensen et al., 2018), so these are reported as *Bonamia*-free in Figure 3.

In Denmark, the main oyster culture area is Limfjord, which remained *Bonamia*-free for a long time (Møllergaard, 2008). *B. ostreae* was recently reported at very low prevalence in the Nissum Bredning, in the western part of Limfjorden (ICES, 2018; Madsen, 2017), which means that Limfjord is now considered a *Bonamia*-infected area, regardless of the fact that there has been no increased mortality (Madsen, 2017).

The *Bonamia* status in Dutch waters is generally 'infected'. However, a small *O. edulis* population was recently discovered in the Dutch Wadden Sea, which was tested by performing DNA analysis on a large number of larvae produced in a hatchery. These were reported free from *Bonamia* (Jacobs et al., to be submitted). However, since no adult oysters were tested, the *Bonamia* status of the area has to be considered as 'uncertain'. Open-sea areas in Dutch waters marked as *Bonamia*-free in Figure 3 represent isolated restoration projects, for which *Bonamia*-free oysters (from Norway) have been employed. In an early restoration (2017) restoration project off the west coast of The Netherlands *Bonamia*-infected oysters were deployed. These oysters could not be retraced, but the location is marked as 'infected' nonetheless.

In December 2019 it was discovered that the Lynn of Lorne, Loch Creran, Loch Etive, and Dornoch Firth oyster populations in Scotland are infected by *Bonamia* (Scottish Government, 2020).



FIGURE 3 Occurrence of *Bonamia ostreae* infection in north-west Europe. The colour of the marked points indicates the infection status of the present oyster population as revealed by our survey: red for 'Bonamia-infected', blue for 'Bonamia-free' and yellow for 'uncertain status'. • When a location is marked as infected, this means that one or more oysters from this area have been tested *B. ostreae* positive. • When a location is marked as *Bonamia*-free, this means that no *B. ostreae* has been detected with regular surveys and tests to date. • When a location is marked as uncertain, this means sources that on the *Bonamia* status are contradictory, not present, or unknown. For details on prevalence of *Bonamia* in marked locations, see Table S2

5 | CHARACTERISTICS OF *B. OSTREA* AND BONAMIOSIS AND THE RELEVANCE FOR RESTORATION PRACTICE

5.1 | Infection and disease development in oysters

Once present, the *Bonamia* parasite spreads rapidly through *O. edulis* beds (Culloty et al., 1999). Although pathways of infection are not fully known, *O. edulis* is susceptible to infection by *B. ostreae* at all life-history stages, including during larval phases (Arzul et al., 2011; Lynch, Armitage, Wylde, Mulcahy, & Culloty, 2005). Male and female oysters are equally susceptible to infection (Culloty & Mulcahy, 1996). An

initial 'latent' period can mask the infection from detection for anything from 4 weeks to several months (Culloty, Cronin, & Mulcahy, 2001).

B. ostreae is an intracellular parasite (2–5 µm) that infects the haemocytes and, occasionally, branchial epithelium (ectoderm) of the oysters (Arzul & Carnegie, 2015). Haemocytes are suspended in the haemolymph fluid, which is a plasma similar to the blood in vertebrates. One of the functions of haemocytes is to detect and destroy pathogens, but *O. edulis* haemocytes fail to destroy *Bonamia*. There is evidence that the parasite inhibits or blocks molecular weapons of oyster haemocytes to destroy pathogens (Gervais et al., 2019; Gervais, Chollet, Renault, & Arzul, 2016; Gervais, Renault, & Arzul, 2018; Hervio, Chagot, Godin, Grizel, & Mialhe, 1991).

The infection usually develops through infiltration of infected haemocytes into the tissues of the gills and mantle and around the gut. In its severe state, it causes loss of the normal architecture of the gills, the digestive gland, the gonad, and other organs leading to general dysfunction and ultimately death of the oyster (Culloty & Mulcahy, 2007). Bonamiosis usually causes highest mortality in oysters that are 3 years or older, although younger infected oysters may also suffer mortality (Lynch et al., 2005). Sometimes, the effect of the disease is sublethal, reducing the host's ability to cope with additional stressors such as changes in water temperature, translocation to other environments, or reproductive activity (Dijkema, 1990; van Banning, 1991) and increasing host susceptibility to other microorganisms. Eradication of, or treatment against, *B. ostreae* is not considered possible (Morga et al., 2017).

The parasite occurs throughout the year, but prevalence of infection tends to be highest in spring and summer, with the peak of prevalence at the end of winter to spring in most of the infected countries in Europe (Culloty & Mulcahy, 1996; Engelsma et al., 2010).

5.2 | Detection methods of infection by *B. ostreae* at individual and population levels

Detection of *Bonamia* presence in the source *O. edulis* population for restoration purposes is essential to avoid accidental spreading of the infection. Testing should also be performed if an *O. edulis* population is already present in the restoration area in order to determine any previous presence of the parasite.

B. ostreae infection is often difficult to detect visually in the oyster, but gross signs can occasionally be observed including yellow discoloration in the gills and extensive lesions, including perforated ulcers in the connective tissues of the gills, mantle, and digestive gland. Standard diagnostic methods use cytology (haemolymph smears or tissue imprints) and histopathology to screen oyster tissues, after staining the sample (da Silva & Villalba, 2004).

DNA techniques, based on the polymerase chain reaction, are now widely used, due to their high specificity and ability to detect very low infection levels (Flannery et al., 2014). New species-specific molecular methods are available (Ramilo, Navas, Villalba, & Abollo, 2013) and their use is recommended in European regulation (EU, 2015). These species-specific tools (Batista et al., 2016; Flannery, Lynch, Longshaw, et al., 2014; Ramilo et al., 2013) confer high sensitivity and can detect a lower degree of infection/presence than histological analysis. However, it may also yield false positive detections. The lower sensitivity of more dated primers that are currently recommended by the World Organization for Animal Health (OIE, 2019) may provide underestimations of prevalence within a population (Helmer et al., 2020).

Thus, compared to histology, DNA-techniques appear to be more sensitive. However, they are indicative of the presence of *B. ostreae* DNA and not of an infection: histology remains a key technique to

confirm an infection especially in a previously *Bonamia*-free population or region.

Even if the prevalence of the infection in a population is low, it is crucial to be able to detect it. An important factor in determining whether a population can confidently be assessed for its *Bonamia* infection status is sample size. The EU prescribes a minimum sample size of 150 individual oysters in Annex I, part 5 of EU (2015). The document does not explain the requirements and assumptions underlying this number, but by using basic statistics as provided by the World Organization for Animal Health (OIE, 2008) these can be reconstructed as:

- required confidence level: 95%;
- *Bonamia* prevalence in the *O. edulis* population to be tested: 2%;
- sensitivity of the testing method: 95%.

More extensive recommendations for *Bonamia* survey and detection methods are given in OIE (2019).

5.3 | Spreading mechanisms of the *Bonamia* infection

Transmission pathways of *B. ostreae* may occur directly from parent oysters to larvae, but also via the water column, probably via filtration (Arzul et al., 2011; Culloty & Mulcahy, 2007). The mechanism of transmission is not fully understood, although some mechanisms and factors are described by Engelsma, Culloty, Lynch, Arzul, and Carnegie (2014).

The maximum transmission distance is also unknown. It could be relatively small, since infection prevalence tends to increase with oyster population density (Engelsma et al., 2010) and *Bonamia*-free and -infected *O. edulis* areas are observed to exist at a close distance to each other, e.g. in bays and inlets in south-west England (Figure 3). However, since the infection can be transferred through water currents and also larvae (which remain in the water phase for 11–30 days) potentially large dispersal distances (10 km or more) can occur, depending on the local hydrogeographic regime. The main infection vector is, however, considered to be shellfish transfers of infected *O. edulis*. Hence, EU regulation against the spreading of the infection focuses on quarantining infected areas where transport of *O. edulis* from infected to non-infected areas is prohibited (EU, 2006).

Given that the infection can be transmitted through larvae and the water phase, once present on an oyster bed, *B. ostreae* cannot be eradicated (van Banning, 1991). *O. edulis* are not the only shellfish species to transmit *B. ostreae* (Engelsma et al., 2014; Laramore et al., 2017). Contrary to initial evidence, which suggested that *C. gigas* was not susceptible to infection (Culloty et al., 1999), it is now believed that it may indeed act as either a paratenic or dead-end host for both *B. ostreae* and *B. exitiosa* (Helmer et al., 2020; Lynch et al., 2010). This should be investigated further as infection and

transmission via this highly abundant and commercially produced species could have implications for restoration of *O. edulis* and the transport of commercial stock could exacerbate the spread of *Bonamia* species. Non-bivalve species may also serve as vectors, such as the brittlestar *Ophiothrix fragilis* (Lynch, Armitage, Coughlan, Mulcahy, & Culloty, 2007).

5.4 | Sensitivity of the *Bonamia* parasite to climate change

General effects of climate change include higher temperature and (through dissolution of CO₂) lower pH (Huthnance et al., 2016). To our knowledge, there are no specific studies of the impact on the prevalence and/or the mortality caused by *Bonamia* infection to *O. edulis* populations under these climate change scenarios. Arzul et al. (2009) have, however, studied the survival of purified *Bonamia* parasites using sea water from three different sources with pH values of 8.06, 7.06 and 6.5 under different temperature regimes. The results showed significantly lower survival at 25°C compared to 4°C and 15°C. Regarding pH, an *ad hoc* experiment was not performed because sea water with different pH values also differed in chemical composition, but the results showed a better survival of purified *B. ostreae* (60–80%) in the sea water with pH = 8.06 and pH = 7.06 than in artificial sea water (survival <40%) with pH = 6.5, regardless of temperature and incubation time. It is worth noting that *B. ostreae* exhibited high survival under the full range of pH and temperature conditions tested. Besides, the tested range of temperature and pH is far greater than the actual changes in the variables predicted in Huthnance et al. (2016) for the end of this century, so it seems unlikely that the *Bonamia* parasite will be strongly negatively impacted by climate change in the near to medium term. However, specific research on the interactions between climate change effects and *Bonamia* is needed to test this hypothesis.

5.5 | Evidence for tolerance or resistance in existing *O. edulis* populations

Disease tolerance and resistance are two physiological defence strategies demonstrated by *O. edulis* in response to infection by the parasite *B. ostreae*. Disease resistance is when the parasite is able to infect the host, but it is unable to multiply, reproduce and therefore to proliferate within the host tissues. Resistant individuals have also demonstrated the ability to reduce parasite burden (Ayres & Schneider, 2008; Morga et al., 2017; Råberg, Sim, & Read, 2007). Disease tolerance is when the host's fitness is not greatly affected by the presence of the parasite, regardless of its successful proliferation in host tissues (Ayres & Schneider, 2008; Råberg, Graham, & Read, 2008). This balance between parasite and tolerant host can be interrupted by stress, as any environmental pressure such as a change in abiotic conditions or food supply can lead to immune imbalance, resulting in host mortality (Mydlarz, Jones, & Harvell, 2006).

Although marine invertebrates lack the ability to develop pathogen specific antibodies, *O. edulis* from *Bonamia*-exposed populations have demonstrated more resistance or tolerance to the parasite than oysters from *Bonamia*-free populations (Culloty et al., 2001; Culloty, Cronin, & Mulcahy, 2004; da Silva, Fuentes, & Villalba, 2005; Hervio et al., 1995). (Morga et al., 2017) demonstrated a degree of disease resistance in *Bonamia*-exposed oysters, with inhibiting phagocytotic activity to reduce the spread of parasites to further tissue, while inducing in haemocytes the expression of genes associated with apoptosis, thus hampering parasite proliferation within haemocytes.

Various studies in different countries have shown that oysters living in areas affected by bonamiosis for a long time (i.e. >20 years) survive exposure to *B. ostreae* much better than oysters living in areas only recently affected by the disease or in non-affected areas, indicating development of natural resistance or tolerance of oysters to infection by the parasite over time (da Silva et al., 2005; Flannery, Lynch, Carlsson, Cross, & Culloty, 2014).

Selective breeding for resistance or tolerance has taken place in Cork Harbour, Ireland (Lynch, Flannery, Hugh-Jones, Hugh-Jones, & Culloty, 2014). This has taken the form of large-scale breeding trials in spatting ponds, using 4–5-year-old survivors of the disease. In laboratory- and field-based trials comparing the susceptibility of the Cork Harbour *O. edulis* with Irish and European populations, the former have performed well (Culloty et al., 2001; Culloty et al., 2004). Again, the mechanism through which this occurs is unknown. Additionally, pilot programmes have been performed in France (Baud, Gerard, & Naciri-Graven, 1997; Naciri-Graven, Haure, Gérard, & Baud, 1999; Naciri-Graven, Martin, Baud, Renault, & Gérard, 1998) and Spain (da Silva et al., 2005), also showing that selective breeding leads to significant increase of tolerance or resistance and survival.

Culloty et al. (2004) compared performance of oysters that had been selectively bred for resistance to *B. ostreae* (Rossmore, Cork Harbour, Ireland), and oysters from two areas where *Bonamia* has been present for a long time (Lake Grevelingen, the Netherlands; Brittany, France) with oysters from four *Bonamia*-free populations (Lough Foyle, Ireland; Tralee, Ireland; Loch Kishorn, Scotland; Mull, Scotland). Oysters from all these locations were translocated to Cork Harbour (Ireland), Lake Grevelingen (the Netherlands), and Brittany (France). The field trials indicated that Rossmore and Lake Grevelingen oysters showed lower mortality compared to other populations. Culloty et al. (2004) concluded that previous exposure in these populations had conferred some reduced susceptibility to the parasite compared to *Bonamia*-free populations. In a follow-up study, spat was produced in the hatchery of Roem van Yerseke with broodstock from long-term exposed populations in Lake Grevelingen and the Oosterschelde, and a *Bonamia*-free population in Limfjord in Denmark. Spat of all three groups were reared for 1 year in Lake Grevelingen. Survival was best in spat from Lake Grevelingen (OYSTERECOVER). It was concluded that Grevelingen should be considered as a candidate stock for starting a breeding programme in the Netherlands. Although this stock had the highest overall prevalence of infection, it also had the greatest growth and survival rate indicating that it may have formed some local

tolerance to the disease. Appropriate design to avoid undesirable side-effects of inbreeding or substantial reduction of the genetic variability of the species should be considered when selecting oysters for resistance or tolerance.

The development of *B. ostreae* resistance and/or tolerance is a hopeful sign. Efforts to understand how oysters become resistant to *B. ostreae* have increased in recent years; studying gene expression associated with *B. ostreae* infection (Gervais et al., 2016; Gervais et al., 2018; Gervais et al., 2019; Morga et al., 2017; Morga, Renault, Faury, Chollet, & Arzul, 2011) and comparing it between *O. edulis* stocks with different susceptibility to the parasite (Morga et al., 2017; Pardo et al., 2016) are providing clues. Decreasing phagocytic activity and increasing apoptosis (i.e. cell suicide) of haemocytes seem to be associated with increased oyster resistance (Gervais et al., 2016; Gervais et al., 2019; Morga et al., 2017), probably by restraining parasite multiplication within haemocytes.

Genetic analysis has so far identified multiple genes indicating bonamiosis immunity, including OelAP and OeFas-ligand gene expression, highlighting differences in wild-type and selectively bred oysters in their ability to regulate apoptosis (Morga et al., 2017). Comparisons of gene expression profiles in *Bonamia*-free and -infected oysters are producing suites of candidate resistance conferring genes (e.g. Ronza et al., 2018; Vera et al., 2019) for testing and screening resistance. Proteomic approaches can also contribute to identify molecular markers of resistance to bonamiosis (de la Ballina, Villalba, & Cao, 2018).

Given the importance of promoting resistance and/or tolerance on the one hand, and the absolute need to avoid the spread of *Bonamia* on the other, this is a critical, although challenging, area of research.

5.6 | Biosecurity measures

As the transfer of stocks of *O. edulis* is considered to be responsible for the introduction of bonamiosis in Europe (Bromley, McGonigle, Ashton, & Roberts, 2016), biosecurity measures rely on the prohibition of transfer of live or dead oysters, of any age class, from an infected area. This is mandatory under current EU regulations (EU, 2006). In accordance to this regulation, all oyster transports are subject to licensing, according to EU and/or national regulation. The project organizer should therefore always apply for a transport licence (and other relevant licences) from the competent authorities in the country where the restoration project is undertaken and adhere to licence conditions at all time.

Upon transfer of oysters to sensitive locations, such as the restoration project area and hatcheries, measures have to be put in place to limit spreading of the disease as much as possible. These should include the quarantine of oysters, combined with analysis for the detection of *B. ostreae* on a sample of the oysters, applying the techniques described above. Most techniques lead to the destruction of the sample, but a non-destructive method (analysing samples of tissue collected from previously anaesthetized oysters [Kamermans et al., submitted]) is being developed.

5.7 | Production of oysters which are simultaneously *Bonamia*-free and *Bonamia*-tolerant/resistant

Production of oysters that are simultaneously *Bonamia*-free and *Bonamia*-tolerant should be technically feasible. Infection of a population by *Bonamia* does not result in the total eradication of that population. Within the remaining population, there will always be uninfected as well as infected individuals. Following long-term exposure to the parasite, these uninfected individuals can be identified within the population, and spat derived from them in a hatchery can be non-infected. *Bonamia*-infection in this new generation can be reliably detected with polymerase chain reaction/DNA analysis, given the correct minimum amount of spat tested. Hence, a *Bonamia*-free broodstock can be established in a hatchery and, if managed properly (with quarantine measures), non-infected spat ready to be relayed can be produced from these. These oysters may have developed tolerance or resistance to the disease (Kamermans et al., submitted).

This is potentially very useful for restoration projects, since international regulations and national policies aim to prevent the transfer of diseases to new areas, but protection against disease is desired, in case it does appear in a newly established bed. Recently, the first step in this process has been taken. A novel, non-destructive screening method to determine the status of the oyster with regard to *Bonamia* was developed and the selected *Bonamia*-free broodstock produced *Bonamia*-free spat (Kamermans et al., submitted). Further analysis into the genetic profile of these spat is underway to identify any genes that can be used as markers for resistance.

5.8 | Maintaining genetic diversity

Genetic differentiation exists between Atlantic, Mediterranean, and Black Sea native oyster populations (Diaz-Almela, Boudry, Launey, Bonhomme, & Lapegue, 2004; Launey, Ledu, Boudry, Bonhomme, & Naciri-Graven, 2002; Sobolewska & Beaumont, 2005). Native oysters have been cultivated since Roman times, and translocations, especially during the 1800s, were most intense between various north-east Atlantic populations, with translocations taking place to a lesser extent between north-east Atlantic and Mediterranean populations (Bromley et al., 2016). This can explain the moderate genetic differentiation between Atlantic and Mediterranean *O. edulis* populations and a tendency for Atlantic populations to be even less differentiated than Mediterranean ones (Launey et al., 2002). However, Vera et al. (2016) studied oyster populations in the Netherlands, Denmark, Ireland, England, France, and Spain and revealed systematic genetic differences between native oysters in three geographical regions: (1) The Netherlands and Denmark; (2) France, Ireland, and England; and (3) Spain. In addition, Gutierrez et al. (2017) showed high genetic similarity in *O. edulis* between Norway, Lake Grevelingen, and Maine.

The selection of resistant oysters involves reproduction with *Bonamia*-free broodstock in a hatchery, but spat produced in a hatchery has a lower genetic diversity than pond production or

spat collection in the field (Lallias, Boudry, Lapegue, Kin, & Beaumont, 2010). Thus, it is important to maintain genetic diversity in hatchery production through regular replacement of broodstock oysters, with new individuals from outside waters (Ryman & Laikre, 1991).

6 | RECOMMENDATIONS FOR RESTORATION PRACTICE

6.1 | Avoidance of spreading diseases in general

Since *O. edulis* have been extirpated from much of its natural range, restoration often involves introduction of a breeding population. Care should be taken that this introduction does not lead to spreading of diseases, impacting shellfish or other species. This article focuses on the *Bonamia*-infection, since this is considered to be the most severe native oyster disease in north-west Europe, but it should be investigated whether other diseases, such as *Marteilia refringens*, are present in the breeding population and whether these can have a negative impact in the project area. If so, the type of measures recommended in this article to avoid spreading of the *Bonamia* infection should be applied to these other diseases.

6.2 | Detection of *Bonamia* presence and adherence to licence procedures

The recommendations in the following paragraph give guidance to using *Bonamia*-exposed or *Bonamia*-free oysters in the relevant circumstances. It should be noted that, even when these recommendations are adhered to, all oyster transports are subject to licensing, according to EU and/or national regulation. The project organizer should therefore always determine the *Bonamia*-infection status of the breeding population, applying the detection methods and following the EU-regulations. In addition, a transport licence (and other relevant licences) should be applied for at the competent authorities in the country where the restoration project is undertaken, and licence conditions should be adhered to at all time. While undergoing the detection process, the oysters to be transported or introduced should be kept in quarantine.

6.3 | Should *Bonamia*-exposed or *Bonamia*-free oysters be used for restoration purposes?

In 2017, NORA members drafted and agreed upon the following set of guidelines when employing *O. edulis* restoration projects (Pogoda et al., 2017, 2019).

1. If an oyster (*O. edulis* or otherwise) population is already present in the restoration area and the population is *Bonamia*-free:

Only *Bonamia*-free oysters can be introduced even if close to a *Bonamia*-infected region or (sub)area. As Figure 3 shows, there are several situations where a *Bonamia*-free area exists close to a *Bonamia*-infected area, and spread of the infection must be avoided by restoration attempts.

2. If an oyster (*O. edulis* or otherwise) population is already present in the restoration area and the population is *Bonamia*-exposed:

Either *Bonamia*-free or *Bonamia*-exposed oysters can be employed, but from a restoration perspective it is recommended to introduce *Bonamia*-exposed oysters in these areas, since they may have developed a certain level of tolerance or even resistance.

3. If an oyster (*O. edulis* or otherwise) population is absent in the restoration area:

Many current or planned restoration projects aim at reintroducing oyster populations in areas where oysters themselves are not present anymore, such as the open North Sea, Channel, or Irish Sea.

Arguments in favour of using *Bonamia*-free oysters in these open sea areas are:

- It is guaranteed that the infection does not spread through the restoration attempt.
- The oysters may be in a better condition, since they do not suffer from the illness, and therefore may better survive displacement stress.

The argument in favour of using *Bonamia*-exposed oysters in the open sea is that the infection is broadly present around these seas and, eventually, the infection may reach the restoration area sometime in the future, not only through *O. edulis*, but also through other hosts, possibly even *C. gigas*. In that case, *Bonamia*-exposed oysters, which may have developed a level of tolerance or resistance, could have an advantage.

A rational decision to use either *Bonamia*-exposed or *Bonamia*-free oysters is therefore subject to an assessment of the risks involved (risk of infection, risk of high mortality due to displacement stress combined with the infection etc.). However, it is impossible to make a reliable risk assessment on the basis of current scientific knowledge so that application of the precautionary principle, i.e. by only introducing *Bonamia*-free oysters in areas where no oyster population previously existed, is strongly recommended. This recommendation holds for the whole open North Sea, Channel, and Irish Sea and other open sea areas.

It should be noted that there is ongoing research into production of *Bonamia*-free oysters, produced from an infected, and therefore possibly *Bonamia*-tolerant or *Bonamia*-resistant population (Kamermaans et al., submitted). Should the rearing of tolerant/resistant and yet *Bonamia*-free oysters become possible, then this represents an opportunity to reduce the risk both of introducing the disease to new areas and of suffering high mortalities should the disease appear

at a later stage. In any case, it should be absolutely guaranteed that these oysters are free from the infection before they can be deployed. How this guarantee can be realized (detection accuracy, quarantine measures, etc.) should be researched and tested in detail and agreed by key experts before application in practice can be considered.

6.4 | Recommendations for future research

There are still many unknowns regarding the impact of *Bonamia* on *O. edulis* restoration activities, such as the impact of oyster density, temperature, and food availability on disease prevalence in natural systems (zu Ermgassen et al., 2020). The importance of developing research to understand both the mechanisms *Bonamia* tolerance or resistance, and ways in which scaling up the production of tolerant or resistant spat for restoration purposes was also identified and remains a pressing issue.

For the time being, it is important to emphasise that current best practice, from a legal as well as nature conservation perspective, is to use *Bonamia*-free *O. edulis* for restoration efforts in situations where no living oysters are currently present.

ORCID

Hein Sas  <https://orcid.org/0000-0002-9407-3950>
 Philine S.E. zu Ermgassen  <https://orcid.org/0000-0002-3409-0644>
 Bernadette Pogoda  <https://orcid.org/0000-0003-3997-426X>
 Joanne Preston  <https://orcid.org/0000-0002-2268-4998>
 Luke Helmer  <https://orcid.org/0000-0002-8315-6056>
 Zoë Holbrook  <https://orcid.org/0000-0002-0967-1770>
 Isabelle Arzul  <https://orcid.org/0000-0001-5436-5927>
 Antonio Villalba  <https://orcid.org/0000-0001-8049-0502>
 Bérenger Colsoul  <https://orcid.org/0000-0002-7891-8036>
 Alice Lown  <https://orcid.org/0000-0002-8331-8019>
 Verena Merk  <https://orcid.org/0000-0002-9034-3431>
 Nadescha Zwerschke  <https://orcid.org/0000-0003-4099-8269>

REFERENCES

- Abollo, E., Ramilo, A., Casas, S. M., Comesaña, P., Cao, A., Carballal, M. J., & Villalba, A. (2008). First detection of the protozoan parasite *Bonamia exitiosa* (Haplosporidia) infecting flat oyster *Ostrea edulis* grown in European waters. *Aquaculture*, 274, 201–207. <https://doi.org/10.1016/j.aquaculture.2007.11.037>
- Alderman, D. J. (1985). Fiche No. 16 Shell diseases of oysters. In C. J. Sinderman (Ed.), *Fiches d'identification des maladies et parasites des poissons, crustacés et mollusques*. Copenhagen: Conseil international pour l'exploration de la mer. ISSN 0109-2510.
- Alderman, D. J., & Jones, B. G. (1970). Shell disease of *Ostrea edulis*. L. *Fisheries Investigations London, Series 11*, Vol. 26, Part 8.
- Arzul, I., & Carnegie, R. B. (2015). New perspective on the haplosporidian parasites of molluscs. *Journal of Invertebrate Pathology*, 131, 32–42. <https://doi.org/10.1016/j.jip.2015.07.014>
- Arzul, I., Gagnaire, B., Bond, C., Chollet, B., Morga, B., Ferrand, S., ... Renault, T. (2009). Effects of temperature and salinity on the survival of *Bonamia ostreae*, a parasite infecting flat oysters *Ostrea edulis*. *Diseases of Aquatic Organisms*, 85, 67–75. <https://doi.org/10.3354/dao02047>
- Arzul, I., Langlade, A., Chollet, B., Robert, M., Ferrand, S., Omnes, E., ... Garcia, C. (2011). Can the protozoan parasite *Bonamia ostreae* infect larvae of flat oysters *Ostrea edulis*? *Veterinary Parasitology*, 179, 69–76. <https://doi.org/10.1016/j.vetpar.2011.01.060>
- Ayres, J. S., & Schneider, D. S. (2008). Two ways to survive an infection: What resistance and tolerance can teach us about treatments for infectious diseases. *Nature Reviews Immunology*, 8, 889–895.
- Bass, D., Ward, G. M., & Burki, F. (2019). Ascetosporea. *Current Biology*, 29, R7–R8. <https://doi.org/10.1016/j.cub.2018.11.025>
- Batista, F. M., López-Sanmartín, M., Grade, A., Navas, J. I., & Ruano, F. (2016). Detection of *Bonamia exitiosa* in the European flat oyster *Ostrea edulis* in southern Portugal. *Journal of Fish Diseases*, 39, 607–611. <https://doi.org/10.1111/jfd.12396>
- Baud, J. P., Gerard, A., & Naciri-Graven, Y. (1997). Comparative growth of *Bonamia ostreae* resistant and wild flat oysters, *Ostrea edulis*, in an intensive upwelling system. I. First year of experiment. *Marine Biology*, 130, 71–79. <https://doi.org/10.1007/s002270050226>
- Beck, M. W., Brumbaugh, R. D., Airoldi, L., Carranza, A., Coen, L. D., Crawford, C., ... Guo, X. (2011). Oyster reefs at risk and recommendations for conservation, restoration, and management. *Bioscience*, 61, 107–116. <https://doi.org/10.1525/bio.2011.61.2.5>
- Berthe, F. C. J., & Hine, P. M. (2003). *Bonamia exitiosa* Hine et al., 2001 is proposed instead of *B. exitiosus* as the valid name of *Bonamia* sp. infecting flat oysters *Ostrea chilensis* in New Zealand. *Diseases of Aquatic Organisms*, 57, 181–181. <https://doi.org/10.3354/dao057181>
- Bougrier, S., Tigé, G., Bachère, E., & Grizel, H. (1986). *Ostrea angasi* acclimatization to French coasts. *Aquaculture*, 58, 151–154. [https://doi.org/10.1016/0044-8486\(86\)90165-1](https://doi.org/10.1016/0044-8486(86)90165-1)
- Bromley, C., McGonigle, C., Ashton, E. C., & Roberts, D. (2016). Bad moves: Pros and cons of moving oysters—A case study of global translocations of *Ostrea edulis* Linnaeus, 1758 (Mollusca Bivalvia). *Ocean and Coastal Management*, 122, 103–115. <https://doi.org/10.1016/j.ocecoaman.2015.12.012>
- Carnegie, R. B., Burreson, E. M., Hine, M. P., Stokes, N. A., Audemard, C., Bishop, M. J., & Peterson, C. H. (2006). *Bonamia perspora* n. sp. (Haplosporidia), a parasite of the oyster *Ostreola equestris*, is the first *Bonamia* species known to produce spores. *Journal of Eukaryotic Microbiology*, 53, 232–245. <https://doi.org/10.1111/j.1550-7408.2006.00100.x>
- Carnegie, R. B., Hill, K. M., Stokes, N. A., & Burreson, E. M. (2014). The haplosporidian *Bonamia exitiosa* is present in Australia, but the identity of the parasite described as *Bonamia* (formerly *Mikrocytos*) *roughleyi* is uncertain. *Journal of Invertebrate Pathology*, 115, 33–40. <https://doi.org/10.1016/j.jip.2013.10.017>
- Carrasco, N., Villalba, A., Andree, K. B., Engelsma, M. Y., Lacuesta, B., Ramilo, A., ... Furones, M. D. (2012). *Bonamia exitiosa* (Haplosporidia) observed infecting the European flat oyster *Ostrea edulis* cultured on the Spanish Mediterranean coast. *Journal of Invertebrate Pathology*, 110, 307–313. <https://doi.org/10.1016/j.jip.2012.03.015>
- Cigarria, J., & Elston, R. (1997). Independent introduction of *Bonamia ostreae*, a parasite of *Ostrea edulis*, to Spain. *Diseases of Aquatic Organisms*, 29, 157–158. <https://doi.org/10.3354/dao029157>
- Culloty, S. C., Cronin, M. A., & Mulcahy, M. F. (2001). An investigation onto the relative resistance of Irish flat oysters *Ostrea edulis* L. to the parasite *Bonamia ostreae* (Pichot et al. 1980). *Aquaculture*, 199, 229–244. [https://doi.org/10.1016/S0044-8486\(01\)00569-5](https://doi.org/10.1016/S0044-8486(01)00569-5)
- Culloty, S. C., Cronin, M. A., & Mulcahy, M. F. (2004). Potential resistance of a number of populations of the oyster *Ostrea edulis* to the parasite *Bonamia ostreae*. *Aquaculture*, 237, 41–58. <https://doi.org/10.1016/j.aquaculture.2004.04.007>
- Culloty, S. C., & Mulcahy, M. F. (1996). Season-, age-, and sex-related variations in the prevalence of bonamiasis in flat oyster (*Ostrea edulis* L.) on the south coast of Ireland. *Aquaculture*, 144, 53–63. [https://doi.org/10.1016/S0044-8486\(96\)01290-2](https://doi.org/10.1016/S0044-8486(96)01290-2)

- Culloty, S. C., & Mulcahy, M. F. (2007). *Bonamia ostreae* in the native oyster *Ostrea edulis*. A Review. *Marine Environment and Health Series*, 29, 3–30.
- Culloty, S. C., Novoa, B., Pernas, M., Longshaw, M., Mulcahy, M. F., Feist, S. W., & Figueras, A. (1999). Susceptibility of a number of bivalve species to the protozoan parasite *Bonamia ostreae* and their ability to act as a vector for this parasite. *Diseases of Aquatic Organisms*, 37, 73–80.
- da Silva, P., & Villalba, A. (2004). Comparison of light microscopic techniques for the diagnosis of the infection of the European flat oyster *Ostrea edulis* by the protozoan *Bonamia ostreae*. *Journal of Invertebrate Pathology*, 85(2), 97–104. <https://doi.org/10.1016/j.jip.2003.12.010>
- da Silva, P. M., Fuentes, J., & Villalba, A. (2005). Growth, mortality and disease susceptibility of oyster *Ostrea edulis* families obtained from brood stocks of different geographical origins, through on-growing in the Ría de Arousa (Galicia, NW Spain). *Marine Biology*, 147, 965–977. <https://doi.org/10.1007/s00227-005-1627-4>
- de la Ballina, N., Villalba, A., & Cao, A. (2018). Proteomic profile of *Ostrea edulis* haemolymph in response to bonamiosis and identification of candidate proteins as resistance markers. *Diseases of Aquatic Organisms*, 128, 127–145. <https://doi.org/10.3354/dao03220>
- Diaz-Almela, E., Boudry, P., Launey, S., Bonhomme, F., & Lapegue, S. (2004). Reduced female gene flow in the European flat oyster *Ostrea edulis*. *Journal of Heredity*, 95, 510–516. <https://doi.org/10.1093/jhered/esh073>
- Dijkema, R. (1990). Balans van de activiteiten om de cultuur van de platte oester in Bretagne weer op gang te krijgen, einde 1989. *Vertaling Van Equinox*, 30.
- Elston, R. A., Kent, M. L., & Wilkinson, M. T. (1987). Resistance of *Ostrea edulis* to *Bonamia ostreae* infection. *Aquaculture*, 64, 237–242. [https://doi.org/10.1016/0044-8486\(87\)90328-0](https://doi.org/10.1016/0044-8486(87)90328-0)
- Engelsma, M., Culloty, S. C., Lynch, S. A., Arzul, I., & Carnegie, R. (2014). *Bonamia* parasites: A rapidly changing perspective on a genus of important mollusc pathogens. *Diseases of Aquatic Organisms*, 110, 5–23. <https://doi.org/10.3354/dao02741>
- Engelsma, M., Kerkhoff, S., Roozenburg, I., Haenen, O., van Gool, A., Siermans, W., ... Hummel, H. (2010). Epidemiology of *Bonamia ostreae* infecting European flat oysters *Ostrea edulis* from Lake Grevelingen, The Netherlands. *Marine Ecology Progress Series*, 409, 131–142. <https://doi.org/10.3354/meps08594>
- EU. (2006). Council directive 2006/88/EC on animal health requirements for aquaculture animals and products thereof, and on the prevention and control of certain diseases in aquatic animals of 24 October 2006.
- EU. (2015). Commission implementing decision 2015/1554, Official Journal of the European Union 23.9.2015 L 247/1.
- European Commission. (2017). Notification of the confirmation of bonamiosis in Kilkieran Bay, Co Galway, Ireland. Marine Institute, January 12th, 2017.
- Farley, C. A., Wolf, P. H., & Elston, R. A. (1988). A long-term study of “microcell” disease in oysters with a description of a new genus, *Mikrocytos* (g. n.), and two new species, *Mikrocytos mackini* and *Mikrocytos roughleyi* (sp. n.). *Fisheries Bulletin*, 86, 581–593.
- Flannery, G., Lynch, S. A., Carlsson, J., Cross, T. F., & Culloty, S. C. (2014). Assessment of the impact of a pathogen, *Bonamia ostreae*, on *Ostrea edulis* oyster stocks with different histories of exposure to the parasite in Ireland. *Aquaculture*, 432, 243–251. <https://doi.org/10.1016/j.aquaculture.2014.04.038>
- Flannery, G., Lynch, S. A., Longshaw, M., Stone, D., Martin, P., Ramilo, A., ... Culloty, S. C. (2014). Interlaboratory variability in screening for *Bonamia ostreae*, a protistan parasite of the European flat oyster *Ostrea edulis*. *Diseases of Aquatic Organisms*, 110, 93–99. <https://doi.org/10.3354/dao02717>
- Gercken, J., & Schmidt, A. (2014). Current status of the European oyster (*Ostrea edulis*) and possibilities for restoration in the German North Sea 2014, Bundesamt für Naturschutz
- Gervais, O., Chollet, C., Dubreuil, C., Durante, S., Feng, C., Hénard, C., ... Arzul, I. (2019). Involvement of apoptosis in the dialogue between the parasite *Bonamia ostreae* and the flat oyster *Ostrea edulis*. *Fish & Shellfish Immunology*, 93, 958–964. <https://doi.org/10.1016/j.fsi.2019.08.035>
- Gervais, O., Chollet, C., Renault, T., & Arzul, I. (2016). Flat oyster follows the apoptosis pathway to defend against the protozoan parasite *Bonamia ostreae*. *Fish & Shellfish Immunology*, 56, 322–329. <https://doi.org/10.1016/j.fsi.2016.07.021>
- Gervais, O., Renault, T., & Arzul, I. (2018). Molecular and cellular characterization of apoptosis in *O. edulis* a key mechanisms at the heart of host-parasite interactions. *Scientific Reports*, 8, 12494–12506. <https://doi.org/10.1038/s41598-018-29776-x>
- Grizel, H. (1985). Etude des récentes épizooties de l'huitre plate *Ostrea edulis* Linne et de leur impact sur l'ostréiculture bretonne, PhD/Dr Thesis. University of Montpellier, France.
- Grizel, H., Comps, M., Bonami, J. R., Cousserans, F., Duthoit, J. L., & LePennec, M. A. (1974). Research on the agent of digestive gland disease of *Ostrea edulis* Linné. *Science et Pêche Bulletin d'Information et de Documentation de l'Institut Scientifique et Technique Des Pêches Maritimes*, 240, 7–30. (In French)
- Gutierrez, A. P., Turner, F., Gharbi, K., Talbot, R., Lowe, N. R., Peñaloza, C., ... Houston, R. D. (2017). Development of a medium density combined-species SNP Array for Pacific and European oysters (*Crassostrea gigas* and *Ostrea edulis*). *G3 Genes Genomes Genetics*, 7, 2209–2218. <https://doi.org/10.1534/g3.117.041780>
- Haenen, O., Engelsma, M., & Beurden, S. (2011). Ziekten van vissen, schaal- en schelpdieren, van belang voor de Nederlandse aquacultuur. *Centraal Veterinair Instituut van Wageningen UR, Drukkerij CABRI BV, Lelystad*.
- Helmer, L., Hauton, C., Bean, T., Hendy, I., Harris-Scott, E., Bass, D., & Preston, J. (2020). Ephemeral detection of *Bonamia exitiosa* (Haplosporidia) in adult and larval European *Ostrea edulis* in the Solent (UK). *Journal of Invertebrate Pathology*, 174, 104217. <https://doi.org/10.1016/j.jip.2020.107421>
- Héral, M. (1990). Traditional oyster culture in France. In G. Barnabé & J. F. de IB Solbe (Eds.), *Aquaculture* (1st ed., Vol. 2, pp. 342–387). Chichester, UK: Ellis Horwood.
- Hervio, D., Bachère, E., Boulo, V., Cochenec, N., Vuillemin, V., Le Coguic, Y., ... Mialhe, E. (1995). Establishment of an experimental infection protocol for the flat oyster, *Ostrea edulis*, with the intrahaemocytic protozoan parasite, *Bonamia ostreae*: Application in the selection of parasite-resistant oysters. *Aquaculture*, 132, 183–194. [https://doi.org/10.1016/0044-8486\(94\)00342-L](https://doi.org/10.1016/0044-8486(94)00342-L)
- Hervio, D., Chagot, D., Godin, P., Grizel, H., & Mialhe, E. (1991). Localization and characterization of acid phosphatase activity in *Bonamia ostreae* (Asctospora), an intrahemocytic protozoan parasite of the flat oyster *Ostrea edulis* (Bivalvia). *Diseases of Aquatic Organisms*, 12, 67–70.
- Hine, P. M., Cochenec-Laureau, B. N., & Berthe, F. C. J. (2001). *Bonamia exitiosus* n. sp.(Haplosporidia) infecting flat oysters *Ostrea chilensis* in New Zealand. *Diseases of Aquatic Organisms*, 47, 63–72. <https://doi.org/10.3354/dao047063>
- Houziaux, J.-S., Kerckhof, F., Degrendele, K., Roche, M., & Norro, A. (2008). The hinder banks: yet an important area for the Belgian marine biodiversity? Final report, EV/45, *Belgian Science Policy*
- Huthnance, J., Weisse, R., Wahl, T., Thomas, H., Pietrzak, J., Souza, A. J., ... Woodworth, P. (2016). Recent Change - North Sea. In M. Quante & F. Colijn (Eds.), *North Sea region climate change assessment. Regional climate studies*. Cham: Springer. https://doi.org/10.1007/978-3-319-39745-0_3

- ICES. (2018). Report of the Working Group on Pathology and Diseases of Marine Organisms (WGPDMO), 13-17 February 2018, Riga, Latvia. ICES CM 2018/ASG:01. 42 pp.
- Korringa, P. (1951). Investigations on shell disease in the oyster, *Ostrea edulis* L. *ICES Rapports et procès-Verbaux Des réunions*, 128, 50–54.
- Korringa, P. (1976). *Farming the flat oysters of the genus Ostrea*. Amsterdam: Elsevier.
- Laing, I., Dunn, P., Peeler, E. J., Feist, S. W., & Longshaw, M. (2014). Epidemiology of *Bonamia* in the UK, 1982 to 2012. *Diseases of Aquatic Organisms*, 110, 101–111. <https://doi.org/10.3354/dao02647>
- Laing, I., Walker, P., & Areal, F. (2006). Return of the native—Is European oyster (*Ostrea edulis*) stock restoration in the UK feasible? *Aquatic Living Resources*, 19, 283–287. <https://doi.org/10.1051/alr:2006029>
- Lallias, D., Boudry, P., Lapegue, S., Kin, J. W., & Beaumont, A. R. (2010). Strategies for the retention of high genetic variability in European flat oyster (*Ostrea edulis*) restoration programmes. *Conservation Genetics*, 11, 1899–1910. <https://doi.org/10.1007/s10592-010-0081-0>
- Laramore, S., Krebs, W., Lave, A. L., & Gallagher, K. (2017). Survey of bivalve molluscs for *Bonamia* spp. and other parasitic pathogens in Florida East Coast Lagoons. *Journal of Shellfish Research*, 36, 379–390. <https://doi.org/10.2983/035.036.0211>
- Launey, S., Ledu, C., Boudry, P., Bonhomme, F., & Naciri-Graven, Y. (2002). Geographic structure in European flat oyster (*Ostrea edulis* L.) as revealed by microsatellite polymorphism. *Journal of Heredity*, 93, 40–47.
- Longshaw, M., Stone, D. M., Wood, G., Green, M. J., & White, P. (2013). Detection of *Bonamia exitiosa* (Haplosporidia) in European flat oysters *Ostrea edulis* cultivated in mainland Britain. *Diseases of Aquatic Organisms*, 106, 173–179. <https://doi.org/10.3354/dao02643>
- Lynch, S. A., Abollo, E., Ramilo, A., Cao, A., Culloty, S. C., & Villalba, A. (2010). Observations raise the question if the Pacific oyster, *Crassostrea gigas*, can act as either a carrier or a reservoir for *Bonamia ostreae* or *Bonamia exitiosa*. *Parasitology*, 137, 1515–1526. <https://doi.org/10.1017/S0031182010000326>
- Lynch, S. A., Armitage, D. V., Coughlan, J., Mulcahy, M. F., & Culloty, S. C. (2007). Investigating the possible role of benthic macroinvertebrates and zooplankton in the life cycle of the haplosporidian *Bonamia ostreae*. *Experimental Parasitology*, 115, 359–368. <https://doi.org/10.1016/j.exppara.2006.09.021>
- Lynch, S. A., Armitage, D. V., Wylde, S., Mulcahy, M. F., & Culloty, S. C. (2005). The susceptibility of young prespawning oysters, *Ostrea edulis*, to *Bonamia ostreae*. *Journal of Shellfish Research*, 24, 1019–1025. [https://doi.org/10.2983/0730-8000\(2005\)24\[1019:TSOYPO\]2.0.CO;2](https://doi.org/10.2983/0730-8000(2005)24[1019:TSOYPO]2.0.CO;2)
- Lynch, S. A., Flannery, G., Hugh-Jones, T., Hugh-Jones, D., & Culloty, S. C. (2014). Thirty-year history of Irish (Rossmore) *Ostrea edulis* selectively bred for disease resistance to *Bonamia ostreae*. *Diseases of Aquatic Organisms*, 110, 113–121. <https://doi.org/10.3354/dao02734>
- Madsen, L. (2017). *Bonamia ostreae* in the Limfjord in Denmark—Where is the parasite hiding? *18th International Conference on Diseases of Fish and Shellfish*, Belfast, UK, 4–8 September 2017, pp. 332.
- Møllergaard, S. (2008). *Declaration for surveillance programme/continuation of disease free status for bonamiosis and marteiliosis obtained in 2005 according to commissions decision 2005/104/EC*. Søborg: Danish Veterinary and Food Administration.
- Möbius, K. A. (1877). *Die Auster und die Austernwirtschaft*. Berlin: Verlag von Wiegandt, Hempel und Parey. 126 S
- Morga, B., Renault, T., Fauray, N., Chollet, B., & Arzul, I. (2011). Cellular and molecular responses of haemocytes from *Ostrea edulis* during in vitro infection by the parasite *Bonamia ostreae*. *International Journal of Parasitology*, 41, 755–764. <https://doi.org/10.1016/j.ijpara.2011.01.013>
- Morga, B., Renault, T., Fauray, N., Lerond, S., Garcia, C., Chollet, B., ... Arzul, I. (2017). Contribution of in vivo experimental challenges to understanding flat oyster *Ostrea edulis* resistance to *Bonamia ostreae*. *Frontiers in Cellular and Infection Microbiology*, 7, 1–13. <https://doi.org/10.3389/fcimb.2017.00433>
- Mortensen, S., Sælemyr, L., Skår, C.K., Bodvin, T., & Jelmert, A. (2016). The surveillance and control programme for bonamiosis and marteiliosis in European flat oysters (*Ostrea edulis*) and blue mussels (*Mytilus* sp.) in Norway 2015, Institute for Marine Research Norway, ISSN 1893-4536
- Mortensen, S., Sælemyr, L., Skår, C.K., Bodvin, T., & Jelmert, A. (2018). The surveillance and control programme for bonamiosis and marteiliosis in European flat oysters, *Ostrea edulis*, and blue mussels, *Mytilus* sp. in Norway in 2017, Havforskningen Instituttet, ISSN 1893-4536
- Mydlarz, L. D., Jones, L. E., & Harvell, C. D. (2006). Innate immunity, environmental drivers, and disease ecology of marine and freshwater invertebrates. *Annual Review of Ecology, Evolution, and Systematics*, 37, 251–288. <https://doi.org/10.1146/annurev.ecolsys.37.091305.110103>
- Naciri-Graven, Y., Haure, J., Gérard, A., & Baud, J. P. (1999). Comparative growth of *Bonamia ostreae* resistant and wild flat oyster *Ostrea edulis* in an intensive system. II. Second year of experiment. *Aquaculture*, 171, 195–208. [https://doi.org/10.1016/S0044-8486\(98\)00493-1](https://doi.org/10.1016/S0044-8486(98)00493-1)
- Naciri-Graven, Y., Martin, A. G., Baud, J. P., Renault, T., & Gérard, A. (1998). Selecting flat oyster *Ostrea edulis* for survival when infected with the parasite *Bonamia ostreae*. *Journal of Experimental Marine Biology and Ecology*, 224, 91–107. [https://doi.org/10.1016/S0022-0981\(97\)00171-8](https://doi.org/10.1016/S0022-0981(97)00171-8)
- Narcisi, V., Arzul, I., Cargini, D., Mosca, F., Calzetta, A., Traversa, D., ... Tiscar, P. G. (2010). Detection of *Bonamia ostreae* and *B. exitiosa* (Haplosporidia) in *Ostrea edulis* from the Adriatic Sea (Italy). *Diseases of Aquatic Organisms*, 89, 79–85. <https://doi.org/10.3354/dao02167>
- NORA. (2019). Information brochure on restoration projects for the native oyster restoration Alliance Edinburgh conference, May 21–23, 2019. Retrieved from <https://noraeurope.eu/restoration-projects/>
- OIE. (2008). Terrestrial manual 2008, Part 1—General information, par. B—Sample size.
- OIE. (2009). Infection with *Bonamia ostreae* in Norway 2009. Retrieved from https://www.oie.int/fileadmin/Home/eng/Health_standards/aahm/current/chapitre_bonamia_ostreae.pdf
- OIE. (2019). Manual of diagnostic tests for aquatic animals, Chapter 2.4.3. Infection with *Bonamia ostreae*. Retrieved from www.oie.int/fileadmin/Home/eng/Healthstandards/aahm/current/chapitre_bonamia_ostreae.pdf
- Olsen, O. T. (1883). *The piscatorial atlas of the North Sea, English and St. George's channels, illustrating the fishing ports, boats, gear, species of fish (how, where, and when caught), and other information concerning fish and fisheries*. London: Taylor and Francis.
- Orton, J. H. (1924a). An account of investigations into the cause or causes of the unusual mortality among oysters in English oyster beds during 1920 and 1921, Part I. *Ministry of Agriculture and Fisheries, Fisheries Investigations, Londonser.2*, 6, 1–199.
- Orton, J. H. (1924b). An account of investigations into the cause or causes of the unusual mortality among oysters in English oyster beds during 1920 and 1921, Part II. *Ministry of Agriculture and Fisheries, Fisheries Investigations, Londonser.2*, 6, 3–14.
- OSPAR Commission. (2009). Background document for *Ostrea edulis* and *Ostrea edulis* beds. OSPAR.
- Oysterecover. Establishing the scientific bases and technical procedures and standards to recover the European flat oyster production through strategies to tackle the main constraint, bonamiosis. Brussels: European Commission. <https://cordis.europa.eu/project/id/243583/reporting>
- Pardo, B. G., Álvarez-Dios, J. A., Cao, A., Ramilo, A., Gómez-Tato, A., Planas, J. V., ... Martínez, P. (2016). Construction of an *Ostrea edulis* database from genomic and expressed sequence tags (ESTs) obtained from *Bonamia ostreae* infected haemocytes: Development of an immune-enriched oligo-microarray. *Fish & Shellfish Immunology*, 59, 331–344. <https://doi.org/10.1016/j.fsi.2016.10.047>

- Pichot, Y., Comps, M., Tige, G., Grizel, H., & Rabouin, M. A. (1980). Recherches sur *Bonamia ostreae* gen. n., sp. n., parasite nouveau de l'huître plate *Ostrea edulis* L. *Revue Des Travaux de l'Institut des Pêches Maritimes*, 43, 131–140.
- Pogoda, B. (2019). Current status of European oyster decline and restoration in Germany. *Humanities*, 8, 9–21. <https://doi.org/10.3390/h8010009>
- Pogoda, B., Brown, J., Hancock, B., & von Nordheim, H. (2017). Berlin Oyster Recommendation on the future of native oyster restoration in Europe. Native oyster restoration in Europe—Current activities and future perspectives. Kick-off workshop Berlin, November 1st – 3rd, 2017, Berlin. Pp. 6
- Pogoda, B., Brown, J. H., Hancock, B., Preston, J., Pouvreau, S., Kamermans, P., ... von Nordheim, H. (2019). The Native Oyster Restoration Alliance (NORA) and the Berlin Oyster Recommendation: Bringing back a key ecosystem engineer by developing and supporting best practice in Europe. *Aquatic Living Resources*, 32, 13–22. <https://doi.org/10.1051/alr/2019012>
- Råberg, L., Graham, A. L., & Read, A. F. (2008). Decomposing health: Tolerance and resistance to parasites in animals. *Philosophical Transactions of the Royal Society B*, 364, 37–49. <https://doi.org/10.1098/rstb.2008.0184>
- Råberg, L., Sim, D., & Read, A. F. (2007). Disentangling genetic variation for resistance and tolerance to infectious diseases in animals. *Science*, 318, 812–814. <https://doi.org/10.1126/science.1148526>
- Ramilo, A., Gonzalez, M., Carballal, M. J., Darriba, S., Abollo, E., & Villalba, A. (2014). Oyster parasites *Bonamia ostreae* and *B. exitiosa* co-occur in Galicia (NW Spain): Spatial distribution and infection dynamics. *Diseases of Aquatic Organisms*, 110, 123–133. <https://doi.org/10.3354/dao02673>
- Ramilo, A., Navas, J. I., Villalba, A., & Abollo, E. (2013). Species-specific diagnostic assays for *Bonamia ostreae* and *B. exitiosa* in European flat oyster *Ostrea edulis*: Conventional, real-time and multiplex PCR. *Diseases of Aquatic Organisms*, 104, 149–161. <https://doi.org/10.3354/dao02597>
- Ronza, P., Cao, A., Robledo, D., Gómez-Tato, A., Álvarez-Dios, J. A., Hasanuzzaman, A. F. M., ... Martínez, P. (2018). Long-term affected flat oyster (*Ostrea edulis*) haemocytes show differential gene expression profiles from naïve oysters in response to *Bonamia ostreae*. *Genomics*, 110, 390–398. <https://doi.org/10.1016/j.ygeno.2018.04.002>
- Ryman, N., & Laikre, L. (1991). Effects of supportive breeding on the genetically effective population size. *Conservation Biology*, 5, 325–329. <https://doi.org/10.1111/j.1523-1739.1991.tb00144.x>
- Scottish Government. (2020). Retrieved from <https://www.gov.scot/publications/confirmed-designation-notice/>
- Sobolewska, H., & Beaumont, A. R. (2005). Genetic variation at microsatellite loci in northern populations of the European flat oyster (*Ostrea edulis*). *Journal of the Marine Biological Association of the U.K.*, 85, 955–960. <https://doi.org/10.1017/S002531540501194X>
- Swedish Board of Agriculture. (2010). Intention of declaration of diseases free status on historical grounds regarding infection with *Bonamia Ostrea*. European Commission, 31-10-2010, Brussels, Belgium
- Tubbs, C. (1999). *The ecology, conservation and history of the Solent*. Chichester: Packard Publishing Limited. 184 pp
- van Banning, P. (1991). Observations on *Bonamiasis* in the stock of the European flat oyster, *Ostrea edulis*, in the Netherlands, with special reference to recent developments in Lake Grevelingen. *Aquaculture*, 93, 205–211. [https://doi.org/10.1016/0044-8486\(91\)90232-V](https://doi.org/10.1016/0044-8486(91)90232-V)
- Vera, M., Carlsson, J., Carlsson, J. E. L., Cross, T., Lynch, S., Kamermans, P., ... Martínez, P. (2016). Current genetic status, temporal stability and structure of the remnant wild European flat oyster populations: Conservation and restoring implications. *Marine Biology*, 163, 1–17.
- Vera, M., Pardo, B. G., Cao, A., Vilas, R., Fernandez, C., Blanco, A., ... Martínez, P. (2019). Signatures of selection for Bonamiosis resistance in European flat oyster (*Ostrea edulis*): New genomic tools for breeding programs and management of natural resources. *Evolutionary Applications*, 12, 1781–1986. <https://doi.org/10.1111/eva.12832>
- WAHIS. (2020). OIE World Animal Health Information System. Retrieved from <https://www.oie.int/animal-health-in-the-world/the-world-animal-health-information-system/data-after-2004-wahis-interface/>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Sas H, Deden B, Kamermans P, et al. *Bonamia* infection in native oysters (*Ostrea edulis*) in relation to European restoration projects. *Aquatic Conserv: Mar Freshw Ecosyst*. 2020;30:2150–2162. <https://doi.org/10.1002/aqc.3430>

Annex

Table 1: Overview of *O. edulis* restoration projects NW Europe (project numbers correspond with Figure 2)

Nr	Project Name	Coordinating institute(s)	Contact person; website
1	RESTORE	Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research	B. Pogoda; www.awi.de/en/science/biosciences/shelf-sea-system-ecology/main-research-focus/european-oyster/restore
2	Borkum Reef NL	WWF-NL	E. Reuchlin; www.wwf.nl/schelpdierbanken
3	Gemini wind farm	WWF-NL, Gemini Windfarm, Edmelja BV	L. Folkerts; www.geminiwindpark.nl
4	Voordelta	ARK	K. van den Wijngaard; www.ark.eu/natuurontwikkeling/dieren/schelpdierbanken
5	Luchterduinen wind farm	North Sea Foundation, Natuur & Milieu	C. van Sluis, N. Strookman; www.natuurenmilieu.nl
6	Essex Native Oyster Restoration Initiative (ENORI)	Blue Marine foundation	M. Uttley; www.essexnativeoyster.com
7	Solent Oyster Restoration Project	Blue Marine foundation	J. Kean-Hammerson; www.bluemarinefoundation.com/project/solent
8	Wales Native Oyster Restoration Project	Natural Resources Wales	B. Wray; www.biodiversitysolutions.org.uk/wales-native-oyster-restoration-project/
9	Galway Bay	Marine Institute	O. Tully; www.cuanbeo.com
10	Lough Swilly Native Oyster Restoration	Marine Institute	www.nativeoysternetwork.org/portfolio/loughswilly/
11	Loch Craignish Native Oyster Restoration	CROMACH	D. Renton; www.nativeoysternetwork.org/portfolio/lochcraignish/
12	Dornoch Environmental Enhancement Project (DEEP)	Heriot-Watt University	B. Sanderson; www.theglenmorangiecompany.com/about-us/deep/
13	Humber Aquaculture Partnership	Yorkshire Wildlife Trust	D. Cowing; www.nativeoysternetwork.org/portfolio/the-humber-aquaculture-partnership-hap/

**Table 2: Spread of *Bonamia Ostreae* in North-West Europe: locations and references
(location numbers correspond with Figure 3)**

Nr	Country	Location	Status	Infection detection year/period	Source
1	Ireland	Cork Harbour (rossmore)	Infected	1987	McArdle et al., 1991; Culloty et al., 2007
2	Ireland	Tralee Bay	Free	-	Culloty et al., 2007
3	Ireland	Galway Bay (Clarenbridge)	Infected	1989	McArdle et al., 1991
4	Ireland	Kilkieran Bay	Infected	2016	European Commission, 2017
5	Ireland	Ballinakill	Infected	1993	Culloty et al., 2007
6	Ireland	Clew Bay	Infected	1988	Culloty et al., 2007
7	Ireland	Achill and Blacksod Bay	Infected	2002	Culloty et al., 2007
8	Ireland	Belmullet	Infected	2003	Culloty et al., 2007
9	Ireland	Lough Swilly	Infected	2006	Culloty et al., 2007
10	United Kingdom	Lough Foyle	Infected	2005	Culloty et al., 2007; Laing et al., 2014
11	United Kingdom	Strangford Lough	Infected	2008	Laing et al., 2014
12	United Kingdom	Holy Island	Free	-	Laing et al., 2014
13	United Kingdom	Humber Estuary	Uncertain	-	C. Gamble, personal communication, 2019
14	United Kingdom	Brancaster	Free	-	Laing et al., 2014
15	United Kingdom	River Blyth/Alde	Free	-	Laing et al., 2014
16	United Kingdom	Walton Backwaters	Infected	1982 - 2012	Laing et al., 2014
17	United Kingdom	River Colne	Infected	1982 - 2012	Laing et al., 2014
18	United Kingdom	River Blackwater	Infected	1982 - 2012	Laing et al., 2014
19	United Kingdom	River Crouch/Roach	Infected	1982 - 2012	Laing et al., 2014
20	United Kingdom	South Thames	Infected	1982 - 2012	Laing et al., 2014
21	United Kingdom	Chichester Harbour	Infected	1982 - 2012	Laing et al., 2014
22	United Kingdom	Langstone Harbour	Infected	1982 - 2012	Laing et al., 2014
23	United Kingdom	Portsmouth Harbour	Infected	1982 - 2012	Laing et al., 2014
24	United Kingdom	The Solent	Infected	1982 - 2012	Laing et al., 2014
25	United Kingdom	Poole Bay	Infected	1982 - 2012	Laing et al., 2014
26	United Kingdom	Poole Harbour	Infected	1982 - 2012	Laing et al., 2014
27	United Kingdom	Portland Harbour	Infected	1982 - 2012	Laing et al., 2014
28	United Kingdom	River Dart	Free	-	Laing et al., 2014
29	United Kingdom	River Avon	Free	-	Laing et al., 2014
30	United Kingdom	River Yealm	Free	-	Laing et al., 2014
31	United Kingdom	Plymouth Harbour	Infected	1982 - 2012	Laing et al., 2014
32	United Kingdom	River Fowey	Free	-	Laing et al., 2014
33	United Kingdom	River Fal	Infected	1982 - 2012	Laing et al., 2014
34	United Kingdom	River Helford	Infected	1982 - 2012	Laing et al., 2014
35	United Kingdom	Swansea Bay	Free	-	Laing et al., 2014

36	United Kingdom	River Cleddau	Infected	1982 - 2012	Laing et al., 2014
37	United Kingdom	Menai Strait	Infected	1982 - 2012	Laing et al., 2014
38	United Kingdom	Morecambe Bay	Free	-	Laing et al., 2014
39	United Kingdom	Loch Ryan	Free	-	B. Sanderson, personal communication, 2019
40	United Kingdom	West Loch Tarbert	Infected	1982 - 2012	Laing et al., 2014
41	United Kingdom	Lynn of Lorne, Loch Creran, Loch Etive	Infected	2019	Scottish Government, 2020
42	United Kingdom	Loch Sunart	Infected	1982 - 2012	Laing et al., 2014
43	United Kingdom	Little Loch Broom	Free	-	B. Sanderson, personal communication, 2019
44	United Kingdom	Durness	Free	-	B. Sanderson, personal communication, 2019
45	United Kingdom	Dornoch Firth	Infected	2019	Scottish Government, 2020
46	Sweden	Västra Götlands län	Free	-	Joyce et al., 2013; Swedish Board of Agriculture, 2010
47	Norway	Langstrand, Aust-Agder	Uncertain	-	S. Mortensen, personal communication, 2018; Mortensen et al., 2016; 2018
48	Norway	Hafrsfjord	Free	-	S. Mortensen, personal communication, 2018; Mortensen et al., 2016; 2018
49	Norway	Sveio, Hordaland	Free	-	S. Mortensen, personal communication, 2018; Mortensen et al., 2016; 2018
50	Norway	Aga, Bømlo, Hordaland	Free	-	S. Mortensen, personal communication, 2018; Mortensen et al., 2016; 2018
51	Denmark	Limfjorden	Infected	2017	Madsen, 2017; ICES, 2018
52	Germany	Helgoland	Free	-	B. Pogoda, personal communication, 2019
53	the Netherlands	Oosterschelde	Infected	1980	Kamermans, 2002; Sas et al., 2018
54	the Netherlands	Grevelingen	Infected	1988	Kamermans, 2002; Sas et al., 2018
55	the Netherlands	Brouwersdam	Infected	2017	Sas et al., 2018
56	the Netherlands	Bollen van de Ooster	Infected	2018	Sas et al., 2018
57	the Netherlands	Luchterduinen Windfarm	Free	-	Natuur & Milieu; North Sea Foundation, personal communication, 2018
58	the Netherlands	IJmuiden	Infected	2017	J. Groot, personal communication, 2017
59	the Netherlands	Wadden Sea	Uncertain	-	P. Jacobs et al., to be submitted

60	the Netherlands	Borkum Reef	Free	-	E. Reuchlin, personal communication, 2018
61	the Netherlands	Gemini Windfarm	Free	-	E. Reuchlin, personal communication, 2018
62	France	Baie du Mont Saint Michel	Infected	1979	I. Arzul, personal communication, 2019; Culloty et al., 2007
63	France	Brittany	Infected	1979	I. Arzul, personal communication, 2019; Culloty et al., 2007