

LETTER • OPEN ACCESS

## Arctic amplification causes earlier onset of seasonal tree growth in northeastern Siberia

To cite this article: Alexander V Kirilyanov *et al* 2024 *Environ. Res. Lett.* **19** 114091

View the [article online](#) for updates and enhancements.

You may also like

- [Gender inequality in global water security](#)  
Kaoru Kakinuma and Yoshihide Wada
- [Tales of river and ice: Indigenous art and water justice in the Arctic and the Amazon](#)  
Antonia Sohns, Alyssa Noseworthy,  
Gordon M Hickey *et al.*
- [Measuring the extent of trees outside of forests: a nature-based solution for net zero emissions in South Asia](#)  
David L Skole, Jay Samek, Swapan Mehra  
*et al.*

ENVIRONMENTAL RESEARCH  
LETTERS

## LETTER

Arctic amplification causes earlier onset of seasonal tree growth  
in northeastern Siberia

## OPEN ACCESS

RECEIVED  
13 May 2024REVISED  
22 September 2024ACCEPTED FOR PUBLICATION  
8 October 2024PUBLISHED  
22 October 2024

Original content from  
this work may be used  
under the terms of the  
[Creative Commons  
Attribution 4.0 licence](#).

Any further distribution  
of this work must  
maintain attribution to  
the author(s) and the title  
of the work, journal  
citation and DOI.



Alexander V Kirilyanov<sup>1,2,3,\*</sup> , Alexey I Kolmogorov<sup>4</sup>, Stefan Kruse<sup>5</sup> , Ulrike Herzschuh<sup>5</sup>, Alberto Arzac<sup>3</sup>,  
Lyudmila A Pestryakova<sup>4</sup>, Anatoly N Nikolaev<sup>4</sup>, Tatiana Bechuk<sup>1</sup> and Ulf Büntgen<sup>1,6,7</sup>

<sup>1</sup> Department of Geography, University of Cambridge, Cambridge CB2 3EN, United Kingdom

<sup>2</sup> Sukachev Institute of Forest SB RAS, Federal Research Center 'Krasnoyarsk Science Center SB RAS', Akademgorodok, Krasnoyarsk 660036, Russia

<sup>3</sup> Siberian Federal University, 79 Svobodnii, Krasnoyarsk 660041, Russia

<sup>4</sup> North-Eastern Federal University, 58 Belinsky str, Yakutsk 677027, Russia

<sup>5</sup> Polar Terrestrial Environmental Systems, Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, 14473 Potsdam, Germany

<sup>6</sup> Department of Geography, Masaryk University, 61137 Brno, Czech Republic

<sup>7</sup> Global Change Research Centre, 61300 Brno, Czech Republic

\* Author to whom any correspondence should be addressed.

E-mail: [ak2118@cam.ac.uk](mailto:ak2118@cam.ac.uk)

**Keywords:** boreal forest, climatic change, dendroecology, global warming, northern treeline, tree rings

Supplementary material for this article is available [online](#)

## Abstract

Although recent warming affects the high-northern latitudes at an unprecedented rate, little is known about its impact on boreal forests because *in situ* observations from remote ecosystems in Siberia are sparse. Here, we analyse the radial growth and climate sensitivity of 54 Cajander larches (*Larix cajanderi* Mayr.) from three sites across the northern treeline ecotone within the Omoloy river basin in northeastern Siberia. Three independent tree-ring width chronologies span 279–499 years and exhibit distinct summer temperature signals. These records further reveal evidence for sufficiently earlier onsets of growing seasons since the middle of the 20th century. This phenological shift coincides with rapidly increasing May temperatures and associated earlier snowmelt. Our findings reinforce the importance of high-precision ground measurements from remote regions in Siberia to better understand how warming-induced changes in the functioning and productivity of the boreal forest influence carbon, nutrient, and water cycle dynamics.

## 1. Introduction

The Arctic and subarctic are warming at historically unprecedented rate, which exceeds the global average temperature increase by four times (Rantanen *et al* 2022). Associated with this Arctic amplification (Francis *et al* 2017, Previdi *et al* 2021), the boreal forest zone is experiencing shifts in vegetation structure and productivity, northward vegetation expansion, permafrost degradation and greenhouse gas emission from permafrost thawing (Anisimov 2007, Schaefer *et al* 2011, Serreze and Barry 2011, Schuur *et al* 2015, Turetsky *et al* 2019). Understanding the speed and magnitude of these ecosystem changes is critical for assessing their impact on global carbon and water cycles (Chapin *et al* 2005, Bala *et al* 2007).

The dynamics of boreal forest ecosystems within the northern treeline ecotone, the transition from northern taiga to treeless tundra, is largely regulated by growing season temperatures (Paulsen and Körner 2014, Hansson *et al* 2023). Due to high climate sensitivity of the forest–tundra transition, these ecotones are widely used to monitor early responses to climate change (Shiyatov 1993, Esper and Schweingruber 2004, Kullman 2007, Dufour-Tremblay *et al* 2012). While a number of biotic and abiotic factors have to be considered when forest dynamics are studied (Rees *et al* 2019), current warming is believed to be among the main drivers initiating an expansion of trees into the tundra as well as forest densification within forest–tundra ecotones (Danby and Hik 2007, Fomin *et al* 2020, Dial *et al* 2022, Kruse *et al* 2023).

Tree-ring width (TRW) data are widely used as an indicator of past forest ecosystem response to climate change. In addition to tree establishment dates allowing identification of tree regeneration dynamics (Shiyatov 1993, Esper and Schweingruber 2004, Kharuk et al 2006, Devi et al 2008, Kirilyanov et al 2012, Grigor'ev et al 2019), TRWs provide approximations of tree growth and forest productivity (Knorre et al 2006, Devi et al 2008, Bouriaud et al 2015). Previous tree ring-based studies in Siberian high-latitude forest–tundra ecotones demonstrated a strong dependence of tree growth on summer temperature (Vaganov et al 1996, Naurzbaev et al 2002, Briffa et al 2004, D'Arrigo et al 2006, Hellmann et al 2016, Kirilyanov et al 2018, Büntgen et al 2021, Hantemirov et al 2021, 2022). However, Briffa et al (1998) described a persistent decoupling of tree growth from rising summer temperatures in northeastern Siberia. This so-called 'Divergence' phenomenon was also observed in northcentral Siberia (Kirilyanov et al 2020). Moreover, there is evidence that other climate-related factors can directly influence tree growth in Siberian forest–tundra ecotones, such as winter precipitation (Kharuk et al 2023), snowmelt dates (Vaganov et al 1999, Kirilyanov et al 2003), within-growing season frosts (Gurskaya 2014, 2021), floods (Tei et al 2019a, Meko et al 2020) and soil temperature and moisture (Nikolaev et al 2011, Fujii et al 2022, Kharuk et al 2023, Liang et al 2023). Despite a long history of local and large-scale tree-ring studies in the high-northern latitudes, uncertainties remain with reference to how current and future climate and environmental changes will affect tree growth and forest productivity in northeastern Eurasia.

Here, we explore the radial growth and climate sensitivity of three larch sites across the northern treeline ecotone within the Omoloy river basin of northern Yakutia. Although hardly accessible, this extremely remote region is of great importance for revealing the direct ecological response of undisturbed forest–tundra ecosystems to rising temperatures. We hypothesize that tree growth in this region is directly affected by Arctic amplification, and should be reflected in changing patterns of radial tree growth climatic response.



## 2. Material and methods

The study area is located at the northern treeline in northeastern Siberia between 70°31'36" N and 70°56'00" N along the 132°47' E longitude (figure 1). This region is dominated by deciduous *Larix cajanderi* Mayr. that is well adapted to growing on nutrient-deficient permafrost soils under harsh winter climate and short growing season with 24 h daylight (Abaimov et al 1997). Climate in this region

is extremely continental, with annual mean temperatures of  $-13.3$  °C (WMO 226 609 'Yubileynaya'; 1936–2022; meteo.ru) (figure 2). The warmest and coldest months are July and January with monthly temperatures of 11.2 °C and  $-35.3$  °C, respectively. June is usually the first month with positive temperature means (7.2 °C) that last until September (2.0 °C). Annual precipitation totals are 276 mm (1936–2015), of which 42% fall between June and August. Seasonal temperature means significantly increased since 1975 at a rate of 0.6 ( $P < 0.005$ ), 0.6 ( $P < 0.005$ ) and 0.5 °C/decade ( $P < 0.05$ ), in June–July, summer (June–August) and annually, respectively (figure 2), whereas summer precipitation totals significantly decreased by 10.9 mm decade<sup>-1</sup> ( $P < 0.05$ ). Importantly, data on monthly precipitation totals available from the All-Russian Research Institute of Hydrometeorological Information—World Data Center (RIHMI-WDC) are corrected for systematic errors according to Bogdanova et al (2007) and Bogdanova and Gavrilova (2008).

Increment core samples were collected in undisturbed, uneven-aged *Cajander* larch stands at three sites along a north–south transect within the forest–tundra ecotone (figure 1). The northernmost site N was established in sparse forest on the first terrace of Omoloy river, the only forested area at this location (Miesner et al 2022a, 2022b). Stand density was 700–800 trees/hectare, with larch trees reaching 6.5 m in height and mean DBH (diameter at breast height, 1.3 m) of 10.0 cm. The middle site M represents a closed forest treeline, with varying stand density from 2100 trees/hectare in open forest to 6400 trees/hectare in closed forest. Trees were up to 8 m in height with a DBH of 11.3 cm. The southernmost site S was established in closed forest with the mean tree stand density of 5600 trees/hectare and individual trees being up to 10 m with a DBH of 10.6 cm. In late June 2014, the mean active layer thickness was 10–25 cm, 33–35 cm and  $\sim 35$  cm at N, M and S, respectively. Detailed information on ground vegetation and stand parameters was provided by Miesner et al (2022a). During sampling, we mostly aimed at coring old-growth trees, which were distributed over an area of a one-two hectares. In total, from 16 to 21 trees per site were cored with the oldest trees reaching from 279 (site N) to 499 (S) and the mean segment length (MSL) ranging from 126 (N) to 281 (S) years (table 1).

All increment cores were air-dried and mounted on wooden supports, with their surface cut and contrasted with chalk to increase the visibility of tree-ring boundaries. TRW was measured on a LINTAB measuring system (RINNTECH e.K., Heidelberg, Germany). The individual TRW series were cross-dated using the TSAP-Win (Rinn 2003), and cross-dating accuracy was statistically verified with COFECHA (Version 6.02P). Individual



**Table 1.** Chronology characteristics (MSL = mean segment length, TRW = tree-ring width, CS = coefficient of sensitivity; Rbar = inter-series correlation, EPS = expressed population signal).

Site	N of series	Period CE	MSL	Mean TRW, mm	Mean TRW >150 y, mm	Mean CS	Mean Rbar	Mean EPS
N	21	1736–2014	126	0.56 ± 0.30	0.12 ± 0.06	0.397	0.415	0.905
M	17	1740–2014	190	0.27 ± 0.20	0.12 ± 0.06	0.415	0.538	0.946
S	16	1516–2014	281	0.21 ± 0.15	0.10 ± 0.05	0.408	0.469	0.914

Note: All statistics except Mean TRW and Mean TRW >150 y. were calculated for standardized chronologies.

TRW series were standardized with cubic smoothing splines of 50% frequency-response cutoff at 2/3 of the individual series lengths. Bi-weight robust

means of the individual measurement series were calculated to produce dimensionless TRW index chronologies (Cook 1985). The standard version

of the chronologies was chosen for further analyses to preserve potential longer-term variability for comparison against different climate parameters. The coefficient of sensitivity (CS), mean inter-series correlation ( $R_{\text{bar}}$ ) and expressed population signal (EPS) were calculated using the latest version of the ARSTAN software ([www.geog.cam.ac.uk/research/projects/dendrosoftware/](http://www.geog.cam.ac.uk/research/projects/dendrosoftware/)), last accessed on 25 March 2024).

To determine the most important climatic factors that control tree radial growth, the TRW standard chronologies were correlated against instrumental records of monthly and seasonal temperature means and precipitation totals from the nearest meteorological station Yubileinaya over the common period 1936–2014. Pairwise correlations were calculated, which ignore missing climate data values. To assess the temporal changes in the relationships between TRW chronologies and climate records, Pearson's correlation coefficients were calculated over 1936–1974 and 1975–2014, as well as using moving 25 year windows. To eliminate the influence of climate data trends on tree growth response to climate, we correlated the residual TRW index chronologies with the detrended climate records (Ols *et al* 2023). We also evaluated time shifts in snowmelt dates available for the meteorological station Yubileinaya.

### 3. Results

#### 3.1. Tree radial growth

Trees at the study sites are characterized by low radial stem growth rate, and up to 3.3% of tree rings can be locally absent (M) (table 1). The mean TRW increases from  $0.21 \pm 0.15$  mm (S) to  $0.56 \pm 0.30$  mm (N) with decreasing chronology length and MSL. The mean growth rate of trees at a mature stage (cambial age >150 years), when the age trend in tree radial growth is least pronounced, is similar among the sites, with slightly lower values at S. All the index chronologies are characterized by high CS >0.3 and  $R_{\text{bar}}$  >0.4. Mean EPS (calculated for the 50 year periods shifted by 25 years) is >0.9 since at least 1840. These statistics indicate a strong common signal in individual TRW series from each site and confirm that the sample depth is sufficient for dendroclimatic analysis at all the study sites.

The three TRW chronologies demonstrate a high coherence of inter-annual and multi-decadal variation (figure 3). The coefficient of correlation for the period since 1809, with data for at least three trees in each chronology, ranges from 0.70 ( $P < 0.001$ ) for the most distant sites N and S to 0.85 ( $P < 0.001$ ) for two southern sites M and S. Since 1900, the highest TRW indices were observed in the 1930s and the early 1940s, followed by a gradual TRW decrease until the 1970–80 s, and a growth increase afterwards (figure 3).

#### 3.2. TRW climate response

The growth-climate response analysis of the local chronologies back to 1936 shows that TRW indices at the northernmost site N correlate significantly positively with mean June and July temperatures ( $P < 0.01$ ) (figure 4(A)). TRW data from the other two sites exhibit positive relations with mean June temperatures ( $P < 0.001$ ). All three local chronologies correlate positively with summer (June–July and June–August) temperature means ( $r$  ranges from 0.26 to 0.42,  $P < 0.05$ ). The strongest correlation is found for the southernmost site S with June temperature ( $r = 0.56$ ,  $P < 0.001$ ). Correlations of standard TRW chronologies with observed monthly and seasonal precipitation (figure S1(A)) and residual TRW chronologies with detrended precipitation (figure S1(B)) do not show any consistent results between the sites.

Detailed dendroclimatic analysis with temperatures shows that TRW mostly depends on June and JJ temperatures during the 1936–1974 period ( $P < 0.05$ ) (figure 4(B)). TRW chronology from the northernmost site N significantly ( $P < 0.05$ ) correlates also with JJA temperature. However, the correlation pattern considerably changes for the later period from 1975 (figure 4(C)). June temperature is still the most important climate variable for tree growth at all three sites ( $P < 0.05$ ), and N continues to depend on July temperatures ( $P < 0.05$ ). However, N and M also show a strong dependence ( $P < 0.01$ ) on May temperature. In the case of seasonal temperature means, correlations remain statistically significant only for N and S ( $P < 0.05$ ). Importantly, correlations of the residual TRW chronologies with detrended temperature variables show a similar pattern with increasing importance of May temperature for N and M over the 1936–1974 period (figure S2).

Running correlation analysis of growth-climate relation shows a continuously increasing influence of May temperatures on TRW at all three sites (figure 5(A)). At site N, correlations become significant during the 1959–1983 period and remain high since then. At sites M and S, May temperature is important only during a short interval, with the first significant correlations observed for the 1987–2011 period. June temperatures significantly influence tree radial growth at all the three sites throughout most of the period from 1936, with the generally higher correlations up to  $r = 0.77$  ( $P < 0.001$ ) at S (figure 5(B)). The role of July temperature fluctuated at the  $P < 0.05$  significance level at N, whereas at M and S, it is mostly insignificant throughout the period (figure 5(C)). Surprisingly, running correlations with monthly precipitation totals do not demonstrate increased positive influence of precipitation for the period of temperature increase (figure S4). On the contrary, TRW chronologies significantly ( $P < 0.05$ ) negatively correlate with June precipitation totals (figure S4(B)) in recent decades. Positive correlations are observed




Figure 3. Standard tree-ring width (TRW) chronologies. The chronologies were smoothed with a 30 year spline.





Figure 4. Correlation coefficients of standard tree-ring width chronologies and monthly temperature means from previous September to current September, and seasonal temperature means for June–July (JJ) and summer (JJA) for the period 1936–2014 (A) and selected months and seasonal means (JJ and JJA) for the 1936–1974 (B) and 1975–2014 (C) periods. Horizontal line indicates the significance level  $P < 0.05$ .

with July (figure S4(C)) and August (figure S4(D)) precipitations during some intervals during the first half of the analysed period from 1936. The running correlations of the residual TRW index chronologies with detrended monthly temperature means (figure S3), and precipitation totals (figure S5) demonstrate similar tendencies, with the only major difference of significant negative correlations between the TRW residual chronology from N and detrended


July precipitation starting from the 1973–1997 period (figure S5(C)).

### 3.3. Changes of climate parameters crucial for tree growth onset

To reveal factors leading to the shift of the temperature dependency of tree growth to earlier dates, the dynamics of May temperatures and snowmelt dates were explored (figure 6). Mean May temper-



**Figure 5.** 25 year window running correlations between the standard tree-ring width chronologies N, M and S and monthly temperature means of May (A), June (B) and July (C). Grey horizontal line indicates the significance level  $P < 0.05$ .



**Figure 6.** May temperature (TM) and the day of the year (DOY) of complete snowmelt (Snowmelt) based on data from the nearest meteorological station Yubileynaya. The lines are smoothed with distance-weighted least squares fitting in STATISTICA 64 (StatSoft, Inc.).

ature demonstrates a clear positive trend from the late 1950s. During the last decades, May temperature was on average higher than for any other decade-long interval over the period of instrumental meteorological observations and increased from a decade-long mean  $-6.8^{\circ}\text{C}$  during the 1956–1965 period to  $-2.0^{\circ}\text{C}$  in 1995–2014. Importantly, positive ( $>0^{\circ}\text{C}$ ) May temperature means started to appear in recent decades and were recorded in 1990 ( $2.8^{\circ}\text{C}$ ), 2007 ( $0.1^{\circ}\text{C}$ ) and 2013 ( $0.2^{\circ}\text{C}$ ). On the contrary, snowmelt dates demonstrate a decreasing trend from the 1960s, evidencing a shift to earlier snowmelt from 154th day of the year (DOY) as a mean for the 1949–1958 period to 135th DOY in 2011–2020.

#### 4. Discussion

Radial tree growth at the northern treeline ecotone in Siberia is usually extremely low due to harsh climate conditions, which synchronize tree growth not only within single sites, but also at distances of several hundred kilometers (Vaganov *et al* 1996, Esper *et al* 2010, Hellmann *et al* 2016, Shestakova *et al*

2016, Büntgen *et al* 2021, Hantemirov *et al* 2021, 2022). Low mean annual increment, high inter-series correlations of tree growth in our study region and the results of the dendroclimatic analysis confirm the importance of summer temperature as a primary climatic driver that modulates year-to-year variability of tree radial growth in northern Yakutia, with the early summer period (June) showing the highest impact. These results coincide with previous findings for Siberia (MacDonald *et al* 1998, Hughes *et al* 1999, Vaganov *et al* 1999, Kirilyanov *et al* 2003, 2018, Hantemirov *et al* 2022, Kolmogorov *et al* 2023), as well as for the wider area of Eurasia and the Northern Hemisphere (Briffa *et al* 2004, 2013, St. George 2014, Hellmann *et al* 2016, Devi *et al* 2020). The period of the growing season with significant correlations between TRW and monthly temperature may considerably vary between regions from one to four-five months (Büntgen *et al* 2021), but early summer temperature usually demonstrates higher correlations, especially in Siberia (Vaganov *et al* 1999, Kirilyanov *et al* 2018). Rapid onset of tree cambial activity after snowmelt and daily temperature increase above the

physiologically defined threshold at the beginning of the growing season is the key adaptation of treeline larch in Siberia (Vaganov *et al* 1999, Kirdyanov *et al* 2003). Trees take advantage of a 24 h photoperiod, which enables them to complete tree-ring formation within several weeks of the short and cold growing season (see data on seasonal tree radial growth at the more southern location, but with short seasonal growth in Bryukhanova *et al* 2013, Rinne *et al* 2015a, 2015b). However, neither the crucial temperature threshold nor the timing of growing seasons of larch at the northern treeline are currently known despite the importance of these parameters for understanding the physiology of larch trees and predicting future vegetation shifts in the boreal zone in Siberia. Remote sensing data (Buitenwerf *et al* 2015, Dronova and Taddeo 2022) and tree-ring growth modelling (Vaganov *et al* 1999, 2006, Shishov *et al* 2016, Kang *et al* 2023, Shishov *et al* 2023) may provide rough estimates of these values, but require on-site validation.

Our TRW chronologies demonstrate a recent shift of tree growth dependence on temperature to earlier dates, which can indicate an earlier start of the growing season (Gao *et al* 2022). This shift coincides with the temporal changes of two climate parameters that directly affect the timing of the growing season onset: snowmelt and temperature preceding and during early growing season (Vaganov *et al* 1999, Kirdyanov *et al* 2003, Livensperger *et al* 2016). An approximately 19 d earlier snowmelt and 4.8 °C higher May temperature in recent decades compared to the middle of the 20th century can indeed induce earlier activation of seasonal growth, which is currently observed in extratropical ecosystems in the Northern hemisphere (Linderholm 2006, Menzel *et al* 2006, Piao *et al* 2015, Büntgen *et al* 2022). Surprisingly, the significant correlations between TRW chronologies and May temperature during the period from 1975 were found for the two northern sites N and M, but not the southern site S, meaning the slower reaction of tree growth at the southernmost location to temperature increase. Unfortunately, there are no instrumental meteorological observations in the region that could be representative to each of the study sites and confirm the site-specific tree growth response to warming. The only meteorological station Yubileynaya is latitudinally located in the middle of the study transect, but within closed forest of the treeline ecotone, correspondent to site S. However, due to the proximity of the sites and the flat topography of the study area we can consider the regional climate rather homogeneous, with slightly lower air temperature at the northern sites.

The site-specificity of TRW climatic response to warming observed in the study region is a known phenomenon for the boreal permafrost zone:

trees at geographically close locations can demonstrate different climate response due to local soil thermo-hydroclimate conditions (Kirdyanov *et al* 2013, Bryukhanova *et al* 2015, Kirdyanov *et al* 2024). Intra- and inter-species competition can also act as a factor influencing tree radial growth dynamics and climatic response because soil temperature conditions, active layer thickness and nutrient availability largely depend on tree stand density and ground vegetation (Brown 1966, Shur and Jorgenson 2007, Yin *et al* 2017, Fedorov *et al* 2019, Knorre *et al* 2019, Stuenzi *et al* 2021). Trees at site S are prone to the highest level of competition due to the highest tree density and most developed ground vegetation (Miesner *et al* 2022a). Although the active layer thickness at site S is similar to site M and 10–25 cm deeper than at N, higher root competition for water and nutrients can explain a delayed reaction of TRW to warming (figure 4(A)) (Wieczorek *et al* 2017). Denser ground vegetation at site S can trap and insulate snow delaying snowmelt and leading to later start of growing season. However, the exact reason of the difference in timing of tree seasonal growth activation between the study sites may not be resolved due to the lack of on-site data.

The statement on site-specific tree growth response in the permafrost zone may seem to contradict earlier observations on high similarity of tree growth at distant sites in northern Siberia (for example, Vaganov *et al* 1996, Hellmann *et al* 2016). However, the approaches to sampling in dendroclimatology and dendroecology are different (Fritts 1976, Schweingruber 1996). While dendroclimatic studies in northern Siberia are often focused on extracting summer temperature signal, and researchers pay special attention to cool and moist sites with similar environmental settings (Vaganov *et al* 1996, Hughes *et al* 1999, Jacoby *et al* 2000, Briffa *et al* 2004, Esper *et al* 2010, Büntgen *et al* 2021, Hantemirov *et al* 2022), dendroecology is focused on understanding the effect of different environmental and climatic drivers on tree growth, and sampling is usually conducted in environmentally diverse ecosystems (Kirdyanov *et al* 2020, Gurskaya *et al* 2021, Buchwal *et al* 2023, Kharuk *et al* 2023).

Our study underlines the importance of the on-ground data for detecting the current changes in vegetation. Such data are especially important for remote regions exposed to rapid climate change, like northern Siberia (Gauthier *et al* 2015, Hantemirov *et al* 2022). Generally, despite recent advances in data collection, analyses, and modelling (Lloyd *et al* 2011, Büntgen *et al* 2014, 2021, Tei *et al* 2019b, Tei and Sugimoto 2020, Miesner *et al* 2022a, 2022b, Buchwal *et al* 2023, Liang *et al* 2023, etc), the way treeline forests behave in northeastern Siberia is still poorly understood. Sufficient efforts are needed to obtain

*in situ* field data and gain knowledge, for example, on tree cambium activity and the exact timing of growing season, vegetation productivity and plant-permafrost interactions to define the adaptation potential of trees growing under extreme environmental conditions that are currently rapidly changing.

## 5. Conclusion

Our study demonstrates that radial growth of Cajander larch trees within the treeline ecotone in northeastern Siberia mainly depends on early summer temperature. In recent decades, a shift to earlier growing seasons is observed due to warmer May conditions and earlier snowmelt. The climatic response of tree growth is site-specific, being partly dependent on local conditions that include ecological settings of the sites and intra- and interspecies competition. To detect current ecosystem changes caused by global warming, on-ground data are required, especially for remote regions that are characterized by low anthropogenic pressure, but sensitive to global climate change.

## Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

## Acknowledgment

This work was supported by the project of the Ministry of Education and Science of the Russian Federation FWES-2024-0023. U H was supported by ERC Consolidator Grant Glacial Legacy (Grant No. 772852), S K received support from DFG (project No. 448651799) and U B received funding from the Czech Science Foundation (# 23-08049S; Hydro8), the ERC Advanced Grant (# 882727; Monostar), and the ERC Synergy Grant (# 101118880; Synergy-Plague).

## ORCID iDs

Alexander V Kirilyanov  <https://orcid.org/0000-0002-6797-4964>

Stefan Kruse  <https://orcid.org/0000-0003-1107-1958>

## References

- Abaimov A P, Bondarev A I, Zyryanova O A and Shitova S A 1997 *Polar Forests of Krasnoyarsk Region* (Novosibirsk Nauka) p 208 in Russian
- Anisimov O A 2007 Potential feedback of thawing permafrost to the global climate system through methane emission *Environ. Res. Lett.* **2** 045016
- Bala G, Caldeira K, Wickett M, Phillips T J, Lobell D B, Delire C and Mirin A 2007 Combined climate and carbon-cycle effects of large-scale deforestation *Proc. Natl Acad. Sci.* **104** 6550–5
- Bogdanova E G and Gavrilova S Y 2008 Correction of the precipitation time series nonhomogeneity caused by replacement of the Nipher shielded rain gauge by a Tretyakov precipitation gauge *Russ. Meteorol. Hydrol.* **33** 525–36
- Bogdanova E G, Il'in B M and Gavrilova S Y 2007 Advanced methods for correcting measured precipitation and results of their application in the polar regions of Russia and North America *Russ. Meteorol. Hydrol.* **32** 229–44
- Bouriaud O, Teodosiu M, Kirilyanov A V and Wirth C 2015 Influence of wood density in tree-ring based annual productivity assessments and its errors in Norway spruce *Biogeosciences* **12** 6205–17
- Briffa K R, Melvin T M, Osborn T J, Hantemirov R M, Kirilyanov A V, Mazepa V S, Shiyatov S G and Esper J 2013 Reassessing the evidence for tree-growth and inferred temperature change during the Common Era in Yamalia, northwest Siberia *Quat. Sci. Rev.* **72** 83–107
- Briffa K R, Osborn T J and Schweingruber F H 2004 Large-scale temperature inferences from tree rings: a review *Glob. Planet. Change* **40** 11–26
- Briffa K R, Schweingruber F H, Jones P D, Osborn T J, Shiyatov S G and Vaganov E A 1998 Reduced sensitivity of recent tree-growth to temperature at high northern latitudes *Nature* **391** 678–82
- Brown R J E 1966 Influence of vegetation on permafrost *Proc.: Permafrost Int. Conf.* vol 1963 pp 20–25
- Bryukhanova M V, Fonti P, Kirilyanov A V, Siegwolf R T W, Saurer M, Pochebyt N P, Churakova (Sidorova) O V and Prokushkin A S 2015 The response of  $\delta^{13}\text{C}$ ,  $\delta^{18}\text{O}$  and cell anatomy of *Larix gmelinii* tree rings to differing soil active layer depths *Dendrochronologia* **34** 51–59
- Bryukhanova M V, Kirilyanov A V, Prokushkin A S and Silkin P P 2013 Specific features of xylogenesis in Dahurian larch, *Larix gmelinii* (Rupr.) Rupr., growing on permafrost soils in Middle Siberia *Russ. J. Ecol.* **44** 361–6
- Buchwal A, Rachlewicz G, Heim B and Juhls B 2023 Trees on the tundra: warmer climate might not favor prostrate *Larix* tree but *Betula nana* shrub growth in Siberian tundra (Lena River Delta) *Agr. For. Meteorol.* **339** 109543
- Buitenwerf R, Rose L and Higgins S 2015 Three decades of multi-dimensional change in global leaf phenology *Nat. Clim. Change* **5** 364–8
- Büntgen U et al 2021 The influence of decision-making in tree ring-based climate reconstructions *Nat. Commun.* **12** 3411
- Büntgen U, Kirilyanov A V, Hellmann L, Nikolayev A and Tegel W 2014 Cruising an archive: on the palaeoclimatic value of the Lena Delta *Holocene* **24** 627–30
- Büntgen U, Pierrattei A, Krusic P J, Esper J, Sparks T and Crivellaro A 2022 Plants in the UK flower a month earlier under recent warming *Proc. R. Soc. B* **289** 20212456
- Chapin F S et al 2005 Role of terrestrial ecosystems in arctic summer warming *Science* **310** 657–60
- Cook E R 1985 A time series analysis approach to tree-ring standardization *PhD Thesis* Lamont-Doherty Geological Observatory, New York
- D'Arrigo R, Wilson R and Jacoby G 2006 On the long-term context for late twentieth century warming *J. Geophys. Res. Atmos.* **111** D03103
- Danby R K and Hik D S 2007 Evidence of recent treeline dynamics in Southwest Yukon from Aerial Photographs *Arctic* **60** 411–20
- Devi N M, Kukarskih V V, Galimova A A, Mazepa V S and Grigoriev A A 2020 Climate change evidence in tree growth and stand productivity at the upper treeline ecotone in the Polar Ural Mountains *For. Ecosyst.* **7** 7
- Devi N, Hagedorn F, Moiseev P, Bugmann H, Shiyatov S, Mazepa V and Rigling A 2008 Expanding forests and changing growth forms of Siberian larch at the polar urals

- treeline during the 20th century *Glob. Change Biol.* **14** 1581–91
- Dial R J, Maher C T, Hewitt R E and Sullivan P F 2022 Sufficient conditions for rapid range expansion of a boreal conifer *Nature* **608** 546–51
- Dronova I and Taddeo S 2022 Remote sensing of phenology: towards the comprehensive indicators of plant community dynamics from species to regional scales *J. Ecol.* **110** 1460–84
- Dufour-Tremblay G, Lévesque E and Boudreau S 2012 Dynamics at the treeline: differential responses of *Picea mariana* and *Larix laricina* to climate change in eastern subarctic Québec *Environ. Res. Lett.* **7** 044038
- Esper J, Frank D, Büntgen U, Verstege A, Hantemirov R M and Kirilyanov A V 2010 Trends and uncertainties in Siberian indicators of 20th century warming *Glob. Change Biol.* **16** 386–98
- Esper J and Schweingruber F H 2004 Large-scale treeline changes recorded in Siberia *Geophys. Res. Lett.* **31** L06202
- Fedorov A N, Konstantinov P Y, Vasiliev N F and Shestakova A A 2019 The influence of boreal forest dynamics on the current state of permafrost in Central Yakutia *Pol. Sci.* **22** 100483
- Fomin V V, Mikhailovich A P and Shiyatov S G 2020 Trees in the upper treeline ecotone in the Polar Urals: centuries-old change and spatial patterns *Mt. Res. Dev.* **40** R32–40
- Francis J A, Vavrus S J and Cohen J 2017 Amplified Arctic warming and mid-latitude weather: new perspectives on emerging connections *WIREs Clim. Change* **8** e474
- Fritts H C 1976 *Tree-Rings and Climate* (Academic Press) pp 576
- Fujii K, Yasue K and Matsuura Y 2022 Tree ring evidence of rapid development of drunken forest induced by permafrost warming *Glob. Change Biol.* **28** 3920–8
- Gao S et al 2022 An earlier start of the thermal growing season enhances tree growth in cold humid areas but not in dry areas *Nat. Ecol. Evol.* **6** 397–404
- Gauthier S, Bernier P, Kuuluvainen T, Shvidenko A Z and Schepaschenko D G 2015 Boreal forest health and global change *Science* **349** 819–22
- Grigor'ev A A, Devi N M, Kukarskikh V V, V'yukhin S O, Galimova A A, Moiseev P A and Fomin V V 2019 Structure and dynamics of tree stands at the upper timberline in the western part of the Putorana Plateau *Russ. J. Ecol.* **50** 311–22
- Gurskaya M A 2014 Temperature conditions of the formation of frost damages in conifer trees in the high latitudes of Western Siberia *Biol. Bull. Russ. Acad. Sci.* **41** 187–96
- Gurskaya M A 2021 Formation of frost damage in larch growing in the northern and southern forest–tundra subzones of Siberia: a comparative analysis *Russ. J. Ecol.* **52** 556–66
- Gurskaya M A, Lange J, Kukarskih V V, Buras A and Wilmking M 2021 Stationarity of climate-growth response is only marginally influenced by the soil moisture regime in Western Siberia *Dendrochronologia* **69** 125873
- Hansson A, Yang W H, Dargusch P and Shulmeister J 2023 Investigation of the relationship between treeline migration and changes in temperature and precipitation for the Northern Hemisphere and sub-regions *Curr. For. Rep.* **9** 72–100
- Hantemirov R M et al 2022 Current Siberian heating is unprecedented during the past seven millennia *Nat. Commun.* **13** 4968
- Hantemirov R M, Shiyatov S G, Gorlanova L A, Kukarskih V V, Yu Surkov A, Hamzin I R, Fonti P and Wacker L 2021 An 8768 year Yamal tree-ring chronology as a tool for paleoecological reconstructions *Russ. J. Ecol.* **52** 419–27
- Hellmann L et al 2016 Diverse growth trends and climate responses across Eurasia's boreal forest *Environ. Res. Lett.* **11** 074021
- Hughes M K, Vaganov E A, Shiyatov S, Touchan R and Funkhouser G 1999 Twentieth-century summer warmth in the northern Yakutia in a 600 year context *Holocene* **9** 629–34
- Jacoby G C, Lovelius N V, Shumilov O I, Raspopov O M, Karbainov J M and Frank D C 2000 Long-term temperature trends and tree growth in the Taymir region of Northern Siberia *Quat. Res.* **53** 312–8
- Kang J, Yang Z, Yu B, Ma Q, Jiang S, Shishov V V, Zhou P, Huang J-G and Ding X 2023 An earlier start of growing season can affect tree radial growth through regulating cumulative growth rate *Agr. For. Meteorol.* **342** 109738
- Kharuk V I, Petrov I A, Im S T, Golyukov A S, Dvinskaya M L, Shushpanov A S, Savchenko A P and Temerova V L 2023 Subarctic vegetation under the mixed warming and air pollution influence *Forests* **14** 615
- Kharuk V I, Ranson K J, Im S T, Oskorbin P A, Dvinskaya M L and Ovchinnikov D V 2006 Tree-line structure and dynamics at the northern limit of the larch forest: anabar Plateau, Siberia, Russia *Arct. Antarct. Alp. Res.* **45** 526–37
- Kirilyanov A V et al 2020 Ecological and conceptual consequences of Arctic pollution *Ecol. Lett.* **23** 1827–37
- Kirilyanov A V, Hagedorn F, Knorre A A, Fedotova E V, Vaganov E A, Naurzbaev M M, Moiseev P A and Rigling A 2012 20th century treeline advance and vegetation changes along an altitudinal transect in the Putorana Mountains, northern Siberia *Boreas* **41** 56–67
- Kirilyanov A V, Piermattei A, Kolář T, Rybníček M, Krusic P J, Nikolaev A N, Reinig F and Büntgen U 2018 Notes towards an optimal sampling strategy in dendroclimatology *Dendrochronologia* **52** 162–6
- Kirilyanov A V, Prokushkin A S and Tabakova M A 2013 Tree-ring growth of Gmelin larch under contrasting local conditions in the north of Central Siberia *Dendrochronologia* **31** 114–9
- Kirilyanov A V, Saurer M, Arzac A, Knorre A A, Prokushkin A S, Churakova (Sidorova) O V, Arosio T, Bechuk T, Siegwolf R and Büntgen U 2024 Thawing permafrost can mitigate warming-induced drought stress in boreal forest trees *Sci. Total Environ.* **912** 168858
- Kirilyanov A, Hughes H, Vaganov E, Schweingruber F and Silkin P 2003 The importance of early summer temperature and date of snow melt for tree growth in Siberian Subarctic *Trees* **17** 61–69
- Knorre A A, Kirilyanov A V, Prokushkin A S, Krusic P J and Büntgen U 2019 Tree ring-based reconstruction of the long-term influence of wildfires on permafrost active layer dynamics in Central Siberia *Sci. Total Environ.* **652** 314–9
- Knorre A A, Kirilyanov A V and Vaganov E A 2006 Climatically induced interannual variability in aboveground production in forest-tundra and northern taiga of central Siberia *Oecologia* **147** 86–95
- Kolmogorov A I, Kruse S, Nikolaev A N, Kirilyanov A V and Pestryakova L A 2023 Dendroclimatic studies of *Larix cajanderi* Mayr. in the Omoloy River Basin *Arct. Subarct. Nat. Resour.* **28** 584–94 in Russian
- Kruse S, Shevtsova I, Heim B, Pestryakova L A, Zakharov E S and Herzsich U 2023 Tundra conservation challenged by forest expansion in a complex mountainous treeline ecotone as revealed by spatially explicit tree aboveground biomass modeling *Arct. Antarct. Alp. Res.* **55** 2220208
- Kullman L 2007 Tree line population monitoring of *Pinus sylvestris* in the Swedish Scandes, 1973–2005: implications for tree line theory and climate change ecology *J. Ecol.* **95** 41–42
- Liang M, Sugimoto A, Tei S, Takano S, Morozumi T, Shingubara R, Murase J and Maximov T C 2023 Arctic plant responses to summer climates and flooding events: a study of carbon and nitrogen-related larch growth and ecosystem parameters in Northeastern Siberia *J. Geophys. Res.* **128** e2022JG007135
- Linderholm H W 2006 Growing season changes in the last century *Agr. For. Meteorol.* **137** 1–14
- Livensperger C, Steltzer H, Darrouzet-Nardi A, Sullivan P F, Wallenstein M and Weintraub M N 2016 Earlier snowmelt and warming lead to earlier but not necessarily more plant growth *AoB Plants* **8** lw021
- Lloyd A H, Bunn A G and Berner L 2011 A latitudinal gradient in tree growth response to climate warming in the Siberian taiga *Glob. Change Biol.* **17** 1935–45
- MacDonald G M, Case R A and Szeicz J M 1998 A 538 year record of climate and treeline dynamics from the lower Lena river

- region of northern Siberia, Russia *Arct. Alp. Res.* **30** 334–9
- Meko D M, Panyushkina I P, Agafonov L I and Edwards J A 2020 Impact of high flows of an Arctic river on ring widths of floodplain trees *Holocene* **30** 789–98
- Menzel A et al 2006 European phenological response to climate change matches the warming pattern *Glob. Change Biol.* **12** 1969–76
- Miesner T et al 2022b Tree data set from forest inventories in north-eastern Siberia PANGAEA (<https://doi.org/10.1594/PANGAEA.943547>)
- Miesner T, Herzsuh U, Pestryakova L A, Wiczorek M, Zakharov E S, Kolmogorov A I, Davydova P V and Kruse S 2022a Forest structure and individual tree inventories of northeastern Siberia along climatic gradients *Earth Syst. Sci. Data* **14** 5695–716
- Naurzbaev M M, Vaganov E A, Sidorova O V and Schweingrube F H 2002 Summer temperatures in eastern Taimyr inferred from a 2427 year late-Holocene tree-ring chronology and earlier floating series *Holocene* **12** 727–36
- Nikolaev A N, Fedorov P P and Desyatkin A R 2011 Effect of hydrothermal conditions of permafrost soil on radial growth of larch and pine in Central Yakutia *Contemp. Probl. Ecol.* **4** 140–9
- Ols C, Klesse S, Girardin M P, Evans M E K, DeRose R J and Trouet V 2023 Detrending climate data prior to climate–growth analyses in dendroecology: a common best practice? *Dendrochronologia* **79** 126094
- Paulsen J and Körner C 2014 A climate-based model to predict potential treeline position around the globe *Alp. Bot.* **124** 1–12
- Piao S et al 2015 Leaf onset in the northern hemisphere triggered by daytime temperature *Nat. Commun.* **6** 6911
- Previdi M, Smith K L and Polvani L M 2021 Arctic amplification of climate change: a review of underlying mechanisms *Environ. Res. Lett.* **16** 093003
- Rantanen M, Karpechko A Y, Lipponen A, Nordling K, Hyvärinen O, Ruosteenoja K, Vihma T and Laaksonen A 2022 The Arctic has warmed nearly four times faster than the globe since 1979 *Commun. Earth Environ.* **3** 168
- Rees W G, Hofgaard A, Boudreau S, Cairns D M, Harper K, Marnett S, Mathisen I, Swirad Z and Tutubalina O 2019 Is subarctic forest advance able to keep pace with climate change? *Glob. Change Biol.* **26** 3965–77
- Rinn F 2003 *TSAP-Win—Time Series Analysis and Presentation Dendrochronology and Related Applications* (Frank Rinn)
- Rinne K T, Saurer M, Kirilyanov A V, Bryukhanova M V, Prokushkin A S, Churakova (Sidorova) O V and Siegwolf R T W 2015a Examining the response of needles carbohydrates from Siberian larch trees to climate using compound-specific  $\delta^{13}\text{C}$  and concentration analyses *Plant Cell Environ.* **38** 2340–52
- Rinne K T, Saurer M, Kirilyanov A V, Loader N, Bryukhanova M V, Werner R and Siegwolf R T W 2015b The relationship between needle sugar carbon isotope ratios and tree rings of Siberian larch *Tree Physiol.* **35** 1192–205
- Schaefer K, Zhang T, Bruhwiler L and Barrett A P 2011 Amount and timing of permafrost carbon release in response to climate warming *Tellus B* **6** 165–80
- Schuur E A G et al 2015 Climate change and the permafrost carbon feedback *Nature* **520** 171–9
- Schweingruber F H 1996 *Tree Rings and Environment: Dendroecology* (Haupt) p 609
- Serreze M C and Barry R G 2011 Processes and impacts of Arctic amplification: a research synthesis *Glob. Planet. Change* **77** 85–96
- Shestakova T A, Gutiérrez E, Kirilyanov A V, Camarero J J, Génova M, Knorre A A, Linares J C, de Dios V R, Sánchez-Salguero R and Voltas J 2016 Forests synchronize their growth in contrasting Eurasian regions in response to climate warming *Proc. Natl Acad. Sci. USA* **113** 662–7
- Shishov V V, Arzac A, Popkova M I, Yang B, He M and Vaganov E A 2023 Experimental and theoretical analysis of tree-ring growth in cold climates *Boreal Forests in the Face of Climate Change (Advances in Global Change Research vol 74)*, ed M M Girona, H Morin, S Gauthier and Y Bergeron (Springer)
- Shishov V V, Tychkov I I, Popkova M I, Ilyin V A, Bryukhanova M V and Kirilyanov A V 2016 VS-oscilloscope: a new tool to parameterize tree radial growth based on climate conditions *Dendrochronologia* **39** 42–50
- Shiyatov S G 1993 *The Upper Timberline Dynamics during the Last 1100 Years in the Polar Ural Mountains. Oscillations of the Alpine and Polar Tree Limits in the Holocene* ed B Frenzel, B Eronen and M Glaser (Gustav Fischer Verlag) pp 195–203
- Shur Y L and Jorgenson M T 2007 Patterns of permafrost formation and degradation in relation to climate and ecosystems *Permafrost. Periglac. Process.* **18** 7e19
- St. George S 2014 An overview of tree-ring width records across the Northern Hemisphere *Quat. Sci. Rev.* **95** 132–50
- Stuenzi S M, Boike J, Gädeke A, Herzsuh U, Kruse S, Pestryakova L A, Westermann S and Langer M 2021 Sensitivity of ecosystem-protected permafrost under changing boreal forest structures *Environ. Res. Lett.* **16** 084045
- Tei S et al 2019a An extreme flood caused by a heavy snowfall over the indigirka river basin in Northeastern Siberia *Hydrol. Process.* **34** 522–37
- Tei S, Nagai S and Sugimoto A 2019b Effects of climate dataset type on tree-ring analysis: a case study for Siberian forests *Polar Sci.* **21** 136–45
- Tei S and Sugimoto A 2020 Excessive positive response of model-simulated land net primary production to climate changes over circumboreal forests *Plant-Environ. Int.* **1** 102–21
- Turetsky M R et al 2019 Permafrost collapse is accelerating carbon release *Nature* **569** 32–34
- Vaganov E A, Hughes M K, Kirilyanov A V, Schweingruber F H and Silkin P P 1999 Influence of snowfall and melt timing on tree growth in subarctic Eurasia *Nature* **400** 149–51
- Vaganov E A, Hughes M K and Shashkin A V 2006 *Growth Dynamics of Conifer Tree Rings: Images of past and Future Environments Ecological Studies* vol 183 (Springer Science & Business Media)
- Vaganov E A, Shiyatov S G and Mazepa V S 1996 *Dendroclimatic Research in the Ural-Siberian Subarctic Zone* (Nauka) p 246 in Russian
- Wiczorek M, Kruse S, Epp L S, Kolmogorov A, Nikolaev A N, Heinrich I, Jeltsch F, Pestryakova L A, Zibulski R and Herzsuh U 2017 Dissimilar responses of larch stands in northern Siberia to increasing temperatures—a field and simulation based study *Ecology* **98** 2343–55
- Yin G, Niu F, Lin Z, Luo J and Liu M 2017 Effects of local factors and climate on permafrost conditions and distribution in Beiluhe basin, Qinghai-Tibet Plateau, China *Sci. Total Environ.* **581** 472–85