

# Blue Carbon Potential in Germany: Status and Future Development

**Koplin, Julian**<sup>a, b \*</sup>

Julian.Koplin@rifs-potsdam.de

**Peter, Corina**<sup>a \* †</sup>

Corina.Peter@awi.de

**Bischof, Kai**<sup>c</sup>

**Böttcher, Michael Ernst**<sup>d,e,f</sup>

**Kuhn, Annegret**<sup>g</sup>

**Logemann, Ella**<sup>h</sup>

**Dolch, Tobias**<sup>a</sup>

**Henkel, Susann**<sup>a</sup>

**McCarthy, Daniel**<sup>i</sup>

**Mueller, Peter**<sup>j</sup>

**Morys, Claudia**<sup>k</sup>

**Pineda-Metz, Santiago E.A.**<sup>a</sup>

**Reents, Svenja**<sup>a</sup>

**Reusch, Thorsten B. H.**<sup>l</sup>

**Röschel, Lina**<sup>b</sup>

**Rupprecht, Franziska**<sup>h</sup>

**Stevenson, Angela**<sup>l</sup>

**Wiltshire, Karen**<sup>a, 1</sup>

**Zimmer, Martin**<sup>m, n</sup>

**Pogoda, Bernadette**<sup>a</sup>

\*Contributed equally as first authors

† Corresponding author

<sup>a</sup> Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research (AWI), D-27570 Bremerhaven, Germany

<sup>b</sup> Research Institute for Sustainability - Helmholtz Centre Potsdam (RIFS), D-14467 Potsdam, Germany

<sup>c</sup> Marine Botany, University of Bremen, D-28359 Bremen, Germany

<sup>d</sup> Geochemistry & Isotope Biogeochemistry, Leibniz IOW, D-18119 Warnemünde, Germany

<sup>e</sup> Marine Geochemistry, University of Greifswald, D-17489 Greifswald, Germany

<sup>f</sup> Interdisciplinary Faculty, University of Rostock, D-18059 Rostock, Germany

<sup>g</sup> Center for Ocean and Society, Kiel University, Christian-Albrechts-Universitaet zu Kiel, D-24118 Kiel, Germany

<sup>h</sup> Lower Saxony Wadden Sea National Park, D-26382 Wilhelmshaven, Germany

<sup>i</sup> Kiel University, Christian-Albrechts-Universitaet zu Kiel, D-24118 Kiel, Germany

<sup>j</sup> Institute for Landscape Ecology, University of Münster, D-48149 Münster, Germany

<sup>k</sup> Federal Agency for Nature Conservation (BfN), D-53179 Bonn, Germany

<sup>l</sup> GEOMAR - Helmholtz Centre for Ocean Research Kiel (GEOMAR), D-24148 Kiel, Germany

<sup>m</sup> Leibniz Centre for Tropical Marine Research (ZMT), D-28359 Bremen, Germany

<sup>n</sup> Faculty 2 Biology/Chemistry, University of Bremen, D-28359 Bremen, Germany

---

<sup>1</sup> Present address: Trinity College Dublin, The University of Dublin, Dublin 2, Ireland

## Abstract

Climate change and biodiversity loss are global challenges that need to be addressed through a combination of measures. However, political and societal action has not yet kept pace with the urgency of these challenges. Marine carbon sequestering habitats ("Blue Carbon habitats") are globally recognized for their role in climate change mitigation and for their co-benefits and ecosystem functions, e.g., as habitat providers. In Germany, research on the Blue Carbon potential of coastal and marine ecosystems has gained momentum in recent years. However, a synergistic approach with an inclusive decision-making process is crucial to ensure political action. Current challenges are considerable knowledge gaps and the limited accessibility and transferability of existing data. Funding of research projects at different administrative levels impacts coordination, output and visibility. Here, we present a general overview of existing knowledge and identified knowledge gaps in Blue Carbon research and focus on potential Blue Carbon ecosystems (BCEs) of the German coast. Furthermore, we identify windows of opportunity and provide actionable recommendations at the science-policy-society interface by examining the current framework for Blue Carbon in Germany. Based on this, ongoing research can be further prioritized and funded in order to simultaneously strengthen the political decision-making process. The results of this study, supported by the lessons learned from a case study on the German coast, recommend a two-pronged strategy to not only avoid additional release of already stored carbon through ecosystem conservation and sustainable governance and management, but also to increase net carbon storage through (re-)establishing BCEs.

## Keywords

Blue Carbon governance framework; seagrass; salt marsh; unvegetated soft sediment; kelp forest; biogenic reef

## 1. Introduction

Against the background of global climate change and the loss of biodiversity, there is an increased need to understand roles and contributions of coastal and marine ecosystems and habitats. Coastal ecosystems and their ecological function provide a range of valuable ecosystem services[1-3]. Coastal vegetated ecosystems (CVEs), like mangrove forests, salt marshes and seagrass meadows, can mitigate climate change processes by capturing and storing carbon as biomass and within sediments and soils over long time scales [3,4]. Carbon stored in marine ecosystems is referred to as Blue Carbon (BC) with regional CVEs being important coastal BC ecosystems (BCEs)[3,5]. BC as nature-based solution for climate mitigation aims to avoid or mitigate greenhouse gas emissions by conserving and restoring marine carbon- and biodiversity-rich habitats [3]. As such, BC is seen as carbon dioxide removal (CDR) which according to the latest assessment report of the Intergovernmental Panel on Climate Change are needed to offset residual emissions and achieve the internationally targeted net-zero scenario [4].

For northern Europe, salt marshes and seagrass meadows have been defined as such relevant coastal BCEs [3,6,7]. Research has primarily focused on these ecosystems, given their relatively high efficiency for carbon storage [8,9]. However, in recent years, several



studies have explored the BC potential of other marine systems [7,10-14], such as unvegetated soft sediments, macroalgae (e.g., kelp forests), biogenic reefs (e.g., mussel and oyster reefs), the role of marine organisms (e.g., fish) in these marine systems, effects of outwelling from BCEs for marine C storage, and coastal transition zones such as non-tidal peatlands along the Baltic Sea coast. Protecting these ecosystems from degradation and spatial loss will avoid the release of very large amounts of stored carbon from the associated sediments and soils [15]. Further, restoration of BCEs will increase carbon dioxide removal over time as well as increase local biodiversity in those habitats [16,17]. The important role of BCEs for climate mitigation is broadly recognized, whereas the feasibility of achieving quantifiable and secure negative emissions from restoration is still under debate [5]. In the light of fighting both crises at the same time, research into marine carbon sequestration and storage has received immediate scientific and political attention in the last decade [18,19]. Estimates of carbon storage indicate that more than 30 Gt of (organic) carbon are stored over  $1.85 \times 10^6$  km<sup>2</sup> of BCEs globally [18]. In contrast to other (technical) approaches to ocean-based carbon removal, where potential side effects to the ocean environment need to be closely examined first, conservation and restoration of BC on an ecological scale provides immediate positive effects [20,21]. However, in densely populated and heavily utilized areas, such as the German coasts, these ecological aspects are overshadowed by conflicts of interest and economic factors, making it crucial to consider the societal and political landscape.

BCEs are directly and indirectly integrated into the global environmental governance framework [22], however it is not specified how states ought to approach their restoration. On the EU level, a new regulation which introduces the necessary framework for ongoing habitat improvement, is the Nature Restoration Law (NRL[23])[24]. With aims to integrate biodiversity preservation and climate change mitigation (Art. 1 NRL), the NRL addresses the ongoing deterioration of nature and unfavorable habitat status within the Natura 2000 network identified by the EU Commission's State of Nature report [25]. It distinguishes itself by emphasizing targets, deadlines, and a specific focus on ecosystem restoration, surpassing conventional conservation legislation [26]. The steps and timetables proposed in the NRL provide a clear reference for a German strategy for the formulation of comprehensive nature restoration plans. Together with Germany's 2023 Federal Action Plan on Nature-Based Solutions for Climate and Biodiversity (ANK, Aktionsprogramm Natürlicher Klimaschutz, [27]), synergies between restoration, conservation and climate mitigation strategies in Germany could be strengthened. To move forward with such a national restoration plan [28,29], collecting and visualizing existing knowledge as well as prioritizing open questions is crucial.

This study aims to provide a first summary of BCEs specifically in Germany and within the national environmental policy regime. Here, we provide an overview of the function of carbon sequestration in national (coastal) BCEs as well as of potential (coastal) BCEs of the German North Sea and Baltic Sea. Further we summarize the state of scientific knowledge and related knowledge gaps that need to be addressed in order to I) integrate BC into a comprehensive national climate strategy and II) evaluate how existing BC potential can be enhanced through, e.g., management and restoration activities. We highlight windows of opportunities within the BC policy framework towards the conservation and restoration of BC environments on the sub-national, national and supranational level [30,31]. By outlining and acknowledging the range of open questions on this relatively new topic in Germany in the context of climate mitigation measures, our study helps to guide integrated and interdisciplinary approaches across ecosystems to understand and reveal underlying processes and their positive or negative effects on OM remineralization/carbon storage. A case study of salt marsh restoration on the

German Wadden Sea Coast exemplifies first steps for the practical implementation for enhancing BC in Germany.

## 2. General principles: Carbon capture and sequestration in marine ecosystems

CVEs are characterized by different compositions of vegetation and, thus, they manifest as different types of habitats and ecosystems such as mangrove forests, seagrass meadows or coastal marshes. Nevertheless, even though they are different, general BC drivers and factors apply to all (**Figure 1**).

- 1) CO<sub>2</sub> is captured via photosynthesis and incorporated into above-ground (leaves, plants) and below-ground plant material (roots, rhizomes) [37, 38]
- 2) Particulate organic carbon (POC) is trapped and captured by the vegetation, enabling a constant burial of organic carbon (C<sub>org</sub>) through vertical soil development, which is also influenced by varying rates of sea-level rise.
- 3) Low rates of microbial decay under reducing soil and sediment conditions [32-34].

The greater part of the vegetation in temperate ecosystems dies off seasonally (this does not apply to mangrove forests or most tropical seagrass beds) and is subsequently remineralized, or transported to adjacent ecosystems, or further into other marine and coastal areas [35-37]. However, a proportion of the dead biomass is buried *in situ* and is microbially degraded, releasing CO<sub>2</sub>, or is stored for decades, centuries or longer. Part of this (autochthonous) carbon is sequestered in deeper sediments or soil due to slower remineralization rates under anoxic conditions and high salinity levels compared to the degradation under oxic conditions. Remineralization leads to the formation and outwelling of alkalinity/dissolved inorganic carbon (DIC) [33] which may lead to the *in situ* formation of authigenic carbonates [38,39] and potentially influences the pH of coastal areas [40]. Above ground three dimensional structures of the vegetation (plant components like stem and leaves) reduce the flow velocity and increase sedimentation. Particles (including C<sub>org</sub> and C<sub>inorg</sub>) are removed from the water column by the vegetation and accumulate [41-43], forming and representing an additional allochthonous carbon pool.

In addition to active carbon capture through high primary production and reduced decomposition of organic matter, dense vegetation in CVEs prevents resuspension of deposited sediments and stabilizes the benthic environment. Coastal sediments or soils can contain several-thousand-year-old peat layers, thereby, upon degradation, linking the past with the present carbon cycle. This is an important aspect for the preservation of carbon storage in the sediments or soil as it hinders the release of carbon from the deposits of CVEs [4,44,45]. The amount of carbon deposited in CVEs depends on factors such as plant species composition, plant primary production, particle accumulation rate [32,46], and the distance to terrestrial carbon sources. The processes involved in carbon capture and burial operate over time-scales of hours to months [47,48]. The amount of captured carbon and longevity of storage depends on sediment type, dry bulk density, sedimentation rates, microbial activity, and on environmental conditions such as, e.g., temperature, oxygen conditions and current speed. All factors are subject to natural and anthropogenic variability, e.g., altering flooding frequency and changing salinity zones due to sea level rise in coastal areas, changes in local climate conditions, and events whirling up sediments like storms or bottom fishing activities

[9,49-52]. As carbon sequestration happens over timescales of centuries to millennia [6,47,53], the BC potential of marine ecosystems is mainly assessed within the (underlying) sediments or soils on which the described CVEs grow.

Fine-grained sediments (mud) become usually anoxic within a few mm depth and are usually characterized by higher  $C_{org}$  contents compared to coarse-grained sediments (e.g., sand) [54-56]. Under oxic conditions that prevail at the sediment surface, carbon is rapidly remineralized [33,37,57]. In coarse coastal sediments or soils, where advective transport dominates, oxygen may reach several cm into the sediment or soil. Muddy sediments are dominated by diffusive fluxes, enabling fast oxygen consumption. Further downward, the organic matter degradation (under anoxic conditions) continues at slower rates [34,58-60] using other electron acceptors such as nitrate ( $NO_3^-$ ), manganese (Mn) and iron (Fe) oxides or sulfate. In principle, high sedimentation rates “dilute” the organic matter content, but may also bury organic matter faster into those depths that are oxygen-depleted [58,61-64]. Therefore, higher sedimentation rates usually enhance carbon preservation [65].

Except for permeable tidal surface sediments during low tide, marine sediments and coastal sediments and soils are typically permanently water-saturated and low in dissolved oxygen compared to drained sediments or soils [66]. As a result, microbial decay of organic matter is slow due to the less efficient use of other electron acceptors, enabling long(er)-term carbon storage [32-34].

In general, the functioning of carbon sequestration in coastal and marine habitats is a complex cycle influenced by multiple factors. Some of these general principles are still being examined, e.g., the interaction between inorganic and organic carbon cycles and processes that could counteract carbon storage potential [67], the role of inorganic carbon [68], the relevance of non- $CO_2$  greenhouse gas emissions in the C-budget of BCEs [69,70] and overall the impact of climate change on (re-established) BCEs in the future.




Figure 1: Blue Carbon (BC) drivers and relevant factors. (A) Vegetation: three-dimensional structures such as stems and leaves reduce flow velocity, increasing sedimentation and particle accumulation. This process captures both autochthonous and allochthonous carbon, allowing a steady burial of  $C_{org}$ . The primary long-term storage in the context of BC occurs in deeper layers. (B)  $CO_2$  is captured via photosynthesis and incorporated into above-ground 3D-structures and below-ground structures (roots, rhizomes). Carbon is partly released back through organic matter decomposition. In the sediment/soil, where less exchange through, e.g., currents occur, these

processes are more locally connected than above-ground. (C) Part of the dead biomass is buried in situ and is degraded, with some of the carbon being locally stored in deeper sediments/soils due to reduced remineralisation rates under anoxic and/or highly saline conditions (autochthonous carbon) (D) A proportion of this carbon is remineralised or is transported to adjacent ecosystems or further into other marine and coastal areas, whereas the most recalcitrant material, is being transported without being degraded (transport of allochthonous carbon). (E) Fine-grained sediments, such as mud, tend to have higher organic carbon contents than coarser-grained sediments, such as sand or gravel. The ratio between autochthonous and allochthonous carbon differs depending on the sediment type. We would expect to see more allochthonous carbon in fine-grained sediments such as mud because material that has been transported between ecosystems (allochthonous carbon) is more likely to have reduced in size than material that has not been transported between ecosystems (autochthonous carbon).

### 3. Blue Carbon in German coastal and marine ecosystems

Factors such as the amount of carbon captured, the origin of this carbon (allochthonous vs. autochthonous), physical oceanography, sediment or soil characteristics and the remineralization rate influence the long-term carbon storage potential of BCEs [49,71-75]. Knowing these factors along with the extent covered by the ecosystems as well as their resilience with regard to, e.g., climate change allows an assessment of the carbon storage potential of BC in Germany. It further supports possible protection and optimization measures as required in a national restoration plan, i.e. the NRL or German Marine Strategy. Classical BCEs in Germany are represented by coastal marshes and seagrass meadows. However, other natural systems that contribute to BC storage are also discussed, both globally and in Europe, and are typically referred to as non-classical BC ecosystems [4]. In the following sections, the long-term carbon storage potential of non-vegetated marine sediments is discussed before the classical BCEs in Germany, in order to first illustrate the distribution of different underlying sediment types in the German coastal and marine ecosystems.

#### 3.1. Non-vegetated marine sediments

The long-term storage of organic carbon in marine non-vegetated sediments accounts for only about 5% of the total carbon inventory in BC systems, while the rest is recycled [76]. Nevertheless, partly due to their large spatial extent [56,77], shelf seas such as the North and Baltic Sea are considered to play an important role in the storage of carbon [78], including carbon that was taken up by marine organisms from the atmosphere [79,80]. The seabed of the German Exclusive Economic Zone (EEZ) covers 41,034 km<sup>2</sup> in the North Sea and 15,507 km<sup>2</sup> in the Baltic Sea [70]. One key ecosystem for BC research in the North Sea is the Wadden Sea, the largest tidal flat system in the world with largely undisturbed natural processes enabling significant local mud deposition [81]. The German sector of the Wadden Sea covers an area of about 4,030 km<sup>2</sup> (based on nautical chart/topographic data, data from NLPVW & LKN SH, 2015-2016).

The German North Sea is mainly characterized by shallow water depths (mean depth of 32 m, maximum of 71 m depth [82]), a wide tidal range, wind-induced turbulence and often high current velocities. Sands with low C<sub>org</sub> content dominate in the German Bight (**Figure 2**) [83,84]. The German North Sea sediments are often mixed and reworked due to the combination of the prevailing environmental conditions, leading to a fairly complete mixing of the water column during wind and storm conditions. The material is usually resuspended several times before final deposition, which means that the organic particles are subject to an enhanced oxic degradation. Therefore, carbon accumulation rates are close to zero over most areas [84], resulting in low C<sub>org</sub> contents (POC dry wt%) of 0.1% (Southern Bight) to 1.9% (south east of the Helgoland mud area) in sediments of the German North Sea [83]. Outside of the German part of the North Sea a first estimate [56] of POC storage in the top 10 cm of subtidal sediments of the north west European shelf is 0.48 (0.21 - 0.79) kgC m<sup>-2</sup>. The major

carbon deposition center of the North Sea (outside of Germany) is the Norwegian Trough (Skagerrak), accounting for about 87% of the total  $C_{org}$  accumulation in the total North Sea because of low current velocities and high sedimentation rates of fine sediments [56,83]. In the German North Sea, the area of particular interest for BC research for non-vegetated sediments is the Helgoland mud area, since this is a major offshore mud sink in the North Sea [81].

The German Baltic Sea is also very shallow (mean depth of 18 m, maximum of 47 m [82]), with mostly silty sediments (**Figure 2**) that are richer in organic material than sediments of the North Sea [77]. Outside of the German Baltic Sea average accumulation rates of  $22 \pm 10 \text{ g } C_{org} \text{ m}^{-2} \text{ yr}^{-1}$  were determined for the Bornholm Deep (south western Baltic Sea, Denmark) [80]. The  $C_{org}$  content (POC dry wt%) varies from 0.1% in shallow, sandy areas to 16% in deep, muddy, suboxic to anoxic areas (e.g., Gotland Basin, Sweden) [77]. For the top 10 cm of Baltic Sea sediments, the modelled  $C_{org}$  storage amounts to  $0.83 \pm 0.09 \text{ kgCm}^{-2}$  [85]. There are currently no studies on the  $C_{org}$  content specifically in the German EEZ. However, there are no anoxic areas present in the German EEZ which could potentially show  $C_{org}$  contents as high as in the Gotland Basin. Areas of particular interest to BC research for non-vegetated sediments in the German Baltic Sea are therefore the deep muddy basins (e.g., Kieler, Lübecker and Mecklenburger Bay, and Arkona Basin).

The potential of annual carbon sequestration and total carbon storage in marine sediments, as well as the origin of organic matter stored in the German EEZ, have not yet been determined in detail for neither the Baltic Sea nor the North Sea. Furthermore, long-term carbon storage in marine sediments is currently mainly considered as  $C_{org}$  pools. For a holistic picture,  $C_{inorg}$  pools and carbon fluxes need to be investigated too [77,80]. Additionally, microphytobenthos is an important stabilizer of marine sediments in shallow coastal waters. In shallow coastal areas, its productivity can be greater than the productivity of the water column. Thus, mud and sandflats may seem to be devoid of photosynthesizing plants, but due to the carbon fixation of these microalgae and bacteria this is not the case. Middelburg et al.[86] have shown that the carbon fixed by microphytobenthos can enter all heterotrophic components. Indeed, it has been shown repeatedly that microphytobenthos can play a central role in carbon flow in coastal sediments. In the context of its potential role in carbon sequestration, it is imperative that it is included in future intertidal and coastal carbon considerations.

To be able to estimate the BC potential of non-vegetated marine sediments, in the sense of the conservation of, e.g., carbon-rich habitats under national restoration plans in Germany, the following knowledge gaps need to be addressed: Absence of comprehensive data of the extent of relevant sediment types, carbon stocks ( $C_{org}$  and  $C_{inorg}$ ) and sequestration rates, non- $\text{CO}_2$  greenhouse-gas dynamics and factors influencing them. Several national projects are working on first estimates on the missing data (**see supplementary data S1**). This basis is crucial to plan and implement future management and conservation plans for BC stocks in marine sediments within the German EEZ as well as on the coasts through protection measures. Marine sediments are named in Annex II in the EU NRL and fall within the monitoring of the Habitats Directive [87] as well as of the Marine Strategy Framework Directive (MSFD).






Figure 2: Distribution and type of marine sediments for the German North Sea (left) and the German Baltic Sea (right). Left part of the Figure based on Laurer et al., 2014 [88]. Right part of the Figure based on BSH, 2016 [89].

### 3.2. Coastal marshes

Coastal marshes are highly productive ecosystems at the interface between land and sea and are characterized by distinctive flora and fauna. They include salt, brackish, and freshwater marshes, with habitats such as tidal flats, estuaries and shallow coastal waters generally considered adjacent to vegetated coastal marshes. All these habitats are relevant to BC, albeit studied to varying degrees, especially in Germany. Coastal marshes are studied on a broader scale than tidal flats simply because they store more carbon per unit area. Salt marshes are the most common type of coastal marshes along the German Wadden Sea on the North Sea coast and cover a small portion along the German Baltic Sea coast (**Figure 3**). For Wadden Sea marshes only, initial evidence suggests that in these habitats, allochthonous carbon accounts for the majority of the long-term sequestered carbon [41] with an input of approx.  $1.84 \text{ Mt yr}^{-1}$  and mud sedimentation rates up to a magnitude higher than in the basins of the Wadden Sea [81]. Autochthonous carbon contributes comparatively little to this pool [73,74,90]. Salt marshes are subject to regular flooding, which is a source of external soils and carbon, and their soils are often anaerobic and highly saline. The combination of these factors leads to a suppression of microbial organic matter decomposition, thus enabling carbon preservation and eventually long-term sequestration [32]. However, the influence of anthropogenic drainage systems by, e.g., the use of artificial ditches in salt marshes of the Wadden Sea, likely results in higher than normal drainage rates which, in turn, increase soil aeration and, thus, organic matter decomposition (and associated carbon release) in the upper soil layers [73,91].

The topsoil of saltmarsh soils contains relatively high densities of organic material, although this is fresh material rather than stored or sequestered carbon. The carbon density decreases with increasing soil depth until it eventually reaches a stable state and can be considered as effectively preserved and long-term sequestered; the depth at which this state is reached varies depending on site hydrology and redox conditions [74]. While a fraction of the topsoil carbon may eventually end up as long-term stored carbon in the subsoil, the carbon stored in the long term is primarily restricted to deeper soil layers [73]. There are several factors affecting the quantity and quality of organic matter throughout the soil column, some of them are: species composition [92], the flooding frequency [93], the distance of the habitat to allochthonous carbon sources, and the availability of electron acceptors [9]. Especially the latter two can be greatly influenced by use and management of the land resources. For the Wadden Sea, at least 50% of the salt marshes are used for livestock grazing, which can

strongly affect plant biomass production, soil microbial activity and carbon sequestration. Studies from the Wadden Sea region suggest a positive impact of livestock grazing on soil carbon sequestration in salt marshes [74,94,95].

Along the Wadden Sea coast, coastal marshes cover about 196.7 km<sup>2</sup>, along the Baltic Sea coast about 50.2 km<sup>2</sup> (monitoring data of the German federal states). Measurements for the 1 m-depth soil layer of coastal marshes along the German North Sea coast indicate an annual C<sub>org</sub> sequestration rate of 75.64 - 165.6 gC m<sup>-2</sup> yr<sup>-1</sup> with a carbon storage potential of 12.2 - 21.7 kgC m<sup>-2</sup> [73]. In contrast, for the German Baltic Sea coast, the carbon storage of a marsh colonized by reeds was estimated to vary between 1.76 – 88.6 kgC m<sup>-2</sup> for the 1 m-depth soil layer [96]. To the best of our knowledge, there are no further examples for comparing the measured carbon storage potential between similar coastal marsh types along the German coast. There are also no further examples for comparing different coastal marsh types within the same coastal basin [74]. It is noteworthy that during the preparation of our study, a first step in this direction was achieved, as a compilation of C<sub>org</sub> for salt marshes on a global scale was carried out [8].

In Germany and the EU, coastal marshes are already acknowledged as important and conserved habitat types, e.g., under the MSFD and the habitats directive (salt marshes: Directive 92/43/EEC habitat types 1310, 1320, 1330). In the EU NRL they are named as part of Coastal Wetlands under Annex I. To be able to estimate the BC potential of coastal marshes in Germany, also to be included in future national conservation and climate mitigation plans, the following knowledge gaps need to be addressed: absence of comprehensive and high resolution spatial and temporal data on carbon stocks, on sequestration and remineralization rates (OC and IC) as well as on non-CO<sub>2</sub> greenhouse-gas dynamics. Several national projects are working on first estimates on the missing data (see **supplementary data S1**). With a view on future management regulations, individual assessment of livestock influence on coastal marshes is needed. In relation to climate change effects on salt marshes in the future this includes, e.g., to evaluate promoting carbon storage versus preparing for rising sea levels.




Figure 3: Distribution map of salt marshes (Directive 92/43/EEC habitat types 1310, 1320 and 1330) along the German North the German North Sea and Baltic Sea coasts, maritime border: German EEZ (Sources: North Sea: Monitoring data NLWKN (2014-2017), monitoring data LKN SH (2021): FFH habitat types 1310, 1320, 1330; Baltic Sea: monitoring data LUNG MV (2005-2021), monitoring data LLUR SH (2012-2020): FFH LRT 1310, 1330), Figure from Koplin et al., 2022 [97].

### 3.3. Seagrass meadows

Seagrass meadows can form large, dense beds and, thus, create extensive habitats on a shallow, sedimentary seafloor from which many associated species benefit (e.g., food, shelter, settlement of sediment) [98-104]. They are highly productive, represent biodiversity hotspots and form an important cornerstone of the food web of the German seas [103,105,106]. The common seagrass *Zostera marina* L. is the dominant seagrass species in the Baltic Sea [107], whereas habitat requirements for the dwarf eelgrass *Z. noltii* Hornem differs and it dominates populations in the Wadden Sea of both Lower Saxony [108] and Schleswig-Holstein [109], the south-east and north-west sectors of the German North Sea, respectively. In the German Wadden Sea, seagrass meadows cover about 191 km<sup>2</sup> and their distribution is limited to intertidal and sheltered locations (**Figure 4**) [110]. In comparison, about 269 km<sup>2</sup> of the German Baltic Sea's sublittoral regions are covered by seagrass meadows, large and dense beds are found especially in sheltered bays (**Figure 4**) [107,111].

Both *Zostera* species structure starkly different meadows (e.g., density and canopy height), which can significantly influence the carbon storage potential [16,44,112,113]. The flow-reducing influence of the seagrass vegetation depends, among others, on the structural characteristics of the grass meadow (e.g., canopy height, and shoot density) [114,115]. Thus, the structural differences between *Z. marina* and *Z. noltii* result in clear variances, in terms of biomass per unit area between the Baltic and North Sea seagrass meadows both above and below ground, with biomass of Baltic seagrass meadows exceeding that of the populations found in the North Sea [111]. In terms of C<sub>org</sub> storage, there is currently no available data for the seagrass meadows of the North Sea, while there are datasets available for the Baltic Sea, e.g., a stock of approximately  $1.9 \pm 0.4 \text{ kgCm}^{-2}$  was calculated for the first 25 cm of the sediment layer [116]. From this stock, 12% corresponds to autochthonous sources, whereas 88% of the C<sub>org</sub> stock originates from allochthonous sources (phytoplankton and macroalgae). Relics of terrestrial peatland material deposited approximately 6,000 years BP during the last deglaciation, represent an unexpected and significant storage of C<sub>org</sub>. The Baltic Sea, in comparison to the North Sea, has comparatively finer sediments, less hydrodynamic energy (currents and waves), more efficient capture of allochthonous organic material through higher seagrass complexity, and the relics of terrestrial peatlands. These factors, in combination with the lower biomass per unit area of the North Sea suggest a generally lower storage potential per unit area for seagrass areas in the North Sea. However, the increased occurrence of seagrass meadows in the Wadden Sea through favoring natural spread or reintroduction as a nature-based solution will alter hydro-morphodynamic conditions in favor of sediment, and therefore potentially also carbon, accumulation [117].

In Germany, seagrass beds are monitored as one environmental parameter for the assessment of the ecological condition of coastal areas within the framework of the MSFD and EU WFD. In the EU NRL they are named as part of Group 1 (seagrass beds) under Annex II, being an important ecosystem also for the implementation of the future National Marine Strategy (NMS) of the German government. In order to be able to define substantial recovery and restoration plans with respect to their BC potential along the German coasts the following knowledge gaps need to be addressed: Lack of comprehensive data in terms of extent of seagrass meadows (specifically in the subtidal areas in the Baltic Sea), carbon stocks and sequestration rates, non-CO<sub>2</sub> greenhouse-gas dynamics and the local effect of seagrass restoration on BC sequestration potential. Interaction between inorganic and organic carbon within local carbon cycles could counteract carbon storage potential (e.g., calcification in ecosystems). With regard to changing environmental factors such as temperature and nutrient



loads also the impact of climate change effects on (re-established) seagrass meadows need to be included into future restoration plans. Several national projects are working on first estimates on the missing data (see supplementary S1) [97].



Figure 4: Distribution map of seagrass meadows along the German North Sea and Baltic Sea coasts, maritime border: German EEZ (Sources: North Sea: KÜFOG & Steuer, 2020 [108]; Dolch, 2020 (degree of coverage > 5%; [118]); Baltic Sea: Schubert et al., 2015 [111] (point data), monitoring data LUNG MV 2017 (all coverage degrees)); Figure from Koplin et al., 2022 [97].

### 3.4. Non-classical Blue Carbon ecosystems

The two ecosystems described in the previous sections (marshes and seagrass meadows) are referred to as classical BCEs. Some non-classical BCEs include intertidal and subtidal benthic habitats (e.g., biogenic reefs), mudflats, mesophotic habitats, and macroalgal forests [47], with the latter currently being of great interest in nature-based CDR research, also in Germany. In the German North Sea, relevant distributions of macroalgal forests in this context are mainly represented as kelp forests and found around the island of Helgoland [119,120]. Another possibly relevant macroalgae, *Fucus* spp., also grows in the intertidal zone at Helgoland and can be found along almost the entire sublittoral Baltic Sea coast [119,121]. Macroalgae fix large amounts of carbon during photosynthesis [13]. However, their distribution in Germany is predominantly limited to rocky coasts and shallow reefs, which excludes long-term carbon sequestration in the underlying sediments. Macroalgal detritus can be transported to adjacent deeper habitats or to surrounding habitats with long-term sequestration capacity, such as seagrass meadows or saltmarshes. While this makes it difficult to assess carbon-storage potential for macroalgal forests, the transport of macroalgal detritus suggests macroalgae to act as carbon donors [122-125]. Further difficulties in assessing carbon storage or transport provided by macroalgal forests are related to the uncertainties in calculating storage of macroalgal detritus in sediments, as these require the inclusion of detritus consumption by the detritivorous fauna, the associated remineralization and the local net primary production of the macroalgae [126], factors for which little or no data are available, as is the case for seagrass and salt marsh detritus.

While presumably of smaller relevance to BC than macroalgal forests, the BC potential of biogenic reefs (e.g., mussel and oyster beds), representing both organic and inorganic depots, is also a current controversially discussed topic [10,127-129]. While the formation of

calcareous shells releases CO<sub>2</sub> during the process of calcification, particulate inorganic carbon (PIC) can be buried and stored in the sediment for decades, centuries or millennia [76,130,131]. Further, the dissolution of this PIC causes an increase in alkalinity and therefore an uptake of CO<sub>2</sub>, possibly also enhancing the sequestration potential in habitats where shells are buried [132]. In addition, mussel beds can have indirect effects on sediment dynamics and consequently enhance carbon burial into the underlying sediment [133]. To understand the potential of biogenic reefs for the German seas, analyses of processes within existing reefs, but also the consideration of relevant adjacent processes and interactions, are needed [129,134]. Biogenic reefs are known to increase biodiversity and are therefore also named as a Directive 92/43/EEC habitat type (1170) as well as in Annex II of the EU NRL. Besides other already well-known ecosystem services supplied by biogenic reefs, they could play a role in long-term carbon storage [129]. Large knowledge gaps here are still remaining, mostly related to the regional influence of biogenic reefs on the carbon cycle and the surrounding ecosystem. Several national projects are working on first estimates and collecting missing data also on non-classical BC ecosystems in order to gain a holistic overview of potentials in the German seas (**see supplementary S1**).

#### 4. Integration of policy, society and research: the implementation of a Blue Carbon Restoration Plan (BCRP)

While BC is emerging as a topic in Germany's environmental policy landscape, global and regional biodiversity conservation and restoration targets for BCEs are still far from being met. Enhancement of BC in Germany requires comprehensive and integrated policy, in addition to scientific knowledge and evidence to inform decision-making and increased societal awareness and interest of Germany's general public in the enhancement of BC. This roadmap reflects on the interaction of policy, society and science and determines windows of opportunity for the comprehensive integration of BC into future decision-making in Germany.

Policy action on BC is critically dependent on existing regulations in Germany's complex multi-level governance structure [135]. The aim is to strengthen synergies between existing restoration, conservation, and climate mitigation strategies by implementing a Blue Carbon Restoration Plan (BCRP).

##### 4.1. Policy landscape: Windows of opportunity for Blue Carbon enhancement in Germany

Germany is a federal parliamentary republic comprising 16 states, each having its own constitution and being largely autonomous. While Germany's EEZ is governed at the national level, most CVEs are governed under the jurisdiction of Germany's five coastal states (see also "5. Case study"). The topic of BC touches upon a multitude of policy topics (e.g., climate, ocean, economy) and therefore the enhancement of BCEs is addressed across multiple entities on federal and national level, further challenging comprehensive governance (e.g., topical fragmentation or doubling of efforts). In addition, Germany, as a Member State of the EU adheres to its supranational environmental policy regime (e.g., European Green Deal) and aims to align its position with other EU Member States in intergovernmental negotiations within the UN System (e.g., nationally determined contributions). The different relevant levels of governance are outlined below, focusing on the national and sub-national political levels, as well as the EU level, on which we concentrate our study given the current relevance of the ANK and NRL.

509 *Sub-national / state level*

510 In order to enhance comprehensive management of marine issues across the five coastal  
511 federal states the Federal/State Working Group on the North Sea and the Baltic Sea (BLANO,  
512 Bund-Länder-Arbeitsgruppe-Nord-und-Ostsee) was established in 2012. BLANO comprises  
513 nine working groups and three expert bodies that focus on data collection, evaluation, and  
514 action related to marine topics and could also contribute significantly to the national  
515 implementation of a BCRP. The establishment of a working group or expert body to explicitly  
516 address climate-related issues, including BC, would help to promote cross-cutting policy  
517 integration. To achieve this, in addition to the ministries of environment, food and agriculture,  
518 digital and transport, the ministry of economy and climate action should be closely integrated  
519 into BLANO proceedings.

520 *National / federal level*

521 After federal elections in 2021, the German government put forth a coalition agreement that  
522 stated the aim of “enhancing the ocean’s natural CO<sub>2</sub> storage capacity through a targeted  
523 restoration programme (seagrass meadows, algae forests)”. In 2023, Germany published the  
524 ANK that recognizes the synergies between nature restoration and climate mitigation. ANK  
525 targets specific measures to be achieved by 2026, such as the evaluation of marine carbon  
526 inventories and the development of standardized measurement methods, with a total financing  
527 budget of four billion euros. The NMS is set to be published by mid-2025 and will potentially  
528 further address these efforts in a synergistic manner as well as multiple aspects of the ocean-  
529 climate-nexus in German waters. In addition to financing research projects to close the above-  
530 mentioned CVE-related knowledge gaps, the establishment of a national knowledge and data  
531 sharing platform to link science with policy is important. Providing policy makers with access  
532 to analyzed data on habitat conditions and monitoring status would enhance effective, science-  
533 based governance thereof. The ANK’s plans to standardize data collection efforts and scientific  
534 sampling must allow for national and international comparisons of data and habitats. They  
535 must be aligned with stakeholders from all five coastal states, as well as international cross-  
536 border cooperation (e.g., with Denmark or the Netherlands). Further, national restoration  
537 efforts should integrate carbon sequestration potential of relevant ecosystems and habitats  
538 while avoiding additional release of stored carbon by conservation measures [97]. The  
539 forthcoming NMS is an opportunity to recognize BCEs and marine ecosystems for their climate  
540 mitigation potential as well as for their ecological value. The ANK foresees the identification of  
541 marine areas with carbon-rich sediments and the development of a possible legal framework  
542 for the future designation of climate protection areas (CPAs) [136]. Finally, inclusion of all  
543 stakeholders, including the general public (see “4.2. Societal landscape”), at appropriate time  
544 scales will minimize conflicts, e.g., arising from area-use-competition.


545 The carbon sequestration potential of German BCEs is rather small in comparison to national  
546 emissions. Therefore, a BCRP should consider climate change mitigation and adaptation  
547 potential at the same time as importance for conserving and increasing biodiversity when  
548 determining which restoration measures to undertake (first). Germany could, for example, give  
549 priority for action in areas where salt marshes and seagrass meadows (as well as other  
550 potential BC habitats) have the greatest potential for carbon sequestration and simultaneous  
551 biodiversity conservation.

552 *Supra-national / EU level*


The EU Biodiversity Strategy for 2030 goal of protecting 30% of terrestrial and marine areas, including BCEs, by the end of the decade is translated into action through several environmental policies. Among those relevant for BC is the EU Habitats Directive, which requires Germany to designate special protection areas as part of the Natura 2000 network to achieve or maintain a 'favorable conservation status' and to monitor and report the status of protected species and habitats to the EU every six years. The German Federal Agency for Nature Conservation (BfN) prepares these reports, while the five federal coastal states develop management plans for each Natura 2000 site which are under their jurisdiction. The MSFD has the objective to achieve or maintain 'good environmental status' in Europe's seas based on 11 indicators, complementing the EU Water Framework Directive (WFD) of inland waters. The German Federal Environment Agency (UBA) is primarily responsible for implementation of the MSFD within the EEZ, while the various ministries and authorities of the five federal coastal states oversee implementation within the 12-mile coastal zone. The new NRL sets multiple binding targets and obligations for listed CVEs (e.g., salt marshes), such as restoring at least 30% of these ecosystems to good ecological condition (**see Figure 5**). The German ANK would transpose the NRL and is put into practice by the Federal Ministry for the Environment, Nature Conservation, Nuclear Safety and Consumer Protection (BMUV). Although Germany has decades of accumulated knowledge to support monitoring efforts [137], coordinating different stakeholders for data collection and assessment is challenging. For comprehensive BC monitoring and management of BCEs in Germany, e.g., under EU legislation, it is important to coordinate and streamline efforts across respective ministries and policy levels. BC restoration and management should encompass a long-term perspective supporting EU and global biodiversity, conservation and restoration goals. Measures need to be tailored to habitat properties including vulnerability to climate change to avoid carbon storage losses over time. Both, ANK and NMS initiatives should be tied to a progress chart for the implementation of the proposed legislation and respective measures. The implementation schedule for the NRL on BCEs (**Figure 5**) became legally binding for all EU countries since the law is passed. These (theoretical) timelines shall serve as a guide for national implementation. Under the NRL, states will have concrete milestones for restoring ecosystems to good ecological condition, while retaining flexibility in the means to achieve them. However, appropriate frameworks should be established in the German National Marine Strategy (NMS) for the development, implementation and updating of restoration plans.

Management or restoration plans should give a strategic overview of the estimated carbon sequestration potential of each measure it plans to implement, including "the estimated co-benefits or climate change mitigation and land degradation neutrality associated with the restoration measures over time, as well as wider socio-economic benefits" (Art. 12(2) (j) NRL).

**a** Blue Carbon Habitats and the implementation timeline for the proposed Nature Restoration Law (NRL):  
*Germany would have to restore salt-marsh habitats that fall under Annex I, Group 1 (Wetlands)*



**b** Blue Carbon Habitats and the implementation timeline for the proposed Nature Restoration Law (NRL):  
*Germany would have to restore seagrass habitats that fall under Annex 2, Group 1 (Seagrass Beds) and marine sediments that fall under Annex 2, Group 7 (Soft Sediments)*



**Figure 5: Theoretical timelines of the implementation of the EU Nature Restoration law in Germany with regard to the coastal Blue Carbon ecosystem types salt-marshes (a) and seagrass meadows and marine soft sediments (b) (Status as of March 2024).**

## 4.2. Societal landscape: Potential barriers, knowledge integration and public participation

Germany's BCRP must carefully balance the ecological, economic, and social functions of ecosystems, as well as their contribution to the sustainable development of the regions and communities concerned (Art. 14(16)(b) NRL). Land-based restoration efforts often face public skepticism, as exemplified by debates over peatland restoration versus agriculture [138]. In coastal areas, e.g., seagrass meadow restoration must contend with eutrophication, particularly in the Baltic Sea, which requires governance measures to reduce nutrient inputs. The governance of eutrophication is a highly contested policy area in Germany, at least with regard to nutrient inputs from the agricultural sector. In addition, marine restoration measures are facing increasing multiple uses, if not outright "industrialization" [139] of marine space by offshore wind farms and other energy infrastructure, military uses, tourism, as well as fishing. The provision of opportunities for public consultation and participation is widely recommended (for example, Clark 1994 [140] see also NRL: Art. 14(20) NRL), and indeed various empirical examples show that under certain contextual conditions, public consultation and participation promote political trust, acceptance and legitimacy of environmental decision-making [141]. A broader public participation with the BCRP could be recommended, e.g., through BLANO-working groups or expert groups that have been limited to researchers until now. However, this is no guarantee for increasing public acceptance, as shown by the recent consultation process in Schleswig-Holstein on the idea of a marine national park [142]. More emphasis on (long-term) awareness and knowledge building seems necessary to complement public participation processes [143]. In addition, different governance approaches and policies may need to be applied to the wider public as opposed to the communities or social groups directly affected by restoration activities. For instance, the commonly used argument of socio-economic benefits of restoration [144] does not apply equally to all societal groups, as socio-economic benefits are unequally distributed. Concrete compensation for economic losses, as well as benefits for the provision of ecosystem services to social groups such as farmers, should be discussed openly and with the involvement of these affected groups.

Finally, transdisciplinary research approaches - as currently initiated e.g. by the inter- and transdisciplinary German research missions CDRmare and SustainMare of the German Marine Research Alliance (DAM) - play a role by promoting (stakeholder) dialogues across disciplines for the co-creation and awareness of relevant knowledge and for triggering social learning processes. Successful formats of knowledge integration within different policy institutions, such as HELCOM, OSPAR or BLANO, will facilitate the identification and implementation of best practices.

## 4.3. Applied BC research: Scientific support and approaches for restoration efforts

636 Researchers can be key actors in providing critical assessments of governance integration  
637 efforts and of implementation barriers for a BCRP. Research will close knowledge gaps and  
638 optimize as well as standardize BC monitoring (see also “3. *Blue Carbon in German coastal*  
639 *and marine ecosystems*”). Restoration plans also need to be accompanied by research in order  
640 to measure their effects and, if necessary, to adopt changes. Within a BCRP, both active and  
641 passive approaches are possible. According to the NRL, active approaches for salt marsh  
642 restoration may include removing longitudinal and lateral barriers (such as dikes and dams)  
643 and for restoring seagrass meadows actively stabilizing the seabed, reducing and where  
644 possible eliminating pressure, or active propagation and planting. Passive measures to favor  
645 the natural spread may include reducing stressors and allowing ecosystems to develop their  
646 own natural dynamics, for example through the abandonment of harvesting and the promotion  
647 of wilderness.

648  
649 As an example of a practical implementation of these restoration approaches, we present a  
650 national case study highlighting a key effort focusing on salt marsh restoration in the Lower  
651 Saxony Wadden Sea National Park. The study summarizes the ecological, socio-cultural and  
652 coastal protection lessons learned from 30 years of dedicated restoration efforts in the German  
653 National Park. In addition, valuable knowledge for policy makers and practitioners worldwide  
654 is provided.

## 655 5. National case study: Salt marsh restoration in the Lower 656 Saxony Wadden Sea National Park: A chance to enhance 657 the Blue Carbon potential of anthropogenically modified salt 658 marshes

### 659 5.1. Policy background

660  
661 Most salt marshes in Germany are part of the Wadden Sea National Park located along the  
662 North Sea Coast of the federal states Lower Saxony, Schleswig-Holstein and Hamburg. In  
663 accordance with Federal Nature Conservation Act (BNatSchG § 24) the primary goal of the  
664 Wadden Sea National Park is to protect and enable the undisturbed course of natural  
665 ecosystem processes wherever possible. With respect to the Blue Carbon potential of salt  
666 marshes this means that the Wadden Sea National Park aims at preserving or restoring the  
667 *natural* C-sequestration function of salt marshes, resulting from their characteristic  
668 hydrological, geomorphological and biochemical processes.

### 672 5.2. Assessment of the state of salt marshes

673 Germany has monitored salt marsh habitats in successive six-year reporting periods, most  
674 recently from 2013 to 2018, as part of its reporting obligations under the Habitats Directive  
675 (**Table 1**). Germany has a general obligation to maintain or restore the favorable conservation  
676 status of designated salt marsh habitats under the Habitats Directive. Under the EU NRL,  
677 Germany will have specific timeframes to implement restoration measures to achieve this  
678 objective and to ensure that each habitat type improves to a state where 90 percent of the total  
679 area of that habitat type is in good condition. In practice, therefore, any German nature  
680 restoration plan will need to include restoration measures to maintain the status of salt marshes  
681 that are already in good condition, as well as to improve those salt marshes that are in poor  
682 (or unknown) condition.

Table 1: Overview of total area and respective environmental condition status of coastal marsh habitats in Germany according to the Habitats Directive (Habitats Directive, Article 17 Report, 2013-2018, Coastal Habitats, Germany, All Bioregions).

Habitat	Est. Total Area (km <sup>2</sup> )	Est. area in Good Condition	% Good Condition	Est. area in Poor Condition (km <sup>2</sup> )	Est. area in Unknown Condition (km <sup>2</sup> )
1310 - Salicornia and other annuals colonizing mud and sand	27.51	23.85	86.68%	3.66	0.62
1320 - Spartina swards ( <i>Spartinion maritimae</i> )	23.31	20.3	87.09%	3.00	0
1330 - Atlantic salt meadows ( <i>Glaucopuccinellietalia maritimae</i> )	262.88	168.99	64.28%	66.60	29.76

### 5.3. Principles of salt marsh management and restoration

Many mainland salt marshes in the Wadden Sea, including those in the Lower Saxony Wadden Sea National Park ("National Park" from now on), have been anthropogenically altered or modified in the past. The primary goal for managing the National Park's salt marshes today is to protect and enhance characteristic habitat processes, such as tidal flooding, sediment deposition, erosion, and succession. However, past human impacts still inhibit natural salt marsh development in many areas. The National Park Authority in Lower Saxony (NPA-LS) actively addresses this challenge through salt marsh restoration, involving singular interventions into the ecosystem. In the long run, such interventions reduce anthropogenic influence on salt marsh processes and habitat properties (**Figure 6 B**). Simultaneously, aligning salt marsh development with the National Park's goals provides an opportunity to potentially enhance their BC potential.






Figure 6: Salt marsh management and restoration in the National Park: A) typical mainland saltmarsh situation with uniform bed-ditch-structures. B) principal of salt marsh restoration with the goal of reducing cultivation impact on the habitats (© Linders). C) restored polder after dike removal D) Former clay pit, where marsh creeks established after elimination of bed-ditch structures.

The following types of salt marsh restoration methods in the National Park can be a blueprint for approaches within a potential BCRP:

1) **Opening or removing summer dikes** facilitates year-round tidal flooding (**Figure 6 C**), enabling the re-establishment of salt marsh vegetation in former summer polders [145]. Regular flooding reduces soil oxygen, slowing microbial decay of organic matter. Increased inundation with saline water suppresses methane production through sulfate input. The sulfate availability shifts soil microbial communities and methanogens are mostly outcompeted [146]. Furthermore, soil deposition increases marsh accretion and introduces allochthonous carbon from marine sources [73,147].

2) **Topsoil removal** lowers the soil surface elevation relative to the mean high tide and thus decreases the impact of artificial structures such as field drains. Consequently, formation of relief and vegetation zonation again align with hydrodynamic conditions and soil deposition patterns [145]. This measure resets the succession in favor of the pioneer/low marsh vegetation. The BC potential of the newly forming salt marsh is expected to exceed that of the anthropogenically drained marsh in the long-term, however further scientific studies are needed to clarify these processes.

3) **Restoring the natural hydrology** in the salt marsh involves deactivating or reducing the artificial drainage system. Unlike natural salt marshes, anthropogenically modified marshes have an oversized drainage system [148]. During the restoration process, field and collector drains are filled or blocked to extend flooding duration [145]. This, in turn, boosts waterlogging in soils, reducing microbial remineralization and promoting a stable state of carbon, and therefore increased storage, in salt marsh soils [74].

#### 5.4. Lessons learned from 30 years of salt marsh restoration to prioritize where to take action first

Since the National Park's establishment in 1986, 17 salt marsh restoration projects covering approximately 1000 ha have been completed [145]. Drawing on over 30 years of experience, the NPA-LS observed successful salt marsh re-establishment following interventions, such as summer dike removal (**Figure 7**). This suggests a potential increase in their BC potential over time.




Figure 7: Habitat development of salt marsh restoration sites prior to and after summer-dike opening. a) Hauener Hooge (monitoring period: 22 years), b) Langwarder Groden (monitoring period: 5 years).

Summer dike openings represent a highly effective restoration measure with minimal operational impact and a significant positive effect on BC potential [149]. This approach improves ecosystem quality without causing remineralization or carbon stocks due to restoration activities. Besides re-establishing a carbon sink, it is likely to reduce CO<sub>2</sub> and CH<sub>4</sub> emissions in formerly disturbed polders [150].

However, restored polders often exhibit a low proportion of mudflat and pioneer zones due to unchanged elevation. To address this, combining summer dike openings with topsoil removal can be beneficial [145]. The latter eliminates artificial drainage structures and initially leads to the development of mudflats and pioneer vegetation in the impact area, followed by lower salt marsh vegetation, within 3-5 years [145]. Old clay pits, where marsh soil has been excavated for reinforcement of dikes, show the establishment of natural marsh creeks and relief and can serve as a proxy to assess potential long-term development (> 30 years) of restoration sites with topsoil removal (**Figure 6 D**). These old clay pits are currently sampled for predicting the long-term effect of topsoil removal on the carbon storage.

When considering topsoil removal as a method for enhancing BC potential, it is crucial to include the initial and local conditions as a baseline for carbon calculations. E.g., initial carbon remineralization due to topsoil removal might be smaller when applied in polders where grasslands are established than when applied in tidally unrestricted salt marshes. To understand this impact on the net carbon balance in both tidally restricted (e.g., summer polders) and tidally unrestricted marshes, scientific studies are being conducted in collaboration with universities and the NPA-LS.

For rewetting areas that are permanently impacted by former cultivation, thorough planning and construction work is essential for a successful deactivation or reduction of the artificial drainage structures. In the Norderney Ostheller restoration site, shifts in plant communities 12-years post restoration indicate successful rewetting. Ongoing investigations will identify the impact of such measures on the BC potential.

The initial conditions of a restoration site significantly influence its later success of restoration. Key factors, as identified by the NPA-LS, include site morphology, drainage systems, and disturbance duration. While studies on BC in Wadden Sea salt marshes are limited and exclude restoration, embankments, such as summer dike openings are relatively well described and consistently show a significant impact on the BC potential. However, measures, such as reducing artificial drainage or topsoil removal, impacting carbon sequestration, are infrequently studied and involve high uncertainty on the long-term influence on carbon storage. The NPA-LS is actively addressing these knowledge gaps through collaborations with academic partners investigating each restoration measure's impact on Wadden Sea salt marshes. Recognizing the potential symbiosis between promoting habitat quality and enhancing BC, the NPA-LS aims to invest and promote both simultaneously.

## 5.5. Societal Acceptance: potential barriers, knowledge integration and public participation

Socio-cultural conflicts with local stakeholders pose a consistent challenge. The mainland salt marshes have deep cultural significance as man-made landscapes, shaped by past generations, creating an emotionally charged conflict when surrendering reclaimed land sites to natural tidal dynamics. Involving local stakeholders is a crucial step in the planning process. However, salt marsh restoration also offers great opportunities for tourism and environmental education, creating a win-win situation for the environment and the economy [151]. For instance, the 150 ha restoration site "Langwarder Groden" (summer dike opening, topsoil removal) includes a comprehensive nature experience concept, attracting approximately 50.000 visitors annually and gaining full acceptance from local communities despite initial resistance before and during construction.

Reconciliation of salt marsh restoration projects with coastal protection issues is a crucial matter. Salt marshes contribute to coastal protection by promoting wave attenuation and vertical accretion of the dike foreland through sediment trapping and deposition [1,152]. However, doubts arise among dike managers and associations regarding interference of natural, highly dynamic salt marsh processes such as succession and sedimentary processes with dike safety. Salt marsh restoration thus requires a careful and coordinated strategy, with planning and construction tailored to the local site conditions.

## 6. Outlook

The robust support for BC research is exemplified by numerous newly funded projects worldwide. In Germany, it is explicitly included in the coalition agreement, in the ANK and in the designated NMS. Collectively, these initiatives underline (and should result in) a strong commitment to fostering restoration as an integral part of overarching climate and marine conservation strategies. BC ecosystems such as CVEs but also non-classical but potential BC ecosystems, e.g., biogenic reefs provide crucial ecosystem functions and services as biodiversity hotspots and key habitats for a multitude of other organisms [153-155]. Conservation and restoration of such ecosystems are powerful tools to fight climate change and biodiversity loss. Both are equally important and critical for achieving climate mitigation goals and halting, preventing, and reversing the continuous decrease of biodiversity. These synergies are a chance which must not be underestimated. Seizing this opportunity well can even mean to co-use intact CVEs as living shorelines to improve coastal protection. NbS

combined with technical solutions such as dikes are flexible and able to adapt to changing climate conditions [117,156,157].

While Germany has recently published a promising “Federal Action Plan on Nature-based Solutions for Climate and Biodiversity” (ANK; 2023),[27]) it is common sense that climate change mitigation strategies cannot rely on NbS alone: (1) decarbonization must be the priority, (2) conservation of current carbon stocks (regional coastal and marine carbon stocks) and potentials must be explored and measured, (3) relevant CVEs must be expanded and managed to increase CO<sub>2</sub> sequestration potential, as shown in the national case study, while simultaneously supporting these biodiversity hotspots. Successful and sustainable mitigation of climate change is complex and multi-layered. A multi-pronged approach is needed to reach net carbon emissions targets and climate mitigation goals. (Potential) BCEs are not the key to achieving net zero, but they are key ecosystems for enhancing carbon storage processes and biodiversity, as well as providing various co-benefits. They are crucial to increasing the resilience of marine habitats and ecosystems to future changes. Germany is moving in the right direction to address the interlinked challenges but needs a temporally and spatially binding framework that a BCRP may provide.

## SUPPLEMENTARY MATERIAL

- S1: Table of ongoing (Blue Carbon related) projects in Germany

## DATA STATEMENT

For this publication no other data other than stated and cited in the manuscript was acquired.

## FUNDING

This work was supported by the Federal Agency for Nature Conservation (BfN) with funds from the Federal Ministry for the Environment, Nature Conservation, Nuclear Safety and Consumer Protection (BMUV) [Project DEFINE I, grant number FKZ: 3521831000, 2022, and DEFINE II grant number FKZ: 3522830300, 2022 - 2025].

## ACKNOWLEDGEMENTS

We would like to thank all experts and various project members who helped to bring together this valuable knowledge for their support and valuable feedback.

## 848 REFERENCES

- 849
- 850 1. Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C., and Silliman, B.R.
- 851 (2011). The value of estuarine and coastal ecosystem services. *Ecological*
- 852 *Monographs* 81, 169-193. 10.1890/10-1510.1.
- 853 2. Liqueste, C., Piroddi, C., Drakou, E.G., Gurney, L., Katsanevakis, S., Charef, A., and
- 854 Egoh, B. (2013). Current Status and Future Prospects for the Assessment of Marine
- 855 and Coastal Ecosystem Services: A Systematic Review. *PLOS ONE* 8, e67737.
- 856 10.1371/journal.pone.0067737.
- 857 3. Nellemann, C., Corcoran, E., Duarte, C., Valdes, L., Young, C., Fonseca, L., and
- 858 Grimsditch, G. (2009). Blue Carbon. A Rapid Response Assessment. ( United Nations
- 859 Environment Programme, GRID-Arendal).
- 860 4. IPCC (2022). Climate Change 2022: Impacts, Adaptation, and Vulnerability.
- 861 Contribution of Working Group II to the Sixth Assessment Report of the
- 862 Intergovernmental Panel on Climate Change.
- 863 5. Williamson, P., and Gattuso, J.-P. (2022). Carbon Removal Using Coastal Blue Carbon
- 864 Ecosystems Is Uncertain and Unreliable, With Questionable Climatic Cost-
- 865 Effectiveness. *Frontiers in Climate* 4. 10.3389/fclim.2022.853666.
- 866 6. Duarte, C.M., Middelburg, J.J., and Caraco, N. (2005). Major role of marine vegetation
- 867 on the oceanic carbon cycle. *Biogeosciences* 2, 1-8. 10.5194/bg-2-1-2005.
- 868 7. Lovelock, C.E., and Duarte, C.M. (2019). Dimensions of Blue Carbon and emerging
- 869 perspectives. *Biology Letters* 15, 20180781. doi:10.1098/rsbl.2018.0781.
- 870 8. Maxwell, T.L., Rovai, A.S., Adame, M.F., Adams, J.B., Álvarez-Rogel, J., Austin,
- 871 W.E.N., Beasy, K., Boscutti, F., Böttcher, M.E., Bouma, T.J., et al. (2023). Global
- 872 dataset of soil organic carbon in tidal marshes. *Scientific Data* 10, 797.
- 873 10.1038/s41597-023-02633-x.
- 874 9. Pollmann, T., Böttcher, M.E., and Giani, L. (2021). Young soils of a temperate barrier
- 875 island under the impact of formation and resetting by tides and wind. *CATENA* 202,
- 876 105275. 10.1016/j.catena.2021.105275.
- 877 10. Filgueira, R., Strohmeier, T., and Strand, Ø. (2019). Regulating services of bivalve
- 878 molluscs in the context of the carbon cycle and implications for ecosystem valuation.
- 879 In *Goods and services of marine bivalves*, A. Smaal, Ferreira, J., Grant, J., Petersen,
- 880 J., Strand, Ø., ed. (Springer, Cham), pp. 231-251. 10.1007/978-3-319-96776-9\_12.
- 881 11. HELCOM (2021). Report of the HELCOM Workshop on Blue Carbon Potential in the
- 882 Baltic Sea region. Baltic Marine Environment Protection Commission, ed. pp. 12.
- 883 12. Jurasinski, G., Janssen, M., Voss, M., Böttcher, M.E., Brede, M., Burchard, H., Forster,
- 884 S., Gosch, L., Gräwe, U., Gründling-Pfaff, S., et al. (2018). Understanding the Coastal
- 885 Ecocline: Assessing Sea–Land Interactions at Non-tidal, Low-Lying Coasts Through
- 886 Interdisciplinary Research. *Frontiers in Marine Science* 5. 10.3389/fmars.2018.00342.
- 887 13. Krause-Jensen, D., Lavery, P., Serrano, O., Marbà, N., Masque, P., and Duarte, C.M.
- 888 (2018). Sequestration of macroalgal carbon: the elephant in the Blue Carbon room.
- 889 *Biology Letters* 14, 20180236. 10.1098/rsbl.2018.0236.
- 890 14. Macreadie, P.I., Anton, A., Raven, J.A., Beaumont, N., Connolly, R.M., Friess, D.A.,
- 891 Kelleway, J.J., Kennedy, H., Kuwae, T., Lavery, P.S., et al. (2019). The future of Blue
- 892 Carbon science. *Nature Communications* 10, 3998. 10.1038/s41467-019-11693-w.
- 893 15. Pendleton, L., Donato, D.C., Murray, B.C., Crooks, S., Jenkins, W.A., Sifleet, S., Craft,
- 894 C., Fourqurean, J.W., Kauffman, J.B., Marbà, N., et al. (2012). Estimating Global “Blue
- 895 Carbon” Emissions from Conversion and Degradation of Vegetated Coastal
- 896 Ecosystems. *PLOS ONE* 7, e43542. 10.1371/journal.pone.0043542.
- 897 16. Greiner, J.T., McGlathery, K.J., Gunnell, J., and McKee, B.A. (2013). Seagrass
- 898 Restoration Enhances “Blue Carbon” Sequestration in Coastal Waters. *PLOS ONE* 8,
- 899 e72469. 10.1371/journal.pone.0072469.
- 900 17. Frigstad, H., Gundersen, H., Andersen, G.S., Borgersen, G., Kvile, K.Ø., Krause-
- 901 Jensen, D., Bostrom, C., Bekkby, T., Anglès d'Auriac, M., and Ruus, A. (2021). Blue
- 902 Carbon–climate adaptation, CO<sub>2</sub> uptake and sequestration of carbon in Nordic blue



- forests: Results from the Nordic Blue Carbon Project. Nordic Council of Ministers. 9289368608. <https://pub.norden.org/temanord2020-541/#50843>.
18. Macreadie, P.I., Costa, M.D.P., Atwood, T.B., Friess, D.A., Kelleway, J.J., Kennedy, H., Lovelock, C.E., Serrano, O., and Duarte, C.M. (2021). Blue carbon as a natural climate solution. *Nature Reviews Earth & Environment* 2, 826-839. 10.1038/s43017-021-00224-1.
19. Merk, C., Grunau, J., Riekhof, M.-C., and Rickels, W. (2022). The need for local governance of global commons: The example of blue carbon ecosystems. *Ecological Economics* 201, 107581. 10.1016/j.ecolecon.2022.107581.
20. Quevedo, J.M.D., Uchiyama, Y., and Kohsaka, R. (2023). Progress of blue carbon research: 12 years of global trends based on content analysis of peer-reviewed and 'gray literature' documents. *Ocean & Coastal Management* 236, 106495. 10.1016/j.ocecoaman.2023.106495.
21. Gattuso, J.-P., Williamson, P., Duarte, C.M., and Magnan, A.K. (2021). The Potential for Ocean-Based Climate Action: Negative Emissions Technologies and Beyond. *Frontiers in Climate* 2. 10.3389/fclim.2020.575716.
22. Röschel, L., and Neumann, B. (2023). Ocean-based negative emissions technologies: a governance framework review. *Frontiers in Marine Science* 10. 10.3389/fmars.2023.995130.
23. Parliament, E. (2024). European Parliament legislative resolution of 27 February 2024 on the proposal for a regulation of the European Parliament and of the Council on nature restoration (COM(2022)0304 – C9-0208/2022 – 2022/0195(COD)). [https://www.europarl.europa.eu/doceo/document/TA-9-2024-0089\\_EN.pdf](https://www.europarl.europa.eu/doceo/document/TA-9-2024-0089_EN.pdf).
24. Hering, D., Schürings, C., Wenskus, F., Blackstock, K., Borja, A., Birk, S., Bullock, C., Carvalho, L., Dagher-Kharrat, M.B., Lakner, S., et al. (2023). Securing success for the Nature Restoration Law. *Science* 382, 1248-1250. doi:10.1126/science.adk1658.
25. (EEA), E.E.A. (2020). State of nature in the EU: Results from reporting under the nature directives 2013-2018. EEA Report 10/2020. 10.2800/705440.
26. Hoek, N. (2022). A Critical Analysis of the Proposed EU Regulation on Nature Restoration: Have the Problems Been Resolved? *European Energy and Environmental Law Review*, 320-333.
27. BMUV (2023). Federal Action Plan on Nature-based Solutions for Climate and Biodiversity.
28. Mengis, N., Paul, A., and Fernández-Méndez, M. (2023). Counting (on) blue carbon—Challenges and ways forward for carbon accounting of ecosystem-based carbon removal in marine environments. *PLOS Climate* 2, e0000148. 10.1371/journal.pclm.0000148.
29. Agency, E.E., Castellari, S., Zandersen, M., Davis, M., Veerkamp, C., Förster, J., Marttunen, M., Mysiak, J., Vandewalle, M., Medri, S., and Picatoste, J. (2021). Nature-based solutions in Europe policy, knowledge and practice for climate change adaptation and disaster risk reduction (Publications Office). doi/10.2800/919315.
30. European, C., and Directorate-General for, C. (2023). The European Green Deal – Delivering the EU's 2030 climate targets (Publications Office of the European Union). doi/10.2775/783179.
31. European, C., Directorate-General for Maritime, A., Fisheries, Joint Research, C., Borriello, A., Calvo Santos, A., Ghiani, M., Guillén, J., Peralta Baptista, A., Petrucco, G., et al. (2023). The EU blue economy report 2023 (Publications Office of the European Union). doi/10.2771/7151.
32. Mcleod, E., Chmura, G.L., Bouillon, S., Salm, R.V., Björk, M., Duarte, C.M., Lovelock, C.E., Schlesinger, W.H., and Silliman, B.R. (2011). A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO<sub>2</sub>. *Frontiers in Ecology and the Environment* 9, 552-560. 10.1890/110004.
33. Froelich, P.N., Klinkhamer, G.P., Bender, M.L., Luedtke, N.A., Heath, G.R., Cullen, D., Dauphin, P., Hammond, D., Hartman, B., and Maynard, V. (1979). Early oxidation of organic matter in pelagic sediments of the eastern equatorial Atlantic: suboxic

958 diagenesis. *Geochimica et Cosmochimica Acta* 43, 1075-1090. 10.1016/0016-  
959 7037(79)90095-4.

960 34. Canfield, D.E., and Des Marais, D.J. (1991). Aerobic Sulfate Reduction in Microbial  
961 Mats. *Science* 251, 1471-1473. doi:10.1126/science.11538266.

962 35. Santos, I.R., Burdige, D.J., Jennerjahn, T.C., Bouillon, S., Cabral, A., Serrano, O.,  
963 Wernberg, T., Filbee-Dexter, K., Guimond, J.A., and Tamborski, J.J. (2021). The  
964 renaissance of Odum's outwelling hypothesis in 'Blue Carbon' science. *Estuarine,  
965 Coastal and Shelf Science* 255, 107361. 10.1016/j.ecss.2021.107361.

966 36. Duarte, C.M., and Krause-Jensen, D. (2017). Export from Seagrass Meadows  
967 Contributes to Marine Carbon Sequestration. *Frontiers in Marine Science* 4.  
968 10.3389/fmars.2017.00013.

969 37. Rullkötter, J. (2006). Organic Matter: The Driving Force for Early Diagenesis. In *Marine  
970 Geochemistry*, H.D. Schulz, and M. Zabel, eds. (Springer Berlin Heidelberg), pp. 125-  
971 168. 10.1007/3-540-32144-6\_4.

972 38. Akam, S.A., Coffin, R.B., Abdulla, H.A.N., and Lyons, T.W. (2020). Dissolved Inorganic  
973 Carbon Pump in Methane-Charged Shallow Marine Sediments: State of the Art and  
974 New Model Perspectives. *Frontiers in Marine Science* 7. 10.3389/fmars.2020.00206.

975 39. Van Dam, B., Lehmann, N., Zeller, M.A., Neumann, A., Pröfrock, D., Lipka, M.,  
976 Thomas, H., and Böttcher, M.E. (2022). Benthic alkalinity fluxes from coastal sediments  
977 of the Baltic and North seas: comparing approaches and identifying knowledge gaps.  
978 *Biogeosciences* 19, 3775-3789. 10.5194/bg-19-3775-2022.

979 40. Reithmaier, G.M.S., Cabral, A., Akhand, A., Bogard, M.J., Borges, A.V., Bouillon, S.,  
980 Burdige, D.J., Call, M., Chen, N., Chen, X., et al. (2023). Carbonate chemistry and  
981 carbon sequestration driven by inorganic carbon outwelling from mangroves and  
982 saltmarshes. *Nature Communications* 14, 8196. 10.1038/s41467-023-44037-w.

983 41. Mueller, P., Kutzbach, L., Mozdzer, T.J., Jespersen, E., Barber, D.C., and Eller, F.  
984 (2023). Minerogenic salt marshes can function as important inorganic carbon stores.  
985 *Limnology and Oceanography* 68, 942-952. 10.1002/lno.12322.

986 42. Koch, E.W., Sanford, L.P., Chen, S.-N., Shafer, D.J., and Smith, J.M. (2006). Waves  
987 in seagrass systems: review and technical recommendations.

988 43. Mateo, M.A., Romero, J., Pérez, M., Littler, M.M., and Littler, D.S. (1997). Dynamics of  
989 Millenary Organic Deposits Resulting from the Growth of the Mediterranean Seagrass  
990 *Posidonia oceanica*. *Estuarine, Coastal and Shelf Science* 44, 103-110.  
991 10.1006/ecss.1996.0116.

992 44. Duarte, C.M., Sintes, T., and Marbà, N. (2013). Assessing the CO<sub>2</sub> capture potential  
993 of seagrass restoration projects. *Journal of Applied Ecology* 50, 1341-1349.  
994 10.1111/1365-2664.12155.

995 45. Fourqurean, J.W., Duarte, C.M., Kennedy, H., Marbà, N., Holmer, M., Mateo, M.A.,  
996 Apostolaki, E.T., Kendrick, G.A., Krause-Jensen, D., McGlathery, K.J., and Serrano,  
997 O. (2012). Seagrass ecosystems as a globally significant carbon stock. *Nature  
998 Geoscience* 5, 505-509. 10.1038/ngeo1477.

999 46. Chmura, G., Anisfeld, S., Cahoon, D., and Lynch, J. (2003). Global carbon  
1000 sequestration in tidal, saline wetland soils. *Global Biogeochemical Cycles* 17 (4), 1-12.  
1001 10.1029/2002GB001917.

1002 47. Bax, N., Sands, C.J., Gogarty, B., Downey, R.V., Moreau, C.V.E., Moreno, B., Held,  
1003 C., Paulsen, M.L., McGee, J., Haward, M., and Barnes, D.K.A. (2021). Perspective:  
1004 Increasing blue carbon around Antarctica is an ecosystem service of considerable  
1005 societal and economic value worth protecting. *Global Change Biology* 27, 5-12.  
1006 10.1111/gcb.15392.

1007 48. Pessarrodona, A., Franco-Santos, R.M., Wright, L.S., Vanderklift, M.A., Howard, J.,  
1008 Pidgeon, E., Wernberg, T., and Filbee-Dexter, K. (2023). Carbon sequestration and  
1009 climate change mitigation using macroalgae: a state of knowledge review. *Biological  
1010 Reviews* 98, 1945-1971. 10.1111/brv.12990.

1011 49. Hansen, K., Butzeck, C., Eschenbach, A., Gröngröft, A., Jensen, K., and Pfeiffer, E.-  
1012 M. (2017). Factors influencing the organic carbon pools in tidal marsh soils of the Elbe



1013 estuary (Germany). *Journal of Soils and Sediments* 17, 47-60. 10.1007/s11368-016-  
1014 1500-8.

1015 50. Kirwan, M.L., and Megonigal, J.P. (2013). Tidal wetland stability in the face of human  
1016 impacts and sea-level rise. *Nature* 504, 53-60. 10.1038/nature12856.

1017 51. Saintilan, N., Rogers, K., Mazumder, D., and Woodroffe, C. (2013). Allochthonous and  
1018 autochthonous contributions to carbon accumulation and carbon store in southeastern  
1019 Australian coastal wetlands. *Estuarine, Coastal and Shelf Science* 128, 84-92.  
1020 10.1016/j.ecss.2013.05.010.

1021 52. Valéry, L., Bouchard, V., and Lefeuvre, J.-C. (2004). Impact of the invasive native  
1022 species *Elymus athericus* on carbon pools in a salt marsh. *Wetlands* 24, 268-276.  
1023 10.1672/0277-5212(2004)024[0268:IOTINS]2.0.CO;2.

1024 53. UNFCCC (2022). Common Metrics. Electronic file available at  
1025 [https://unfccc.int/process-and-meetings/transparency-and-reporting/methods-for-](https://unfccc.int/process-and-meetings/transparency-and-reporting/methods-for-climate-change-transparency/common-metrics)  
1026 [climate-change-transparency/common-metrics](https://unfccc.int/process-and-meetings/transparency-and-reporting/methods-for-climate-change-transparency/common-metrics). .

1027 54. Bockelmann, F.-D., Puls, W., Kleeberg, U., Müller, D., and Emeis, K.-C. (2018).  
1028 Mapping mud content and median grain-size of North Sea sediments – A geostatistical  
1029 approach. *Marine Geology* 397, 60-71. 10.1016/j.margeo.2017.11.003.

1030 55. Delafontaine, M., Bartholoma, A., Flemming, B., and Kurmis, R. (1996). Volume-  
1031 specific dry POC mass in surficial intertidal sediments: a comparison between biogenic  
1032 muds and adjacent sand flats. *Senckenbergiana maritima* 26, 167-178.

1033 56. Diesing, M., Kroger, S., Parker, R., Jenkins, C., Mason, C., and Weston, K. (2017).  
1034 Predicting the standing stock of organic carbon in surface sediments of the North-West  
1035 European continental shelf. *Biogeochemistry* 135, 183-200. 10.1007/s10533-017-  
1036 0310-4.

1037 57. Berner, R.A. (1981). A new geochemical classification of sedimentary environments.  
1038 *Journal of Sedimentary Research* 51, 359-365. 10.1306/212f7c7f-2b24-11d7-  
1039 8648000102c1865d.

1040 58. Al-Raei, A.M., Bosselmann, K., Böttcher, M.E., Hespenheide, B., and Tauber, F.  
1041 (2009). Seasonal dynamics of microbial sulfate reduction in temperate intertidal surface  
1042 sediments: controls by temperature and organic matter. *Ocean Dynamics* 59, 351-370.  
1043 10.1007/s10236-009-0186-5.

1044 59. Jørgensen, B.B. (1982). Mineralization of organic matter in the sea bed—the role of  
1045 sulphate reduction. *Nature* 296, 643-645. 10.1038/296643a0.

1046 60. Keil, R. (2017). Anthropogenic Forcing of Carbonate and Organic Carbon Preservation  
1047 in Marine Sediments. *Annual Review of Marine Science* 9, 151-172. 10.1146/annurev-  
1048 marine-010816-060724.

1049 61. Böttcher, M.E., Hespenheide, B., Llobet-Brossa, E., Beardsley, C., Larsen, O.,  
1050 Schramm, A., Wieland, A., Böttcher, G., Berninger, U.-G., and Amann, R. (2000). The  
1051 biogeochemistry, stable isotope geochemistry, and microbial community structure of a  
1052 temperate intertidal mudflat: an integrated study. *Continental Shelf Research* 20, 1749-  
1053 1769. 10.1016/S0278-4343(00)00046-7.

1054 62. DeFlaun, M.F., and Mayer, L.M. (1983). Relationships between bacteria and grain  
1055 surfaces in intertidal sediments1. *Limnology and Oceanography* 28, 873-881.  
1056 10.4319/lo.1983.28.5.0873.

1057 63. Llobet-Brossa, E., Rosselló-Mora, R., and Amann, R. (1998). Microbial Community  
1058 Composition of Wadden Sea Sediments as Revealed by Fluorescence In Situ  
1059 Hybridization. *Applied and Environmental Microbiology* 64, 2691-2696.  
1060 doi:10.1128/AEM.64.7.2691-2696.1998.

1061 64. Llobet-Brossa, E., Rabus, R., Böttcher, M.E., Könneke, M., Finke, N., Schramm, A.,  
1062 Meyer, R.L., Grötzschel, S., Rosselló-Mora, R., and Amann, R. (2002). Community  
1063 structure and activity of sulfate-reducing bacteria in an intertidal surface sediment: a  
1064 multi-method approach. *Aquatic Microbial Ecology* 29, 211-226.

1065 65. Tyson, R.V. (2001). Sedimentation rate, dilution, preservation and total organic carbon:  
1066 some results of a modelling study. *Organic Geochemistry* 32, 333-339.  
1067 [https://doi.org/10.1016/S0146-6380\(00\)00161-3](https://doi.org/10.1016/S0146-6380(00)00161-3).

66. de Beer, D., Wenzhöfer, F., Ferdelman, T.G., Boehme, S.E., Huettel, M., van Beusekom, J.E.E., Böttcher, M.E., Musat, N., and Dubilier, N. (2005). Transport and mineralization rates in North Sea sandy intertidal sediments, Sylt-Rømø Basin, Wadden Sea. *Limnology and Oceanography* 50, 113-127. 10.4319/lo.2005.50.1.0113.
67. Van Dam, B., Zeller, M., Lopes, C., Smyth, A., Böttcher, M., Osburn, C., Zimmerman, T., Pröfrock, D., Fourqurean, J., and Thomas, H. (2021). Calcification-driven CO<sub>2</sub> emissions exceed “Blue Carbon” sequestration in a carbonate seagrass meadow. *Science Advances* 7, eabj1372. 10.1126/sciadv.abj1372.
68. Turrell, W.R., Austin, W.E.N., Philbrick, S.P., Tilbrook, C., and Kennedy, H. (2023). Clarifying the role of inorganic carbon in blue carbon policy and practice. *Marine Policy* 157, 105873. <https://doi.org/10.1016/j.marpol.2023.105873>.
69. Asplund, M.E., Bonaglia, S., Boström, C., Dahl, M., Deyanova, D., Gagnon, K., Gullström, M., Holmer, M., and Björk, M. (2022). Methane Emissions From Nordic Seagrass Meadow Sediments. *Frontiers in Marine Science* 8, 811533. 10.3389/fmars.2021.811533.
70. Al-Haj, A.N., and Fulweiler, R.W. (2020). A synthesis of methane emissions from shallow vegetated coastal ecosystems. *Global Change Biology* 26, 2988-3005. <https://doi.org/10.1111/gcb.15046>.
71. Novak, A.B., Pelletier, M.C., Colarusso, P., Simpson, J., Gutierrez, M.N., Arias-Ortiz, A., Charpentier, M., Masque, P., and Vella, P. (2020). Factors Influencing Carbon Stocks and Accumulation Rates in Eelgrass Meadows Across New England, USA. *Estuaries and Coasts* 43, 2076-2091. 10.1007/s12237-020-00754-9.
72. Verduin, J.J., and Backhaus, J.O. (2000). Dynamics of Plant–Flow Interactions for the Seagrass *Amphibolis antarctica*: Field Observations and Model Simulations. *Estuarine, Coastal and Shelf Science* 50, 185-204. 10.1006/ecss.1999.0567.
73. Mueller, P., Ladiges, N., Jack, A., Schmiedl, G., Kutzbach, L., Jensen, K., and Nolte, S. (2019). Assessing the long-term carbon-sequestration potential of the semi-natural salt marshes in the European Wadden Sea. *Ecosphere* 10, e02556. 10.1002/ecs2.2556.
74. Mueller, P., Do, H.T., Jensen, K., and Nolte, S. (2019). Origin of organic carbon in the topsoil of Wadden Sea salt marshes. *Marine Ecology Progress Series* 624, 39-50. 10.3354/meps13009.
75. Lavery, P.S., Mateo, M.A., Serrano, O., and Rozaimi, M. (2013). Variability in the carbon storage of seagrass habitats and its implications for global estimates of blue carbon ecosystem service. *PLoS One* 8, e73748. 10.1371/journal.pone.0073748.
76. Legge, O., Johnson, M., Hicks, N., Jickells, T., Diesing, M., Aldridge, J., Andrews, J., Artioli, Y., Bakker, D.C.E., and Burrows, M.T. (2020). Carbon on the northwest European shelf: Contemporary budget and future influences. *Frontiers in Marine Science* 7, 143. 10.3389/fmars.2020.00143.
77. Leipe, T., Tauber, F., Vallius, H., Virtasalo, J., Uścińowicz, S., Kowalski, N., Hille, S., Lindgren, S., and Myllyvirta, T. (2011). Particulate organic carbon (POC) in surface sediments of the Baltic Sea. *Geo-Marine Letters* 31, 175-188. 10.1007/s00367-010-0223-x.
78. Graves, C.A., Benson, L., Aldridge, J., Austin, W.E.N., Dal Molin, F., Fonseca, V.G., Hicks, N., Hynes, C., Kröger, S., Lamb, P.D., et al. (2022). Sedimentary carbon on the continental shelf: Emerging capabilities and research priorities for Blue Carbon. *Frontiers in Marine Science* 9, 22. 10.3389/fmars.2022.926215.
79. Chen, C.-T.A., and Borges, A.V. (2009). Reconciling opposing views on carbon cycling in the coastal ocean: Continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO<sub>2</sub>. *Deep Sea Research Part II: Topical Studies in Oceanography* 56, 578-590. 10.1016/j.dsr2.2009.01.001.
80. Winogradow, A., and Pempkowiak, J. (2014). Organic carbon burial rates in the Baltic Sea sediments. *Estuarine, Coastal and Shelf Science* 138, 27-36. 10.1016/j.ecss.2013.12.001.
81. Colina Alonso, A., van Maren, D.S., Oost, A.P., Esselink, P., Lepper, R., Kösters, F., Bartholdy, J., Bijleveld, A.I., and Wang, Z.B. (2024). A mud budget of the Wadden Sea

and its implications for sediment management. *Communications Earth & Environment* 5, 153. 10.1038/s43247-024-01315-9.

82. GEBCO (2022). GEBCO 2022 Grid. 10.5285/e0f0bb80-ab44-2739-e053-6c86abc0289c.

83. de Haas, H., Boer, W., and van Weering, T.C.E. (1997). Recent sedimentation and organic carbon burial in a shelf sea: the North Sea. *Marine Geology* 144, 131-146. 10.1016/S0025-3227(97)00082-0.

84. Diesing, M., Thorsnes, T., and Bjarnadóttir, L.R. (2021). Organic carbon densities and accumulation rates in surface sediments of the North Sea and Skagerrak. *Biogeosciences* 18, 2139-2160. 10.5194/bg-18-2139-2021.

85. Scheffold, M.I.E., and Hense, I. (2020). Quantifying contemporary organic carbon stocks of the Baltic Sea ecosystem. *Frontiers in Marine Science* 7, 1052. 10.3389/fmars.2020.571956.

86. Middelburg, J.J., Barranguet, C., Boschker, H.T.S., Herman, P.M.J., Moens, T., and Heip, C.H.R. (2000). The fate of intertidal microphytobenthos carbon: An in situ <sup>13</sup>C-labeling study. *Limnology and Oceanography* 45, 1224-1234. 10.4319/lo.2000.45.6.1224.

87. EU Council Directive, H. (1992). Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna and flora. *Official Journal of the European Union* 206, 50.

88. Laurer W., N.M., Zeiler M. (2014). Erstellung der Karte zur Sedimentverteilung auf dem Meeresboden in der deutschen Nordsee nach der Klassifikation von FIGGE (1981). *Geopotenzial Deutsche Nordsee (Modul B.)*.

89. BSH (2016). Meeresbodensedimente in Maßstab 1:100.00 in der deutschen Ostsee. [https://data.bsh.de/SpatialData/Main/SGE\\_SeabedSediments\\_1\\_to\\_100000/](https://data.bsh.de/SpatialData/Main/SGE_SeabedSediments_1_to_100000/)

90. Granse, D., Wanner, A., Stock, M., Jensen, K., and Mueller, P. (2024). Plant-sediment interactions decouple inorganic from organic carbon stock development in salt marsh soils. *Limnology and Oceanography Letters* n/a. <https://doi.org/10.1002/lol2.10382>.

91. Esselink, P., van Duin, W., Bunje, J., Cremer, J., Folmer, E.O., Frikke, J., Glahn, M., de Groot, A., Hecker, N., Hellwig, U., et al. (2017). Salt marshes. [qsr.waddensea-worldheritage.org/reports/salt-marshes](http://qsr.waddensea-worldheritage.org/reports/salt-marshes).

92. Schulte Ostermann, T., Heuner, M., Fuchs, E., Temmerman, S., Schoutens, K., Bouma, T.J., and Minden, V. (2024). Identifying Key Plant Traits and Ecosystem Properties Affecting Wave Attenuation and the Soil Organic Carbon Content in Tidal Marshes. *Estuaries and Coasts* 47, 144-161. 10.1007/s12237-023-01266-y.

93. Butzeck, C., Eschenbach, A., Gröngroft, A., Hansen, K., Nolte, S., and Jensen, K. (2015). Sediment Deposition and Accretion Rates in Tidal Marshes Are Highly Variable Along Estuarine Salinity and Flooding Gradients. *Estuaries and Coasts* 38, 434-450. 10.1007/s12237-014-9848-8.

94. Mueller, P., Granse, D., Nolte, S., Do, H.T., Weingartner, M., Hoth, S., and Jensen, K. (2017). Top-down control of carbon sequestration: grazing affects microbial structure and function in salt marsh soils. *Ecological Applications* 27, 1435-1450. 10.1002/eap.1534.

95. Elschot, K., Bakker, J.P., Temmerman, S., van de Koppel, J., and Bouma, T.J. (2015). Ecosystem engineering by large grazers enhances carbon stocks in a tidal salt marsh. *Marine Ecology Progress Series* 537, 9-21.

96. Buczek, U., Jurasinski, G., Glatzel, S., and Karstens, S. (2022). Blue Carbon in Coastal Phragmites Wetlands Along the Southern Baltic Sea. *Estuaries and Coasts*. 10.1007/s12237-022-01085-7.

97. Koplin J., P.C., Pogoda B. (2024). Blue-Carbon-Potenziale der deutschen Nord- und Ostsee. Status und Trends vor dem Hintergrund des Klimaschutzes. *BfN-Schriften* 699, 116. 10.19217/skr699.

98. Bell, J.D. (1989). Ecology of fish assemblages and fisheries associated with seagrasses. *Biology of seagrasses*, 565-609.

99. Bertelli, C.M., and Unsworth, R.K.F. (2014). Protecting the hand that feeds us: Seagrass (*Zostera marina*) serves as commercial juvenile fish habitat. *Marine Pollution Bulletin* 83, 425-429. 10.1016/j.marpolbul.2013.08.011.
100. Ford, J.R., Williams, R.J., Fowler, A.M., Cox, D.R., and Suthers, I.M. (2010). Identifying critical estuarine seagrass habitat for settlement of coastally spawned fish. *Marine Ecology Progress Series* 408, 181-193.
101. Heck, K.L., and Thoman, T.A. (1981). Experiments on predator-prey interactions in vegetated aquatic habitats. *Journal of Experimental Marine Biology and Ecology* 53, 125-134. 10.1016/0022-0981(81)90014-9.
102. Heck Jr, K., Hays, G., and Orth, R.J. (2003). Critical evaluation of the nursery role hypothesis for seagrass meadows. *Marine Ecology Progress Series* 253, 123-136.
103. Larkum, A.W., Orth, R.J., and Duarte, C.M. (2006). Seagrasses: biology, ecology and conservation. *Phycologia* 45, 5.
104. Nacken, M., and Reise, K. (2000). Effects of herbivorous birds on intertidal seagrass beds in the northern Wadden Sea. *Helgoland Marine Research* 54, 87-94. 10.1007/s101520050006.
105. Horn, S., Coll, M., Asmus, H., and Dolch, T. (2021). Effects of seagrass recovery on the food web of the northern Wadden Sea.
106. Orth, R.J., Heck, K.L., and van Montfrans, J. (1984). Faunal communities in seagrass beds: A review of the influence of plant structure and prey characteristics on predator-prey relationships. *Estuaries* 7, 339-350. 10.2307/1351618.
107. Kuhwald, K., Schneider von Deimling, J., Schubert, P., and Oppelt, N. (2021). How can Sentinel-2 contribute to seagrass mapping in shallow, turbid Baltic Sea waters? *Remote Sensing in Ecology and Conservation* 8, 328-346. 10.1002/rse2.246.
108. KÜFOG GmbH, and Steuer, J. (2020). Eulitorale Seegrasbestände im niedersächsischen Wattenmeer 2019.
109. Dolch, T., Buschbaum, C., and Reise, K. (2013). Persisting intertidal seagrass beds in the northern Wadden Sea since the 1930s. *Journal of Sea Research* 82, 134-141. 10.1016/j.seares.2012.04.007.
110. Dolch, T., Folmer, E., Frederiksen, M., Herlyn, M., van Katwijk, M., Kolbe, K., Krause-Jensen, D., Schemedes, P., and Westerbeek, E. (2017). Wadden Sea Quality Status Report. Wilhelmshaven: Common Wadden Sea Secretariat.
111. Schubert, P.R., Hukriede, W., Karez, R., and Reusch, T.B.H. (2015). Mapping and modeling eelgrass *Zostera marina* distribution in the western Baltic Sea. *Marine Ecology Progress Series* 522, 79-95. 10.3354/meps11133.
112. Ribaudo, C., Plus, M., Ganthy, F., and Auby, I. (2016). Carbon sequestration loss following *Zostera noltei* decline in the Arcachon Bay (France). *Estuarine, Coastal and Shelf Science* 179, 4-11. 10.1016/j.ecss.2016.01.024.
113. Röhr, M.E., Holmer, M., Baum, J.K., Björk, M., Boyer, K., Chin, D., Chalifour, L., Cimon, S., Cusson, M., and Dahl, M. (2018). Blue carbon storage capacity of temperate eelgrass (*Zostera marina*) meadows. *Global Biogeochemical Cycles* 32, 1457-1475. 10.1029/2018GB005941.
114. Gacia, E., Granata, T.C., and Duarte, C.M. (1999). An approach to measurement of particle flux and sediment retention within seagrass (*Posidonia oceanica*) meadows. *Aquatic Botany* 65, 255-268. 10.1016/S0304-3770(99)00044-3.
115. Jankowska, E., Michel, L.N., Zaborska, A., and Włodarska-Kowalczyk, M. (2016). Sediment carbon sink in low-density temperate eelgrass meadows (Baltic Sea). *Journal of Geophysical Research: Biogeosciences* 121, 2918-2934. 10.1002/2016JG003424.
116. Stevenson, A., Ó Corcora, T.C., Hukriede, W., Schubert, P.R., and Reusch, T.B.H. (2023). Substantial seagrass blue carbon pools in the southwestern Baltic Sea include relics of terrestrial peatlands. *Frontiers in Marine Science* 10. 10.3389/fmars.2023.1266663.
117. Jacob, B., Dolch, T., Wurpts, A., and Staneva, J. (2023). Evaluation of seagrass as a nature-based solution for coastal protection in the German Wadden Sea. *Ocean Dynamics* 73, 699-727. 10.1007/s10236-023-01577-5.



- 1233 118. Dolch, T., Buschbaum, C., and Reise, K. (2020). Seegrass-Monitoring im Schleswig-  
1234 Holsteinischen Wattenmeer 2017-2019.
- 1235 119. Bartsch, I., and Tittley, I. (2004). The rocky intertidal biotopes of Helgoland: present  
1236 and past. *Helgoland Marine Research* 58, 289-302. 10.1007/s10152-004-0194-2.
- 1237 120. Pehlke, C., and Bartsch, I. (2008). Changes in depth distribution and biomass of  
1238 sublittoral seaweeds at Helgoland (North Sea) between 1970 and 2005. *Climate*  
1239 *Research* 37, 135-147. 10.3354/cr00767.
- 1240 121. Johannesson, K., Johansson, D., Larsson, K.H., Huenchunir, C.J., Perus, J., Forslund,  
1241 H., Kautsky, L., and Pereyra, R.T. (2011). FREQUENT CLONALITY IN FUCOIDS  
1242 (FUCUS RADICANS AND FUCUS VESICULOSUS; FUCALES, PHAEOPHYCEAE) IN  
1243 THE BALTIC SEA1. *Journal of Phycology* 47, 990-998. 10.1111/j.1529-  
1244 8817.2011.01032.x.
- 1245 122. Hill, R., Bellgrove, A., Macreadie, P.I., Petrou, K., Beardall, J., Steven, A., and Ralph,  
1246 P.J. (2015). Can macroalgae contribute to blue carbon? An Australian perspective.  
1247 *Limnology and Oceanography* 60, 1689-1706. 10.1002/lno.10128.
- 1248 123. Krause-Jensen, D., and Duarte, C.M. (2016). Substantial role of macroalgae in marine  
1249 carbon sequestration. *Nature Geoscience* 9, 737-742. 10.1038/ngeo2790.
- 1250 124. Krumhansl, K.A., and Scheibling, R.E. (2012). Production and fate of kelp detritus.  
1251 *Marine Ecology Progress Series* 467, 281-302. 10.3354/meps09940.
- 1252 125. Queirós, A.M., Stephens, N., Widdicombe, S., Tait, K., McCoy, S.J., Ingels, J., Rühl,  
1253 S., Airs, R., Beesley, A., and Carnovale, G. (2019). Connected macroalgal-sediment  
1254 systems: blue carbon and food webs in the deep coastal ocean. *Ecological*  
1255 *Monographs* 89, e01366. 10.1002/ecm.1366.
- 1256 126. Gallagher, J.B., Shelamoff, V., and Layton, C. (2022). Seaweed ecosystems may not  
1257 mitigate CO2 emissions. *ICES Journal of Marine Science* 0, 1-8.  
1258 10.1093/icesjms/fsac011.
- 1259 127. Burrows, M.T., Kamenor, N.A., Hughes, D.J., Stahl, H., Howe, J.A., and Tett, P. (2014).  
1260 Assessment of carbon budgets and potential blue carbon stores in Scotland's coastal  
1261 and marine environment.  
1262 [https://www.nature.scot/sites/default/files/Publication%202014%20-%20SNH%20Co](https://www.nature.scot/sites/default/files/Publication%202014%20-%20SNH%20Commissioned%20Report%20761%20-%20Assessment%20of%20carbon%20budgets%20and%20potential%20blue%20carbon%20stores%20in%20Scotland%27s%20coastal%20and%20marine%20environment.pdf)  
1263 [mmissioned%20Report%20761%20-%20Assessment%20of%20carbon%20budgets](https://www.nature.scot/sites/default/files/Publication%202014%20-%20SNH%20Commissioned%20Report%20761%20-%20Assessment%20of%20carbon%20budgets%20and%20potential%20blue%20carbon%20stores%20in%20Scotland%27s%20coastal%20and%20marine%20environment.pdf)  
1264 [%20and%20potential%20blue%20carbon%20stores%20in%20Scotland%27s%20coa](https://www.nature.scot/sites/default/files/Publication%202014%20-%20SNH%20Commissioned%20Report%20761%20-%20Assessment%20of%20carbon%20budgets%20and%20potential%20blue%20carbon%20stores%20in%20Scotland%27s%20coastal%20and%20marine%20environment.pdf)  
1265 [stal%20and%20marine%20environment.pdf](https://www.nature.scot/sites/default/files/Publication%202014%20-%20SNH%20Commissioned%20Report%20761%20-%20Assessment%20of%20carbon%20budgets%20and%20potential%20blue%20carbon%20stores%20in%20Scotland%27s%20coastal%20and%20marine%20environment.pdf).
- 1266 128. Fodrie, F.J., Rodriguez, A.B., Gittman, R.K., Grabowski, J.H., Lindquist, N.L., Peterson,  
1267 C.H., Piehler, M.F., and Ridge, J.T. (2017). Oyster reefs as carbon sources and sinks.  
1268 *Proceedings of the Royal Society B: Biological Sciences* 284, 20170891.  
1269 10.1098/rspb.2017.0891.
- 1270 129. Lee, H.Z.L., Davies, I.M., Baxter, J.M., Diele, K., and Sanderson, W.G. (2023). A blue  
1271 carbon model for the European flat oyster (*Ostrea edulis*) and its application in  
1272 environmental restoration. *Aquatic Conservation: Marine and Freshwater Ecosystems*  
1273 *n/a*. 10.1002/aqc.4030.
- 1274 130. Burrows, M., Hughes, D., Austin, W., Smeaton, C., Hicks, N., Howe, J., Allen, C.,  
1275 Taylor, P., and Vare, L. (2017). Assessment of Blue Carbon Resources in Scotland's  
1276 Inshore Marine Protected Area Network. [http://www.snh.gov.uk/publications-data-and-](http://www.snh.gov.uk/publications-data-and-research/publications/search-the-catalogue/publication-detail/?id=2453)  
1277 [research/publications/search-the-catalogue/publication-detail/?id=2453](http://www.snh.gov.uk/publications-data-and-research/publications/search-the-catalogue/publication-detail/?id=2453).
- 1278 131. Sander, L., Hass, H.C., Michaelis, R., Groß, C., Hausen, T., and Pogoda, B. (2021).  
1279 The late Holocene demise of a sublittoral oyster bed in the North Sea. *PLOS ONE* 16,  
1280 e0242208. 10.1371/journal.pone.0242208.
- 1281 132. Wallmann, K., Diesing, M., Scholz, F., Rehder, G., Dale, A.W., Fuhr, M., and Suess, E.  
1282 (2022). Erosion of carbonate-bearing sedimentary rocks may close the alkalinity  
1283 budget of the Baltic Sea and support atmospheric CO2 uptake in coastal seas.  
1284 *Frontiers in Marine Science* 9. 10.3389/fmars.2022.968069.
- 1285 133. Sea, M.A., Hillman, J.R., and Thrush, S.F. (2022). The influence of mussel restoration  
1286 on coastal carbon cycling. *Global Change Biology* 28, 5269-5282. 10.1111/gcb.16287.
- 1287 134. Filgueira, R., Byron, C.J., Comeau, L.A., Costa-Pierce, B., Cranford, P.J., Ferreira,  
1288 J.G., Grant, J., Guyondet, T., Jansen, H.M., and Landry, T. (2015). An integrated

- ecosystem approach for assessing the potential role of cultivated bivalve shells as part of the carbon trading system. *Marine Ecology Progress Series* 518, 281-287. 10.3354/meps11048.
135. Boettcher, M., Schenuit, F., and Geden, O. (2023). The formative phase of German carbon dioxide removal policy: Positioning between precaution, pragmatism and innovation. *Energy Research & Social Science* 98, 103018. <https://doi.org/10.1016/j.erss.2023.103018>.
136. Pogoda, B., Rothe, M., Bakker, F., Hauser, S., Hausen, T., Colsohl, B., Krause, J., Prinz, K., Heinicke, K., Dureuil, M., and Pesch, R. (2023). Come, tell me how you live: Habitat suitability analysis for *Ostrea edulis* restoration. *Aquatic Conservation: Marine and Freshwater Ecosystems*
137. Vadrot, A.B.M., Langlet, A., and Tessnow-von Wysocki, I. (2022). Who owns marine biodiversity? Contesting the world order through the 'common heritage of humankind' principle. *Environmental Politics* 31, 226-250. 10.1080/09644016.2021.1911442.
138. O'Riordan, M., McDonagh, J., and Mahon, M. (2016). Local knowledge and environmentality in legitimacy discourses on Irish peatlands regulation. *Land Use Policy* 59, 423-433. <https://doi.org/10.1016/j.landusepol.2016.07.036>.
139. Wolfenden, A., and Penjueli, M. (2023). Blue Economy: Industrialization and Militarization of Oceans? *Development* 66, 40-45. 10.1057/s41301-023-00365-4.
140. Clark, G. (1994). Rescuing the Discourse of Community. *College Composition and Communication* 45, 61-74. 10.2307/358587.
141. Richardson, B.J., and Razzaque, J. (2011). Public participation in environmental decision making (SSRN).
142. Crew, E.C. (2023). Konsultationsprozess für einen Nationalpark Ostsee. [https://www.schleswig-holstein.de/DE/landesregierung/themen/kueste-wasser-meer/konsultationsprozess-ostsee/downloads/downloadsKonsultationsprozess/download\\_abschlussbericht\\_ecc.pdf?blob=publicationFile&v=2](https://www.schleswig-holstein.de/DE/landesregierung/themen/kueste-wasser-meer/konsultationsprozess-ostsee/downloads/downloadsKonsultationsprozess/download_abschlussbericht_ecc.pdf?blob=publicationFile&v=2).
143. Fink, M., and Ratter, B. (2024). Blurring societal acceptance by lack of knowledge—insights from a German coastal population study on blue carbon. *Frontiers in Climate* 6. 10.3389/fclim.2024.1283712.
144. Aubert, G., Christopoulou, I. (2022). Climate mitigation and adaptation benefits. Institute for European Environmental Policy. (2022). [https://ieep.eu/wp-content/uploads/2022/12/Summary-of-policy-brief-on-the-socio-economic-benefits-of-restoration\\_english.pdf](https://ieep.eu/wp-content/uploads/2022/12/Summary-of-policy-brief-on-the-socio-economic-benefits-of-restoration_english.pdf).
145. Rupprecht, F., Reichert, G., Merling, B., Oltmanns, B. (2023). Renaturierung von Salzwiesen im Nationalpark Niedersächsisches Wattenmeer. *Nationalparkverwaltung Niedersächsisches Wattenmeer*, 54.
146. Poffenbarger, H.J., Needelman, B.A., and Megonigal, J.P. (2011). Salinity Influence on Methane Emissions from Tidal Marshes. *Wetlands* 31, 831-842. 10.1007/s13157-011-0197-0.
147. Koppenaar, E.C., Esselink, P., van Duin, W.E., and Bakker, J.P. (2022). Temporal and Spatial Accretion Patterns and the Impact of Livestock Grazing in a Restored Coastal Salt Marsh. *Estuaries and Coasts* 45, 510-522. 10.1007/s12237-021-00963-w.
148. Hartmann, K., and Stock, M. (2019). Long-term change in habitat and vegetation in an ungrazed, estuarine salt marsh: Man-made foreland compared to young marsh development. *Estuarine, Coastal and Shelf Science* 227, 106348. 10.1016/j.ecss.2019.106348.
149. Martens, M., Müller, P., Gösele, C., Jensen, K., and Schuchardt, B. (2021). Blue Carbon im Nationalpark Niedersächsisches Wattenmeer. *Grundlagen für eine Prüfung und Bewertung möglicher Maßnahmen zur Förderung der Kohlenstofffixierung in Salzwiesen*.
150. Kroeger, K.D., Crooks, S., Moseman-Valtierra, S., and Tang, J. (2017). Restoring tides to reduce methane emissions in impounded wetlands: A new and potent Blue Carbon climate change intervention. *Scientific Reports* 7, 11914. 10.1038/s41598-017-12138-4.

- 1345 151. Bax, V., van de Lageweg, W.I., Terpstra, T., Buijs, J.-M., de Reus, K., de Groot, F., van  
1346 Schaik, R., Habte, M.A., Schram, J., and Hoogenboom, T. (2023). The impact of  
1347 coastal realignment on the availability of ecosystem services: gains, losses and trade-  
1348 offs from a local community perspective. *Journal of Environmental Management* 345,  
1349 118675. 10.1016/j.jenvman.2023.118675.
- 1350 152. Möller, I., Kudella, M., Rupprecht, F., Spencer, T., Paul, M., van Wesenbeeck, B.K.,  
1351 Wolters, G., Jensen, K., Bouma, T.J., Miranda-Lange, M., and Schimmels, S. (2014).  
1352 Wave attenuation over coastal salt marshes under storm surge conditions. *Nature*  
1353 *Geoscience* 7, 727-731. 10.1038/ngeo2251.
- 1354 153. Doolan, G., and Hynes, S. (2023). Ecosystem Service Valuation of Blue Carbon  
1355 Habitats: A Review for Saltmarshes and Seagrasses. *Journal of Ocean and Coastal*  
1356 *Economics* 10, 2.
- 1357 154. Gutiérrez, J., Jones, C., Byers, J., Arkema, K., Berkenbusch, K., Commito, J., Duarte,  
1358 C., Hacker, S., Lambrinos, J., and Hendriks, I. (2011). 7.04—Physical ecosystem  
1359 engineers and the functioning of estuaries and coasts. *Treatise on estuarine and*  
1360 *coastal science*, 53-81.
- 1361 155. Smith, P., Arneth, A., Barnes, D.K.A., Ichii, K., Marquet, P.A., Popp, A., Pörtner, H.-O.,  
1362 Rogers, A.D., Scholes, R.J., Strassburg, B., et al. (2022). How do we best synergize  
1363 climate mitigation actions to co-benefit biodiversity? *Global Change Biology* 28, 2555-  
1364 2577. 10.1111/gcb.16056.
- 1365 156. Kiesel, J., Honsel, L.E., Lorenz, M., Gräwe, U., and Vafeidis, A.T. (2023). Raising dikes  
1366 and managed realignment may be insufficient for maintaining current flood risk along  
1367 the German Baltic Sea coast. *Communications Earth & Environment* 4, 433.  
1368 10.1038/s43247-023-01100-0.
- 1369 157. Vogelsang, L.G., Weikard, H.-P., van Loon-Steensma, J.M., and Bednar-Friedl, B.  
1370 (2023). Assessing the cost-effectiveness of Nature-based Solutions under climate  
1371 change uncertainty and learning. *Water Resources and Economics* 43, 100224.  
1372 10.1016/j.wre.2023.100224.  
1373

1374 S1: Table of ongoing (Blue Carbon related) projects in Germany

1375

Habitat/ Topic	Project name	Info/ Funding	Link
Non-vegetated marine sediments	APOC	Anthropogenic influences on the sequestration of carbon in North Sea sediments; German Federal Ministry of Education and Research (BMBF)	<a href="http://www.apoc-project.de">www.apoc-project.de</a>
	CARBOSTORE	Stability and vulnerability of different carbon reservoirs in the North Sea and Baltic Sea, relevant pathways of carbon storage and the prediction of their future development under different scenarios of climate change and anthropogenic changes; German Federal Ministry of Education and Research (BMBF)	<a href="http://www.carbostore.de">www.carbostore.de</a>
Coastal marshes	sea4soCiety	Innovative approaches to improving the carbon storage potential of vegetated coastal ecosystems; DAM Mission CDRmare; German Federal Ministry of Education and Research (BMBF)	<a href="http://www.sea4society.cdrmare.de">www.sea4society.cdrmare.de</a>
	GREENTRIALS	Greenhouse-gas fluxes in Wadden Sea Tidal Marshes – A Trilateral (NL, DE, DK) Assessment of Natural Climate Solutions; Bauer Hollmann-Stiftung	<a href="http://www.biologie.uni-hamburg.de/forschung/oekologie-biologische-ressourcen/angpfloek/drittmittelprojekte/greentrials.html">www.biologie.uni-hamburg.de/forschung/oekologie-biologische-ressourcen/angpfloek/drittmittelprojekte/greentrials.html</a>
	DFG RTG2530	Fill existing knowledge gaps on biota-mediated effects on estuarine C cycling under current conditions and with respect to global change scenarios; German Research Foundation	<a href="http://www.biologie.uni-hamburg.de/en/forschung/grk2530/overview.html">www.biologie.uni-hamburg.de/en/forschung/grk2530/overview.html</a>
Seagrass meadows	SEASTORE	Investigating and developing restoration guidelines for seagrass meadows in the southern Baltic Sea coast; German Federal Ministry of Education and Research (BMBF)	<a href="http://www.seegrasswiesen.de">www.seegrasswiesen.de</a>
	sea4soCiety	Innovative approaches to improving the carbon storage potential of vegetated coastal ecosystems; DAM Mission CDRmare; German Federal Ministry of Education and Research (BMBF)	<a href="http://www.sea4society.cdrmare.de">www.sea4society.cdrmare.de</a>
Non-classical Blue Carbon ecosystems	DEFINE II	Regional influence of biogenic reefs on the carbon cycle and the surrounding ecosystem in the German North Sea; German Federal Agency for Nature Conservation (BfN)	<a href="http://www.awi.de/forschung/biowissenschaften/oekologie-der-schelfmeere/schwerpunkte/europaeische-auster/define.html">www.awi.de/forschung/biowissenschaften/oekologie-der-schelfmeere/schwerpunkte/europaeische-auster/define.html</a>
	STATUS	Investigation of three main topics: 1) Expansion and assessment of biogenic reefs in the Baltic Sea, 2) Measurement of their carbon storage potential using selected examples, 3) Modelling of the CO2 balance under current and future conditions for the entire area of the German Baltic Sea; German Federal Agency for Nature Conservation (BfN)	<a href="http://www.io-warnemuende.de/projekt/350/status.html">www.io-warnemuende.de/projekt/350/status.html</a>
	ARKOBI	Investigation of the carbon sink capacity of the Iceland mussel Arctica islandica, its associated biotopes and the influence of climate change and anthropogenic utilisation on their biodiversity and the associated carbon storage capacity; German Federal Agency for Nature Conservation (BfN)	<a href="http://www.io-warnemuende.de/projekt/326/arkobi.html">www.io-warnemuende.de/projekt/326/arkobi.html</a>
Storage enhancement through management measures	MGF Nordsee & MGF Ostsee	Influence of anthropogenic activities (mobile bottom-contact fishing) on benthic marine habitats in the German nature conservation areas; DAM Mission Sustainmare; German Federal Ministry of Education and Research (BMBF)	<a href="http://www.mgf-nordsee.de">www.mgf-nordsee.de</a> <a href="https://www.sustainmare.de/104235/index.php.de">https://www.sustainmare.de/104235/index.php.de</a>
	GREENTRIALS	Assessment of livestock management and tidal rewetting on coastal marshes; Bauer Hollmann-Stiftung	<a href="http://www.biologie.uni-hamburg.de/forschung/oekologie-biologische-ressourcen/angpfloek/drittmittelprojekte/greentrials.html">www.biologie.uni-hamburg.de/forschung/oekologie-biologische-ressourcen/angpfloek/drittmittelprojekte/greentrials.html</a>

1376